

Regulation of Kinin Receptor Expression in Heart Failure with a Focus on Vascular Endothelium

Inka Liesmaa

**University of Helsinki
Wihuri Research Institute
Helsinki, Finland**

Academic Dissertation

To be publicly discussed with the permission of the Faculty of Medicine, University of Helsinki,
in auditorium 1, Meilahti Hospital, Haartmaninkatu 4,
on February 12th , 2010, at 12 noon.

Helsinki 2010

Supervisors

Docent Ken A. Lindstedt, PhD
Wihuri Research Institute,
Helsinki, Finland

Docent Jorma O. Kokkonen, MD, PhD
Department of Medicine
Central Finland Central Hospital
Jyväskylä, Finland

Reviewers

Professor (Emeritus) Heikki Vapaatalo, MD, PhD
Institute of Biomedicine,
University of Helsinki
Helsinki, Finland

Docent Risto Kerkelä, MD, PhD
Department of Pharmacology and Toxicology
Faculty of Medicine,
University of Oulu
Oulu, Finland

Opponent

Docent Ilkka Tikkanen, MD, PhD
Department of Medicine
Helsinki University Central Hospital
Helsinki, Finland

ISBN 978-952-92-6782-8 (paperback)

ISBN 978-952-10-6051-9 (PDF)

<http://ethesis.helsinki.fi>

Yliopistopaino
Helsinki 2010

Contents

LIST OF ORIGINAL PUBLICATIONS	8
MAIN ABBREVIATIONS	9
ABSTRACT	11
REVIEW OF THE LITERATURE.....	13
1. Heart Failure	13
1.1 Heart Failure – General.....	13
1.2 Pathogenesis of Heart Failure	14
1.3 Remodeling → Heart Failure	15
1.4 Cardiotoxic Pathways	16
1.4.1 Renin-Angiotensin System	16
1.4.2 Endothelins	16
1.4.3 Cytokines	17
1.4.4 Adrenergic System.....	17
1.5 Cardioprotective Pathways	18
1.5.1 Natriuretic Pathway	18
1.5.2 Adrenomedullin	19
1.5.3 Erythropoietin (EPO).....	19
1.5.4 Other Cardioprotective Proteins	19
2. The Kininogen-Kallikrein-Kinin Pathway.....	20
2.1 History	20
2.2 Kininogens.....	21
2.3 Kallikreins.....	21
2.4 Bradykinin and Kallidin.....	23
2.5 Kinin-degrading Enzymes	24
2.5.1 Aminoterminal-end-cleaving Enzymes.....	25
2.5.2 Carboxyterminal-end-cleaving Enzymes.....	25
3. Kinin Receptors.....	26
3.1 BK-2 Receptors.....	27
3.1.1 History	27
3.1.2 Distribution and signaling of BK-2 Receptors	27
3.1.3 Physiological Effects of BK-2Rs	28
3.1.4 Induction of BK-2 Receptors	29
3.1.4.1 Inflammation	29
3.1.4.2 ACE	30
3.1.5 Inhibition of BK-2 Receptors	31
3.1.6 Polymorphisms	32
3.1.7 Second Messengers of BK-2 Receptors.....	33

3.2 BK-1 Receptors.....	35
3.2.1 History	35
3.2.2 Distribution and Physiological Effects of BK-1 Receptors	35
3.2.3 Induction of BK-1 Receptors	37
3.2.3.1 Inflammation	37
3.2.3.2 Allergy and Pain	38
3.2.3.3 Diabetes	38
3.2.3.4 Other Inducing Factors	38
3.2.4 Inhibition of BK-1 Receptors	39
3.2.5 Second Messengers of BK-1 Receptors.....	39
3.3 Knock-out Models	40
3.3.1 BK-2 Receptor Models.....	40
3.3.2 BK-1 Receptor Models.....	41
3.3.3 Double KO.....	41
4. Smooth Muscle Cells.....	42
5. Angiogenesis	43
6. Endothelial Dysfunction	44
6.1 Endothelium General	44
6.2 Endothelium -derived Vasoactive Substances.....	45
6.2.1 Nitric Oxide	45
6.2.2 EDHF.....	46
6.2.3 PGI ₂	46
6.2.4 PGE ₂	47
6.3 Endothelial Dysfunction	47
6.3.1 Hypoxia	47
6.3.2 Other Factors Contributing to Endothelial Dysfunction	48
6.3.3 Endothelial Dysfunction in Heart Failure	49
7. Treatment.....	50
7.1 Statins	50
7.2 ACE and AT-Receptor Inhibitors	52
7.3 Anti-inflammatory Agents.....	53
7.3.1 COX -Inhibitors.....	53
8. BK- Receptors in Heart Failure.....	55
8.1 Lessons from Human Studies	55
8.1.1 BK-2 Receptors	55
8.1.2 BK-1 Receptors	57
8.2 Lessons from Animal Studies	58
8.2.1 BK-2 Receptors	58
8.2.2 BK-1 Receptors	59
9. Aims of the Study.....	59

10. Materials and Methods	60
Human Heart Samples (I, IV)	60
RNA isolation and PCR (I-IV)	61
Protein Extraction and Western Blotting (I-IV)	61
Histological and Immunocytochemical Staining	62
Culture of Human Coronary Arterial Endothelial Cells	63
Isolation and Culture of Rat Coronary Artery Microvascular Endothelial Cells	63
Measurement of NO in Human Coronary Arterial Endothelial Cells	63
cGMP Immunoassay	64
Wound-healing Assay	64
- / + 9 Polymorphism in Exon 1 of the BK-2 Receptor Gene	64
-58 T/C BK-2 Receptor Promoter Polymorphism by PCR and SSCP	65
11. Statistical analysis	65
12. Results	65
BK-1 Receptors in Human Heart Failure	65
BK-2 Receptor Correlation with Age and Polymorphism	66
Induction of BK -Receptors by Lovastatin in Cultured Endothelial Cells	67
Induction of BK-2 Receptors in Hypoxia	68
13. Discussion	70
BK-1Rs and Inflammation	70
BK-2Rs and Aging	72
BK-2 Receptor Polymorphism	72
BK-2Rs and Statins	73
BK-2Rs and COX-2	74
Endothelial Hypoxia and Preconditioning	75
Clinical Implications	77
14. Summary and Conclusions	78
15. Acknowledgments	80
16. References	82

LIST OF ORIGINAL PUBLICATIONS

- I** Liesmaa I, Kuoppala A, Shiota N, Kokkonen JO, Kostner K, Mäyränpää M, Kovanen P, Lindstedt KA: Increased expression of bradykinin type-1 receptors in endothelium of intramyocardial coronary vessels in human failing hearts: *Am J Physiol Heart Circ Physiol*. 2005;288:H2317-2322
- II** Liesmaa I, Kokkonen JO, Kovanen PT, Lindstedt KA: Lovastatin induces the expression of bradykinin type-2 receptors in cultured human coronary artery endothelial cells. *J Mol Cell Cardiol*. 2007;43:593-600
- III** Liesmaa I, Leskinen HK, Kokkonen JO, Ruskoaho H, Kovanen PT, Lindstedt KA: Hypoxia-induced expression of bradykinin type-2 receptors in endothelial cells triggers NO production, cell migration, and angiogenesis. *J Cell Physiol*. 2009;221:359-66
- IV** Liesmaa I, Shiota N, Kokkonen JO, Kovanen PT, Lindstedt KA: Bradykinin type-2 receptor expression correlates with age and is subjected to transcriptional regulation. Submitted for publication

MAIN ABBREVIATIONS

ACE	=	Angiotensin converting enzyme 1
ACEI	=	Angiotensin converting enzyme inhibitor
ANG II	=	Angiotensin II
APM	=	Aminopeptidase M
ANP	=	Atrial natriuretic peptide
APP	=	Aminopeptidase P
ARB	=	Angiotensin receptor 1 inhibitor
AT	=	Angiotensin
BNP	=	Brain-type natriuretic peptide
cGMP	=	Cyclic guanosine monophosphate
CHD	=	Coronary heart disease
CNP	=	C-type natriuretic peptide
CPM	=	Carboxypeptidase M
CPN	=	Carboxypeptidase N
COX	=	Cyclooxygenase
COXIB	=	Cox-enzyme inhibitor
CTGF	=	Connective tissue factor
EDHF	=	Endothelium derived hyperpolarizing factor
EC	=	Endothelial cell
FGF	=	Fibroblast growth factor
HF	=	Heart failure
HIF	=	Hypoxia inducing factor
HMG-CoA	=	3-hydroxy-3-methylglutaryl-coenzyme A
HMWK	=	High molecular weight kininogen
HTk	=	Human tissue kallikrein
HUVEC	=	Human umbilical vein endothelial cell
IDC	=	Idiopathic dilating cardiomyopathy
IFN- γ	=	Interferon gamma
IL-1 β	=	Interleukin 1- beta
IL-2	=	Interleukin 2
IL-10	=	Interleukin 10
LMWK	=	Low molecular weight kininogen
LPS	=	Lipopolysaccharide
LVH	=	Left ventricular hypertrophy
KO	=	Knockout
MAPK	=	Mitogen-activated protein kinase
MVEC	=	Microvascular endothelial cell
NA	=	Noradrenaline

NEP	=	Neutral endopeptidase
NOS	=	Nitric oxide synthase
NSAID	=	Non-steroidal anti-inflammatory drug
RAS	=	Renin-angiotensin-system
RCMEC	=	Rat coronary microvascular endothelial cell
SMC	=	Smooth muscle cell
SNP	=	Single nucleotide polymorphism
TNF- α	=	Tumor necrosis factor–alpha
VEGF	=	Vascular endothelial growth factor

ABSTRACT

Accumulating evidence, in human and experimental animal models, show that kinins, notably bradykinin (BK) and kallidin, have cardioprotective effects. To these include reduction of left ventricular hypertrophy (LVH) and progression of heart failure. Their effects are mediated through two G protein-coupled receptors- bradykinin type-2 receptor (BK-2R) and bradykinin type-1 receptor (BK-1R). The cardioprotective effects of BK-receptors relate to triggering the production and release of vasodilating nitric oxide (NO) by endothelial cells. They also exert anti-proliferative effects on fibroblasts and anti-hypertrophic effects on myocytes, and thus may play an essential role in the cardioprotective response to myocardial injury. The cardioprotective role of BK-2Rs is widely accepted. The role for BK-1Rs in HF, is based on experimental animal models, where the receptors have been linked to cardioprotective- but also to cardiotoxic -effects. The BK-1Rs are induced under inflammatory and ischemic conditions. The induction has been shown in animal models; no previous reports, concerning BK-1Rs in human heart failure, have been presented.

Previously, it has been shown, that the expression of cardioprotective BK-2Rs were down-regulated in human end-stage heart failure. Present results showed that, in end-stage heart failure patients, the BK-1Rs were up-regulated, suggesting that also BK-1Rs may be involved in the pathogenesis of human heart failure. The receptors were localized mainly in the endothelium of the intramyocardial coronary vessels, and correlated with the increased TNF- α expression in the myocardial coronary vessels. Moreover, in cultured endothelial cells, it was shown, that TNF- α is a potent trigger of BK-1Rs. These results suggest, that cytokines, such as TNF- α may be responsible for the up-regulation of BK-1Rs in human heart failure.

A linear relationship was discovered between BK-2R mRNA and protein expression in normal and failing human left ventricles, implying that the BK-2Rs are regulated on the transcriptional level, at least in human myocardium. The expression of BK-2Rs correlated positively with age in normal and dilated hearts (IDC), though the increase in receptors was clearly lower in dilated hearts. The results suggest, that human hearts adapts to age-related changes, by up-regulating the expression of cardioprotective BK-2Rs. It was also presented, that in the promoter polymorphism -58 T/C, the C-allele was accumulated in cardiomyopathy patients which may partially explain the reduced number of BK-2Rs, in these patients. The BK-receptors seem to contribute to the pathogenesis of human heart failure and they localize mainly in the endothelium of the myocardial coronary vessels.

Statins are cardioprotective drugs, which reduce the level of plasma cholesterol, but also exert several non-cholesterol-dependent effects. The effects of statins were studied in human coronary arterial endothelial cells (hCAEC) and incubation with lovastatin induced both BK-1 and BK-2Rs in a time and concentration-dependent way. Moreover, the induced BK-2Rs were functionally active, thus NO production and cGMP signaling was increased. The induction was abrogated by mevalonate, a direct HMG-CoA metabolite. Lovastatin is known to inhibit Rho activation, and by a selective RhoA kinase inhibitor Y27632, we showed a similar induction of BK-2R expression as with lovastatin. Interestingly a COX-2-inhibitor NS398 inhibited this lovastatin-induction of BK-2Rs, suggesting that COX-2 inhibitors may affect the endothelial BK-2Rs, in a negative fashion.

Hypoxia is a common denominator in HF but also in other cardiovascular diseases. An induction of BK-2Rs in mild hypoxic conditions was shown in cultured hCAECs, which was abolished by a specific BK-2R inhibitor Icatibant. These receptors were functionally active, thus BK increased and Icatibant inhibited the production of NO. In rat myocardium the expression of BK-2R was clearly increased in the endothelium of vessels, forming at the border zone, between the scar tissue and the healthy myocardium. Moreover, in *in vitro* wound-healing assay, endothelial cells were cultured under hypoxic conditions and BK significantly increased the migration of these cells and as Icatibant inhibited it. These results show, that mild hypoxia triggers a temporal expression of functionally active BK-2Rs in human and rat endothelial cells, supporting a role for BK-2Rs, in hypoxia induced angiogenesis.

Our and previous results show, that BK-Rs have an impact on the cardiovascular diseases. In humans, at the end stage of heart failure, the cardioprotective BK-2Rs are down-regulated and BK-1Rs induced. Whether the up-regulation of BK-1Rs, is a compensatory mechanism against the down-regulation of BK-2Rs, or merely reflects the end point of heart failure, remains to be seen. In a clinical point of view, the up-regulation of BK-2Rs, under hypoxic conditions or statin treatment, suggests that, the induction of BK-2Rs is protective in cardiovascular pathologies and those treatments activating BK-2Rs, might give additional tools in treating heart failure.

REVIEW OF THE LITERATURE

1. Heart Failure

1.1 Heart Failure – General

Heart failure (HF) is traditionally viewed as a hemodynamic syndrome characterized by fluid retention, high venous pressure, and low cardiac output. It is usually, but not always, caused by a defect in myocardial contraction – myocardial failure – and it is a principal complication of virtually all forms of heart disease (93). The failing heart presents abnormal growth responses – namely myocyte hypertrophy, progressive remodeling and increased myocardial fibrosis. These structural abnormalities, which now are recognized to play a central role in determining the poor prognosis in this clinical syndrome, result from changes in the size, shape, and molecular composition of both cardiac myocytes and non-myocytes (221).

The major causes of HF are coronary heart disease (CHD), hypertension, and valvular diseases, which account for over 80% of the underlying causes. Other causes include entities such as diseases of the myocardium (myocarditis, cardiomyopathies) and pericardium, diseases of the lung and pulmonary vasculature, arrhythmias, and other diseases such as hyperthyreosis, alcohol abuse and diabetes.

During the last decades the understanding and treatment of heart failure has changed dramatically. Large, randomized clinical trials in the 1980s revealed the poor prognosis of HF patients and changed the view of HF being merely a hemodynamic disorder (220). It has now been realized that changes in the structure and composition of the myocardium play a key role in determining the survival of these patients (220). Patients often remain asymptomatic or minimally symptomatic after initial pump dysfunction has been present for many years. The progression of HF is related to the progression of underlying diseases, neurohumoral abnormalities, and LV remodeling (93).

Heart failure is an increasing cause of morbidity and mortality in the Western world (15). Despite improved treatment and a relative reduction in mortality, the prognosis remains poor. The 5-year survival rate of all HF patients is only 50%, and it is even worse in the more severe cases (15).

The prevalence of heart failure in Europe varies between 0.4% and 2.0% (95). It increases dramatically with age, being 1-2% for persons aged 50-59 years and 6-8% for individuals over 75 years old (93). The mean age of the heart failure population in Europe is 74 years (485). Thus, HF is the leading hospital discharge diagnosis in patients over 65 years old and the prevalence is increasing, which is shown by the 55% increase in the number of hospitalizations

due to HF between 1985 and 1995 (93). Fifteen million individuals in Europe and about five million in America (2.2% of the population) currently have HF and 550 000 patients are diagnosed with HF annually (138;420).

The clinical manifestations of heart failure depend on the rate at which the syndrome develops and specifically on whether sufficient time has elapsed for compensatory mechanisms to develop. Chronic HF can be described as a progressive disease where acute HF episodes with repetitive hospital admissions disrupt the normal progression of the disease.

Since the Framingham study, there have been efforts to improve the prognosis of HF and the treatments have indeed improved. Before ACE inhibitors, the treatment of HF consisted of merely relieving the symptoms with diuretics and nitrates. ACE inhibitors were the first to actually have an impact on the symptoms, but more important on morbidity and mortality. In the last ten years the treatment of heart failure has advanced and whole new categories of drugs have been developed. Despite these new drugs mortality has not diminished dramatically.

1.2 Pathogenesis of Heart Failure

When the heart has difficulty pumping blood at a rate commensurate with the requirements of metabolizing tissues or does so only by elevating the filling pressure, heart failure develops. At the same time the heart switches on a number of adaptive mechanisms for maintaining its pumping function (93). Essential among the adaptive systems are the Frank-Starling mechanism, activation of the neurohumoral systems and myocardial remodeling.

Increased preload helps to sustain the cardiac performance: reduced cardiac output activates neurohumoral responses, leading to vasoconstriction, fluid retention, and cardiac stimulation (93). These adaptations occur rapidly and may be adequate to maintain pumping performance for a short time. Myocardial hypertrophy and progressive dilation (remodeling) of the failing heart account for abnormal proliferative responses and play an important role in long-term adaptation to hemodynamic overload. The structural responses play a crucial role in the poor prognosis of HF, which results from changes in the shape, size, and molecular composition of both cardiac myocytes and other cardiac cells (219).

The development of HF is generally progressive where the initial step challenges cardiac performance- either systolic or diastolic function. Regardless of the initial stimulus, adaptive, compensatory mechanisms are activated in both the heart and the circulatory system. These include activation of the neurohumoral pathway, specifically the renin-angiotensin-aldosterone (RAA) system, and the sympathetic nervous system. This activation leads to production of several neurohormones, which have both hemodynamical

and biological effects on the myocardium and the systemic vascular bed. At first these mechanisms are compensatory, protecting the heart by balancing cardiac output with demands. As the deleterious activation is prolonged the compensatory mechanisms become noxious to the myocardium, leading to alteration of the existing structures (93).

The activation of adaptive mechanisms also leads to the development and progression of LV dysfunction and ultimately to a clinical syndrome of HF (93).

The purpose of these adaptive mechanisms is to sustain cardiac performance in spite of the changed hemodynamic conditions.

1.3 Remodeling → Heart Failure

Left ventricular remodeling is a compensatory process in which mechanical, neuro- hormonal, and possibly genetic factors contribute to alteration of ventricular size, shape, and function (93). It develops over months and years and is crucial to hemodynamic adaptation. It is also irreversible and initially adaptive, thus finally leading to left ventricular hypertrophy (LVH) and clinical HF.

Two major components characterize LVH: hypertrophy of the myocytes and myocardial fibrosis, mainly located in the interstitium but also in the perivascular areas of the myocardium (347;497).

LVH initially compensates for volume or pressure overload by increasing either the size of the left ventricle or its capacity to produce force. Finally, LVH impairs the diastolic function by increasing the stiffness of the left ventricle, working consecutively against the active phase of relaxation and atrial filling. Later, the stiffness produced by LVH begins to interfere with contraction, thereby also affecting the systolic function of the heart.

LVH is associated with ischemia, arrhythmias, and heart failure and represents an independent risk factor of cardiovascular mortality (305). The transition from compensated hypertrophy to manifested HF involves a complex of events at the cellular and molecular level – hypertrophy of cardiac myocytes, changes in gene expression with re-expression of fetal programs and decreased expression of adult programs, changes in the nature and quantity of interstitial matrix, and cellular death by apoptosis and necrosis. HF is progressive, though the velocity of its progression varies depending on factors ranging from underlying cause to ethnical factors (93;141).

It seems very likely that the key to the progressive nature of LVH and HF is the activated signaling peptide systems. Several of these peptides have been shown to induce LVH when infused into animals, and their inhibition has succeeded in improving the otherwise very poor prognosis of HF patients. The most important of these signaling systems include the RA system, the

endothelin system, inflammatory cytokines and the kinin system, as well as the adrenergic system (93).

The balance between cardioprotective elements and components that are deleterious to the myocardium (cardiotoxic) determines the progression of heart failure and its velocity.

1.4 Cardiotoxic Pathways

Cardiotoxic pathways can be defined as signaling pathways that have deleterious effects on the myocardium and the cardiovascular system. Though the presence of these pathways is essential for normal cardiovascular physiology, the transition point from protection to toxicity is define.

1.4.1 Renin-Angiotensin System

The renin-angiotensin system (RAS) plays a prominent role in cardiac remodeling during HF development. Activation of the RA system induces formation of angiotensin II, which promotes myocyte hypertrophy, fibrosis, and chamber dilation in the heart, exerting similar effects to the β -adrenergic system. It also promotes water and sodium retention and exerts electrolytic disturbances (93).

The RAS involves systemic and local components. Although all the components of RAS are present in the heart, not all of them are believed to be synthesized there (341;494). The local RAS seems to be activated differentially depending on the activating signal and regulated independently from the circulating RAS (180;490).

Both local and systemic RA cascades are activated early during the progression of HF, and they are considered the major contributor to most cardiovascular diseases. In fact, the most important drugs for HF are based on blocking the RAS with ACE inhibitors, aldosterone inhibitors, or AT receptor inhibitors. Angiotensin converting enzyme (ACE), which converts Ang I into Ang II, also degrades bradykinin into its metabolites.

1.4.2 Endothelins

Endothelins are the most powerful vasoconstrictors identified. The effects of four different endothelin peptides (ET-1-4) are mediated through two G-protein-coupled receptors (ETA, ETB) that are distributed throughout tissues, notably the vascular endothelium, smooth muscle cells, and cardiomyocytes. In HF patients circulating ET-1 is elevated and correlates with the severity of hemodynamic disturbances and symptoms (300). Though endothelins are known to exert different effects on myocardial contractility in both normal and

failing hearts (272;442), vasoconstriction is the most apparent. However, the endothelin system is not as important as the adrenergic system and RAS for regulation of myocardial performance (312).

1.4.3 Cytokines

There is convincing evidence suggesting that inflammation plays a significant role in the pathogenesis of heart failure. Cytokines are highly potent endogenous peptides produced by a variety of cell types and act as essential mediators and modulators of many biological systems. Tumor necrosis factor (TNF)- α , interleukin (IL)-1 α , IL-1 β , and IL-6 have been classified as proinflammatory cytokines that play important roles in primary host responses and tissue repair. In experimental models, inflammatory cytokines promote left ventricular remodeling and acute reversible contractile dysfunction (299;429). In patients with chronic heart failure, increased plasma levels of proinflammatory cytokines, including TNF- α and IL-6, correlate with both the severity of disease symptoms and clinical outcome (256;456).

The cytokines act in concert with certain cytokine inhibitors and soluble cytokine receptors to regulate the human immune response. Cytokines such as TNF- α have an immediate negative inotropic effect on the myocardium. Some of them are partly responsible for the increased oxygen consumption related to the impaired economy of contraction (189). Some interferons (IFN- γ , IFN- α) also depress the contractility of the myocardium and are involved in the progression of LVH and HF (88;422).

1.4.4 Adrenergic System

One of the most powerful deleterious systems is the sympathetic or adrenergic system. The effects of adrenergic system include activation of α - and β -adrenergic receptors, which are targets of endogenous adrenaline or noradrenaline as well as exogenously-administered β -blockers (243). In chronic HF, increased adrenergic signaling is present and in continuous stimulation, a down-regulation especially of the β -1 adrenergic receptors is present (58). Desensitization of these receptors together with the reduced number of receptors, seen in HF, can be seen as a protective mechanism in the failing myocardium (58).

The adrenergic pathway is activated early in the course of HF and at first serves to compensate for the functional impairment of the myocardium, but ultimately results in deterioration of cardiac function, altered structure, and heart failure. Activation of the adrenergic system induces peripheral vasoconstriction, which in turn increases after-load and oxygen consumption. It causes hypoperfusion and increased anaerobic metabolism in tissues.

Excess noradrenaline increases intracellular calcium, which is directly lethal to cardiomyocytes. Sympathetic activation with excess noradrenalin stimulation promotes eccentric myocyte hypertrophy, enhances apoptosis of cardiomyocytes, and produces detrimental changes in contractile and metabolic proteins via alterations of gene expression in cardiomyocytes (90;135). Activation of the sympathetic system also promotes fibrosis and activates numerous other signaling systems such as the renin-angiotensin (RA) system.

1.5 Cardioprotective Pathways

When cardiotoxic pathways are activated in the myocardium, also opposing systems that protect the myocardium from deleterious signals are activated. To these protecting systems include, for example, the kinin, natriuretic, and adrenomedullin systems as well as the newly discovered urocortin and adiponectin systems. The kinin system is the main focus of this review and it will be closely examined later.

1.5.1 Natriuretic Pathway

In the late 1970s it was discovered that infusing atrial extracts into hypertensive rats lowered blood pressure and resulted in natriuresis (101). The peptide was later isolated and named atrial natriuretic peptide (ANP). Natriuretic peptides are a peptide family constituting ANP, brain natriuretic peptide (BNP), and C-type natriuretic peptide (CNP). ANP and BNP are mostly synthesized in the myocardium and secreted in response to muscle stress. CNP is considered a neurotransmitter in the central nervous system, but is also widely distributed throughout the vasculature and is found in high concentrations in the endothelium where it plays a role in the regulation of local vascular tone (495;506). ANP and BNP are widely used as indicators of myocardial ischemia and heart failure, where they can be used as prognostic biomarkers (320). The effects of natriuretic peptides are mediated through three receptors, namely NPR-A, NPR-B, and NPR-C. ANP and BNP bind preferentially to NPR-A and CNP prefers NPR-B. All three peptides bind to NPR-C, which seems to clear the natriuretic peptides from the circulation.

The peptides induce diuresis and natriuresis and also regulate blood pressure mainly via vasodilation by inhibiting the synthesis and secretion of aldosterone and renin. They inhibit vascular SMC proliferation and hypertrophy of cardiac myocytes and regulate systemic vascular permeability (203). Peptides act as circulating hormones, but also as autocrine and paracrine factors. In heart failure, the levels of ANP and BNP in plasma are increased and they are considered to compensate for heart failure (267).

1.5.2 Adrenomedullin

Adrenomedullin is a potent vasodilating peptide that was originally isolated from pheochromocytoma tissue (227). It is secreted by many cardiovascular tissues and cells including the myocardium, vascular endothelium and vascular smooth muscle cells (191). Adrenomedullin has similar effects as natriuretic peptides thus, it promotes vasodilation, natriuresis, diuresis, angiogenesis, antifibrotic effects, and anti-oxidative actions (227;324). Plasma adrenomedullin levels are increased in patients with various cardiovascular diseases, including hypertension, heart failure, and renal failure. Adrenomedullin has also showed to have an impact on the cardiac contractility and depending on the downstream signaling cascades, it either decreases or increases the contractility of cardiomyocytes (57;253;430). Adrenomedullin as well as natriuretic peptides are mainly degraded in the human heart by neural endopeptidase (NEP).

1.5.3. Erythropoietin (EPO)

In response to ischemia, the cells upregulate erythropoietin through increased expression of hypoxia-inducible factor 1 α (HIF-1 α). Patients, suffering from chronic heart failure, are often anemic and recombinant-erythropoietin (EPO) has been successfully used in the treatment of HF patients (414).

Increased Hb in HF patients improves cardiac function, but EPO seems to exert also some non-hematopoietic effects. Erythropoietin has shown to have anti-apoptotic and antioxidative effects as well as angiogenesis promoting effects (373;418).

Recent studies showed that EPO treatment reduced the infarct size (both brain and heart infarction) in ischemia/reperfusion experiments. *In vivo* and *in vitro* studies reported reduced apoptosis in myocardial cells and normalized hemodynamic functions after reperfusion in rodents. Although the mechanisms of EPO in preserving cardiac functions are mainly unknown, the anti-apoptotic role of EPO has been speculated (69;463).

1.5.4 Other Cardioprotective Proteins

Urocortins (UCN) represent a recently discovered family of cardiac peptides. The name applies to three peptides (urocortin I-III) belonging to the corticotrophin-releasing-factor (CRF) super family. Three urocortins have so far been identified, and they exert their effects through two G-protein coupled receptors (269). Urocortin-I has shown to have potent coronary vasodilator and cardiac inotropic effects. The effects resemble the effects of natriuretic peptides, and in animal models Urocortin I stimulates ANP and BNP secretion from cardiomyocytes (113).

Adiponectin is a protein secreted by adipocytes that has also been

connected to cardioprotective systems (340). It exhibits favorable effects on atherogenesis, endothelial function, and vascular remodeling by modulating the signaling cascades in vascular cells. Recent findings have shown that it can directly affect the signaling of cardiac cells and is beneficial in cardiac remodeling and also in acute cardiac injuries such as myocardial infarction (190).

Cardiotrophin-1 (CT-1) is a cytokine, also shown to have cardioprotective effects. The plasma level of CT-1 is increased in HF and it seems to have diagnostic and prognostic value in heart failure. The level of CT-1 is increased in hypertension, valvular diseases, hypoxia, and myocardial infarction and it stimulates cardiomyocyte proliferation. CT-1 has also shown to exert protective effects against ischemia/reperfusion injury, infused both prior to ischemia and at reperfusion. The results suggest that CT-1 has a protective role in myocardial diseases and it may play a role in decreasing myocardial damage (66;258).

On the basis of these findings, it has been hypothesized that a balance exists between harmful signaling systems and cardioprotective systems, which stabilizes the homeostasis of the normal heart. This equilibrium is disturbed in the beginning of the pathogenesis of HF, resulting in the onset of LVH, accompanied by remodeling and followed by impaired function of the heart, finally leading to heart failure.

2. The Kininogen-Kallikrein-Kinin Pathway

2.1 History

Kininogens are defined as circulating proteins that contain BK or lys-BK. These peptides and their active metabolites assume particular importance as natural ligands of two BK receptors (BK-1R and BK-2R). Interest in the kinin system originates from experiments done a hundred years ago by two surgeons who observed that an alcohol-insoluble fraction of human urine caused hypotension when injected into dogs (480). In 1937 Dr Werle discovered that kallikrein is a proteolytic enzyme that liberates the active polypeptide, kallidin, from the plasma protein kininogen (480). He also noticed the degradation of kinins and identified them as peptides (481).

In the late 1940s Rocha e Silva discovered that trypsin, when incubated with blood, released a substance that contracted the ileum of a guinea pig, and the substance was named bradykinin. He later succeeded in purifying the substance and determined that it was a peptide (378).

2.2 Kininogens

Kininogens are mainly synthesized by hepatocytes, and after posttranslational modifications they are secreted into the circulation (433). Two forms of kininogens have been identified in humans, namely high-molecular-weight kininogen (HMWK) and low-molecular-weight kininogen (LMWK), which differ in size, structure, and function. Their common basic structure – an aminoterminal heavy chain – is linked to varying carboxyterminal light chains, allowing the distinction between low- and high- molecular-weight kininogens (42).

BK and kallidin are formed from either HMWK or LMWK, respectively, by means of the enzymatic action of kallikreins. Several studies have shown that, at least, endothelial cells, platelets and granulocytes have binding sites for HMWK (168;399;465).

The binding of kininogens requires the presence of Zn^{2+} cations, suggesting it is necessary for the expression of the kininogen receptor (510). The affinity of LMWK and HMWK for binding is similar, but the number of binding sites for kininogens varies greatly between cells. Thus, platelets have approximately 1000 binding sites/cell and endothelial cells have between one and ten million binding sites/cell (92). When HMWK is bound to an endothelial cell it activates prekallikrein to release kallikrein, which in turn cleaves BK from HMWK (318).

The kininogen system is involved in inflammation and blood coagulation. In tissue destruction or thrombus formation the contact activation and coagulation activation occur *in vivo*, but the physiological function of the kallikrein system in coagulation has been challenged.




Fig 1. The human kininogen-kallikrein-kinin -system.

2.3 Kallikreins

Kallikreins are serine proteases found in glandular cells, neutrophils, and biological fluids. Kallikreins are divided into two groups: plasma and tissue

kallikreins, which differ in their molecular weight, pI (isoelectric point), substrate specificity, immunological characteristics, type of kinin released, and functional importance. Kallikreins release vasoactive kinin peptides through enzymatic action (321). Enzymes that possess the capacity to release kinins from kininogens are collectively called kininogenases, which include enzymes such as kallikreins, trypsin, neutrophil elastase, and plasmin (236).

Plasma prekallikrein (PPK) is synthesized and secreted by hepatocytes, forms a complex with HMWK, and circulates in the plasma attached to the external surface membrane of neutrophils (182). When the Hageman factor is activated, either by interaction with neutrophil or endothelial membranes or by binding to negatively charged surfaces, it enzymatically converts plasma prekallikrein to plasma kallikrein. A feedback mechanism causes further and rapid activation of the Hageman factor and together with cyclic conversion of PPK to kallikrein, results in local formation of kinins from kallikreins (37).

The major function of plasma kallikrein is to release BK from HMWK by hydrolysis of Lys-Arg and Arg-Ser bonds (217). It also participates in surface-dependent activation of blood clotting, fibrinolysis, regulation of vascular tone, and inflammation. Molecular events in blood clotting involve a number of proenzymes, namely plasma prekallikrein, factor XII (Hageman factor), factor XI, and HMWK, of which HMWK is essential for the coagulation cascade (217). Although LMWK is a poor substrate for plasma kallikrein, it can form BK in the presence of neutrophil elastase (387).

Tissue kallikrein (TK) is a single-chain acidic glycoprotein that differs completely from plasma kallikrein (37). Tissue kallikrein has strong kininase activity, and it is locally synthesized and releases mainly kallidin from both kininogens, though the main substrate is considered to be LMWK (184;273). Tissue kallikrein can also cleave other precursors to release active peptides (37). Tissue kallikrein has been isolated from various tissues, including the heart and arteries (34;302). Unlike with plasma kallikrein, continuous synthesis and secretion of kallikrein can occur in synthesizing tissues, producing kallidin from local and plasma-derived LMWK. This also means that as tissue kallikrein produces kinins continuously, it has an important role in the endothelial function in arteries. It also participates in blood pressure regulation and cardioprotection in early and late ischemic preconditioning (357).

Kallikreins can also activate BK-2Rs independently of kinin generation. This introduces a “shunt” concept into the kallikrein-kinin system in which the normal multistep-cascade to release bradykinin is effectively bypassed as kallikrein directly activates BK-2 receptors (39).

2.4 Bradykinin and Kallidin

Kallidin	Lys ₁ -Arg ₂ -Pro ₃ -Pro ₄ -Gly ₅ -Phe ₆ -Ser ₇ -Pro ₈ -Phe ₉ -Arg ₁₀
Bradykinin	Arg ₁ -Pro ₂ -Pro ₃ -Gly ₄ -Phe ₅ -Ser ₆ -Pro ₇ -Phe ₈ -Arg ₉

Fig 2. The structure of kinins

The two peptides usually referred to as kinins are bradykinin (BK) and kallidin (lys-BK). Their amino acid sequences are shown in **Fig 2**.

In human heart kallidin is rapidly cleaved into BK, and BK is rapidly metabolized into BK-(1-8), BK-(1-7), and BK-(1-5), shown in **Fig 3** (234).

Of the kinin metabolites, des-Arg⁹-BK and des-Arg¹⁰-KD can activate BK-1 receptors, and BK-2 receptors can be activated by both BK and KD. Of the bradykinin degradation products, BK-(1-7) is considered to be inactive and BK-(1-5) to be active in the coagulation cascade (177).




Fig 3. Kinins, their degradation products, and the degrading enzymes in human heart

In 1960 the first amino acid sequence proposed as bradykinin lacked a single proline residue in position 7; the correct sequence was established later that year (50).

The half-life of kinins is very short and they are rapidly degraded into inactive metabolites. In plasma the half-lives of BK and KD are estimated to be < 30 s (144;294). The concentration of BK in the circulation and tissues is very small, and the concentration of kallidin is even smaller (50). The concentrations (25-50 pM) of kinins in tissues are higher than in the circulation, reflecting the fact that tissues produce kinins and also that BK does not participate in systemic blood pressure regulation (314). Bradykinin

can be found in all secretions of the body, i.e. urine, saliva, sweat, and lacrimal fluid, as well as in tissues such as the heart, vasculature, blood, kidney, gastro-intestine track, lungs, skin, brain, and nervous system (72;302;403).

The most important hemodynamic effect of bradykinin is vasodilation, and vascular endothelial cells in the heart are the primary source of BK. Endogenous BK participates in the regulation of coronary vascular tone through BK-2Rs, thus vasodilation can be abolished by a BK-2R antagonist Icatibant. Exogenously-administered BK is also a potent coronary artery vasodilator (84;172;195;348). Vasodilation is partly achieved by stimulating vascular endothelial cells to produce NO but also by activating ECs to produce prostacyclin (PGI₂). The endothelium- derived hyperpolarizing factor (EDHF) is produced by activating BK-2Rs (172;226).

BK-induced vasodilation increases local hyperemia and reduces blood pressure, thus reducing systemic and local vascular resistance (172;397). It also stimulates nociceptive afferent nerves, promotes formation of edema by increasing vascular permeability, and participates in coagulation and fibrinolysis. BK induces antiproliferative effects via BK-2Rs by inhibiting several growth factors that inhibit the hypertrophy of myocytes, thus indicating that an intact kinin system is essential for the regulation of the myocardium (207;293;381).

Kallidin can be detected in the heart and circulation as well as in urine and the kidneys (127). Kallidin concentrations are lower than BK in both plasma (< 1 pM) and tissues (0.5- 4.0 pM), suggesting that the formed kallidin is rapidly degraded into BK (70).

Kinin concentrations are regulated by their formation, but especially by their degradation. There are many enzymes responsible for degradation of kinins, which differ from tissue to tissue, thus kinin profiles vary greatly between tissues. Also both BK-1 and BK-2 receptors, their secondary messenger pathways, and their responses to ligands differ, depending on the cell type and tissue observed. So, the same stimulation can cause entirely different activation in another tissue with the same ligand.

2.5 Kinin-degrading Enzymes




Fig 4. Sites and enzymes which degrade BK

Kinin-degrading enzymes are collectively called kininases. There are four major kininases that cleave kinins from the carboxyterminal end. There are also enzymes that cleave kinins from the aminoterminal end, but their significance seems to be less important – at least in the heart.

In humans, plasma BK is mainly metabolized by three metallopeptidases – ACE, APP, and CPN (41;99). In addition to ACE 1 a homologue of ACE 1 -ACE2-has recently been presented. In this review, ACE refers to ACE 1.

2.5.1 Aminoterminal-end-cleaving Enzymes

Aminopeptidase M (APM) is an enzyme detected in vascular endothelial and epithelial cells in the intestine and renal tubules, and it degrades kallidin into BK (**Fig 4**) (233;337;488). The concentration of kallidin in the human heart is very low, and kallidin is rapidly degraded into BK through the action of APM, likely by fibroblasts (233).

Aminopeptidase P (APP) is a membrane-bound enzyme located in endothelial cells, and particularly high activities have been located in the kidney, heart, and spleen. APP cleaves the first amino acid of BK, converting it into inactive BK-(2-9) (81).

In rats, APP together with ACE are the predominant kininases that participate equally in kinin degradation (108;137). A recent study with rats showed that inhibition of APP potentiated the cardioprotective actions of kinins (489). The importance of APP in humans emerges in the presence of ACE inhibition, when APP represents the main inactivating pathway of BK (99).

2.5.2 Carboxyterminal-end-cleaving Enzymes

The four major enzymes responsible for cleaving kinins from the carboxyterminal end are: angiotensin-converting enzyme 1 (ACE), carboxypeptidase's N and M (CPN, CPM), and neutral endopeptidase (NEP). Based on their enzymatic action, kininases are divided into kininase I and II. To the first belong CPN and CPM and to the latter, ACE and NEP. Kininase I cleaves carboxyterminal Arg⁻⁹ and kininase II cleaves terminal dipeptide Phe⁸-Arg⁹ of kinins. **Fig 4** shows the main cleavage sites of BK.

ACE is mainly localized in the pulmonary vasculature, where it destroys more than 99% of the circulating BK in a single passage through the pulmonary circulation (367). It has generally been thought to be the most important BK-degrading enzyme in the heart and plasma (108;125). Accumulating evidence suggest that in human myocytes, BK is degraded mostly by NEP into inactive BK-(1-7), and only about 10% is degraded by ACE (126;233;354). The activity of kinins seems to be constant in humans in spite of cardiovascular diseases (233). In animals there is some evidence that the level of degrading enzymes

differs in diseased hearts (367). The degradation of kinins in the human heart is due to ACE, NEP, and APM, of which ACE is probably the most important in the endothelium, NEP in the myocardium, and APM in fibroblasts.

Kinins are degraded in plasma mainly by ACE, which cleaves bradykinin into inactive BK-(1-7) and into BK-(1-5), which accompanies the coagulation process.

3. Kinin Receptors




Fig 5. Schematic diagram of a G-protein receptor

Two types of 7-transmembrane G-protein-coupled receptors mediate the cellular effects of kinins, namely BK-1R and BK-2R. A third G-protein-coupled receptor named GPR-100 was recently detected and suspected to be a kinin receptor (49). Bradykinin binds to GPR-100 acts and it resembles BK-2R with a sequence homology of 27%. The receptor has now been confirmed as an insulin-like receptor with relaxin-3 (insulin-like peptide), a ligand (154;265).

The existence of BK receptors was suspected in the early 1970s (100;474;479). The first definition of the receptor was made in 1974 (29). In 1977 two different BK receptors were described in rabbit and rat aortas(369). Later the existence of the two receptors was confirmed, one being constitutively expressed (BK-2R) and the other being inducible (BK-1R) (369). The first findings of BK receptors in the heart were described in the myocardial nerve endings of dogs (425), and almost 20 years later in rat cardiomyocytes (307).

These pharmacologically different receptor subtypes are found interspecies, which explains some of the discrepancies in previous studies done with various animals.

Kinin receptors have been under investigation now for more than 30 years. The scientific interest in kinins arose after the recognition that kinins induce contraction of smooth muscle cells in many vessels, but at the same time

stimulate the production of endothelial vasodilators, thus inducing vasodilation and relating kinins to cardiovascular diseases (85;445). The known positive cardiovascular effects are mainly mediated via BK-2Rs. Substantially less information exists about the functionality and significance of BK-1Rs.

3.1 BK-2 Receptors

3.1.1 History

BK-2Rs were first discovered in cat interstitium, rat stomach and uterus, and rabbit aorta and jugular vein by a in 1975 (28;29;369). The receptor encoding for 366 amino acid residues was cloned from rat uterus SMCs in 1991 (295).

Subsequently, the human receptor was cloned from a fibroblast cell line by PCR and cDNA library screening (183), and the human gene was encoded (128;215). The gene consists of three exons and two introns located in chromosome 14, specifically in 14q32. It codes for a 391 bp amino acid protein with a molecular mass of 41 Da, which has about 80% homology with mouse and rat BK-2 receptors (215;271;359).

3.1.2 Distribution and signaling of BK-2 Receptors

The “native” kinins – BK and kallidin – generated by plasma or tissue kallikreins in all mammalian species have a high affinity for BK-2Rs. None of the fragments of BK retain a significant affinity for BK-2Rs.

The constitutively expressed BK-2Rs have been identified in most tissues and cells: endothelial cells (146), smooth muscle cells (vascular and non-vascular) (36;38;381), and fibroblasts (183). They have also been detected in mesengial cells (13;285), neurons (426), neutrophils (27), and tumor cells, such as adenocarcinoma, lymphoma and hepatoma (491).

BK-2R signaling in the vasculature leads to either vasoconstriction or vasodilation (85;381) and promotes antithrombotic effects (59;407). Depending on the cell type, activation induces proliferative or antiproliferative responses in SMCs and fibroblasts, respectively (160).

Kinin receptors undergo multiple post-translational modifications, including glycosylations and disulfide bridge (cystine) formation, in their extracellular domains. Most of these modifications, including phosphorylation, occur in a ligand-independent manner, although phosphorylation and possibly also acetylation appears to be regulated primarily through receptor activation (355;424).

Of the post-translational modifications in BK-2Rs, glycosylation does not appear to affect ligand-receptor interactions (502). BK-2Rs contain two

cysteine residues in the tail region, which can be modified by palmitoylation, which further modifies receptor coupling, internalization, resensitization, and intracellular trafficking (355;424). Agonist-induced reversible phosphorylation plays an important role in the desensitization of GPCRs, which may also affect endocytosis, recycling, trafficking, and coupling of the receptors.

BK-2 receptors are phosphorylated upon stimulation, leading to receptor desensitization (147;210). Repeated or continuous stimulation leads to internalization of the desensitized receptors. Upon stimulation the receptors are redistributed into plasmalemmal caveolae, and the receptor-caveolae-containing vesicles are thereafter endocytosed and taken into cells. The receptors are recycled back to plasma membranes after a refracting period (43;46)

3.1.3 Physiological Effects of BK-2Rs

In animals and humans the vasodilation under normal conditions is mediated by BK-2Rs (43;46) and thus BK-2Rs mediate hypotensive effects. In the heart, BK-2Rs appear to be antiarrhythmic (262), reduce infarct size, and contribute to preconditioning of the heart against ischemia (61;89;166;504)

BK-2Rs also have an impact on angiogenesis. In animal models receptor activation leads to angiogenesis via several downstream signaling pathways, notably activating eNOS, COX-2, ACE, and VEGF (133;310;415;443).

In the kidneys, BK-2Rs are expressed in all nephron segments, and receptor activation increases renal blood flow, and induces diuresis and natriuresis with no change in the glomerular filtration rate (205;279). Animal studies have shown that BK-2R activation in nephrons promotes antifibrotic effects (389;395). BK-2R activation prevented tubulo-intestinal fibrosis but not perivascular or glomerular fibrosis in hypertensive rats (185). In hypertension it has been shown that the local kinin system plays a significant role in renal protection (11;503). The protective effect is augmented with ACE inhibitors and vasopeptidase (dual ACE+NEP) inhibitors (405). Some conflicting results on BK-2R renoprotection have been published with BK-2R knock-out mice (392).

BK-2Rs affect glucose metabolism by stimulating glucose uptake in the presence of insulin (206;411), and hyperglycemia upregulates BK-2Rs in SMCs (86). The role of BK-2Rs in the development of type-1 diabetic nephropathy has been suggested but is controversial (208). Icatibant reduced glomerular hyper filtration, a phenomenon that is typical in the early phase of diabetes, but another study didn't find any differences in the progression of diabetic nephropathy in BK-2R -antagonized rats (12;453).

Studies of diabetic rats show activation of BK-2Rs and contribution to diabetic nephropathy, where activation of the connective tissue factor,

collagen I, and the transforming growth factor- β (TGF- β) are involved (435). The results have been confirmed in BK-2R $-/-$ diabetic mice in which the renoprotective effect was associated with increased renal expression of BK-1Rs and AT-2Rs (435;436).

In the gastro-intestinal tract, BK and BK-2Rs induce SMCs to either relax or contract (17;145). The same phenomenon is also seen in smooth muscle cells in the uterus, bladder, and bronchi (209;450). The activation of BK-2 receptors has also been linked to inflammation and pain (116;117).

In the respiratory tract, kinins induce bronchoconstriction, and BK and BK-2Rs are associated with the pathogenesis of asthma (116;117).

A potentially important role of BK-2Rs in asthma was demonstrated by a clinical trial in patients where icatibant dose-dependently increased pulmonary function test values. The effect was considered anti-inflammatory rather than bronchodilatory (9).

3.1.4 Induction of BK-2 Receptors

The stimulation of BK-2Rs does not influence the formation of kinins, implicating that there is no feed-back regulation between kinins and BK-2Rs (71;281).

The native ligand of BK-2Rs is bradykinin, but the receptors can also be stimulated by kallidin, kallikreins, or cathepsin G and trypsin (39;179).

Substantially divergent binding affinities have been reported for BK-2Rs in various tissues and preparations. The receptor affinity also depends on receptor coupling, and the phosphorylation state and extent of receptor occupation (43;251;377).

3.1.4.1 Inflammation

Inflammatory cytokines can up-regulate BK-2Rs. In a T24 transitional carcinoma cell line, interferon- γ (IFN- γ) stimulated BK-2R expression (270). Interleukin 1- β (IL1- β) and the tumor necrosis factor- α (TNF- α) induce BK-2Rs through a post-transcriptional mechanism by activating PKA and p38 MAP kinase pathways (173;270;401).

In human airway smooth muscle cells, BK induces the release of IL1- β via cAMP, IL-8, prostaglandin E₂ (PGE₂), COX-2, and IL-6 by a BK-2R dependent pathway (327;344;507). As BK is considered proinflammatory in lungs, BK-2 receptors are possibly involved in the pathogenesis of asthma (30;78).

It has been suggested that BK-2Rs play a role in acute inflammatory response in contrast to BK-1Rs, which account for chronic inflammation (327).

3.1.4.2 ACE




Fig. 6 The renin-angiotensin system. For abbreviations see the text below.

The renin -angiotensin system is a complex system that regulates the cardiovascular physiological processes. The primary effector molecule of this system is the potent vasopressor angiotensin II (Ang II), which has both beneficial and deleterious effects. Angiotensin-converting enzyme (ACE) is an ectoenzyme that generates Ang II by catalyzing its conversion from Ang I. In addition, ACE hydrolyses both bradykinin and kallidin (498). ACE preferentially hydrolyses bradykinin rather than Ang I and is especially abundant in the endothelial cells of lung vessels, but is also expressed in other endothelial cells, some smooth muscle cells, and monocytes (75). After a single passage through the pulmonary circulation, ACE degraded over 95% of administered BK in humans (51). ACE is heterogeneously distributed in the vascular tree, which may be of importance to regional blood flow (282).

ACE inhibitors exert beneficial effects on vascular endothelial function and vascular remodeling, many of which many are considered to be BK-2R-mediated effects (278;396). The beneficial effects are generally attributed to a decrease in Ang II and an accumulation of BK (278;483). ACE is also able to hydrolyse Ang II into Ang (1-7), which potentiates the vasodilating effects of BK, by inhibiting degradation but also by resensitizing BK-2Rs (142;446).

Ang (1-7) is a vasodilating peptide that counteracts the effects of Ang II. It acts by binding to a receptor of its own,- an orphan receptor,-Mas-,or to an AT-1 or AT-2 receptor, and participates in the regulation of cardiac function, coronary perfusion, and endothelial function (268;386). The vasoconstricting effects, described with high Ang (1-7) concentrations, are likely due to stimulation of AT-1R (358;446;459). Ang (1-7) is also formed by degradation of Ang I or II by a novel enzyme, -ACE-2 (115;471).

ACE-2 is a newly discovered enzyme that under normal conditions is located in the myocardium or vascular endothelium. It hydrolyzes Ang II into angiotensin (1-7), angiotensin I into angiotensin (1-9), cleaves kallidin but not BK, and is insensitive to ACE inhibitors (471). It has been suggested that ACE-2 counteracts the function of ACE. ACE-2 - KO mice have severely impaired cardiac function and thinning of the left ventricle wall, but not altered blood pressure or increased cardiac fibrosis. HIF-1- α related genes, induced by hypoxia (PAI-1 and BNIP3), are up-regulated, suggesting a role for ACE-2 in cardiac ischemia (96;115). There is also evidence, that ACE-2 is induced in the myocardium of heart failure patients (62;511).

Over the last decade there have been a number of reports showing that ACE and ACEIs amplify the responses of BK and BK-2Rs. Their cardioprotective effects are partly due to the reduction of BK degradation, but clearly there are other potentiating effects (193;483). ACEIs not only facilitate the accumulation of locally formed BK, they also directly affect BK-2R signaling, resulting in an enhanced response to BK. They stabilize BK-2Rs in a high affinity state, thereby preventing and/or reversing BK-induced sequestration to caveolae before internalization (33). The BK effects have also been suggested to be augmented by ACE-BK-2R cross -talk at the cell membrane (33;309). This could result from a steric relationship between the enzyme and the receptor or from heterodimer formation (283;308).

Both Ang (1-7) and Ang (1-9) induced reactivation of BK-2Rs in Chinese hamster ovary cells (CHO). The effect was attributed to a conformational change in the ACE-BK-2 receptor heterodimer (82;283).

All these results refer to a complex RAA-kinin system interaction, and despite the numerous studies on this subject, its physiological relevance is still not evident.

3.1.5 Inhibition of BK-2 Receptors

The first weak, non-receptor-specific antagonists of the BK-2Rs were generated in 1985 (467). A specific and potent antagonist HOE-140 or icatibant was introduced in 1991 (188) and is still the most commonly used antagonist, despite various chemical subclasses of BK-2R antagonists discovered later (14;383).

BK-2R inhibitors have been developed for clinical use and icatibant is currently used as a drug for treating angioedema. Unfortunately the results have not been as promising as hoped (31;175).

Stimulation of BK-2Rs with agonists reduces the affinity of the receptor towards ligands, thereby requiring higher ligand concentrations for stimulation (107). Long- term (> 24 h) stimulation of BK-2Rs reduces the surface binding sites by means of a post-translational mechanism that reduces receptor protein

synthesis and increase receptor instability (45).

The activation of BK-2Rs involves phosphorylation and desensitization of the receptor and subsequent internalization where the quantity of receptors decreases, which is also observed as the aminoterminal end of the receptor dimer (43;102). Depending on the ligand, recycling of the receptor can take a long time or it can be shortly restored to the plasma membrane. Internalization of BK-2Rs seems to be short-lived, since upon withdrawal of the agonists, a complete restoration of the receptor follows in 30 minutes (486). It also appears that BK-2Rs are almost completely recycled and therefore do not involve permanent down-regulation of the receptors (25).

BK-2R down-regulation is observed in some intense and chronic inflammatory states, but the mechanism is unknown (197). Nitric oxide can also inhibit BK-2R induction by inhibiting coupling with G-proteins (G_i, G_q) via a cGMP-dependent pathway (311).

3.1.6 Polymorphisms

Several polymorphisms of the BK-2R gene have been described. Point mutations occur in both the promoter region and the exons (56). Rieder's group recently sequenced 24 African-American and 23 Caucasian subjects and identified 77 SNPs of BK-2Rs. Twenty-five of them have a frequency greater than 10% and most of the discovered SNPs were located in the intron region. Two relatively highly frequent SNPs, with frequencies of 37% and 40%, were located at 102bp upstream and 277 bp downstream of exon 2. It has not yet been determined if these SNPs alter the function of the BK-2R gene and underlying physiological responses (374).

At least three single-nucleoside polymorphisms (SNP) of the BK-2R gene are known to have an impact on cardiovascular diseases. First, the promoter region -58T/C BK-2R polymorphism has been linked to hypertension in African-American, Chinese and Japanese populations. The C allele is an independent risk factor of essential hypertension (158;478). Fu *et al.* showed that the polymorphism is related to the development of LVH in hypertensive Japanese, but it didn't correlate with hypertension (155). Also the susceptibility to developing a cough with ACE inhibitors is associated with the -58T/C polymorphism (319). It is not clear if the T/ C transition causes an altered expression at the level of BK-2Rs.

Second, a point mutation at +181 C/T in exon 2 determines the Arg → Cys substitution in the extracellular N-terminal domain. The potency of BK (EC_{50}) on BK-2Rs is selectively higher in the presence of the T allele. Furthermore, the T allele has been proposed to be protective during the development of renal failure (196).

And third, the deletion of 9 base pairs (-9/-9) in exon 1 seems to be the most

important polymorphism of the BK-2R gene. A lack of this deletion (+9/+9) is connected to a lower BK-2 receptor expression and is strongly associated with LVH growth response in normotensive men undergoing physical training (196). It also relates to impaired LV mass reduction during antihypertensive treatment (155;174).

3.1.7 Second Messengers of BK-2 Receptors




Fig 7. The intracellular signaling cascades of BK-2Rs
(For abbreviations see the text below)

BK receptors are seven trans-membrane G-protein-coupled receptors GPCR that, when activated transduce the provided information into intracellular second messengers. This process involves the coupling of activated GPCRs to various effector systems via their interaction with heterotrimeric G-proteins. Upon binding, the responses of the intracellular and also the intercellular mechanisms vary according to cell and tissue types as well as activators. The receptors form homo- and heterodimers and oligomers, which also modify their imminent responses. Another modifying factor is the initial stimulus, which can activate several different secondary pathways. The binding of a ligand activates G-proteins, namely $G\alpha_{i2}$ and $G\alpha_q$, which activate phospholipases PLA_2 , PLD , and/or PLC (257). The activation of the phospholipases induces the formation of inositol triphosphate (IP_3) and diacylglycerol (DAG). Upon stimulation, PLA_2 activates arachidonic acid (AA) release and prostaglandin (PGI_2 and PGE_2) production through COX-1 and COX-2 enzymes (160;496). The phospholipases induce an increase in

cytosolic Ca^{2+} and together with IP_3 and DAG they activate the protein kinase (PKC) pathway (43). The PKC pathway in turn activates various kinase and phosphate pathways. The increase of intracellular Ca^{2+} also activates eNOS to synthesize NO (63). eNOS is located in the plasma membrane near BK-2R and caveolin-1 complex. By BK-2R activation, eNOS dephosphorylates and initiates NO production (165;469).

The G-protein $\text{G}\alpha_i$ is important in BK-2R signaling, thus the knockout model ($\text{G}\alpha_{i2-/-}$) displayed a two-fold increase in resting cAMP levels, which seems to be a BK-2R up-regulating factor, at least in fibroblasts (291).

In endothelial cells, BK-2Rs are located near the plasmalemmal caveolae (102). Caveolae are cholesterol- and glycosphingolipid-enriched invaginations of plasma membrane that function to bind, organize, and conserve a variety of signaling molecules (264). Several signaling proteins, including eNOS, interact with caveolin, an oligomeric integral membrane protein that appears to serve as the structural “scaffold” within caveolae (16).

When a ligand activates BK-2Rs, they normally transfer into the caveolae (102;211). The activated receptors are phosphorylated at Ser/Thr residues, located in the C-terminal domains and uncoupled from G-proteins (47;356). The receptors are internalized to an intracellular compartment by endocytosis and relocated in vesicles. By these actions the receptors are desensitized and can not respond to ligands. After internalization, dephosphorylated BK-2Rs can be resensitized and recycled back in to the plasma membrane (43). The cycle from receptor activation to resensitization and the capacity to bind a new ligand takes minutes, thus being a regulative element in BK-2R activation. The complex regulation of receptor sequestration and desensitization contributes to the differential use of the phosphorylation sites in BK-2Rs (46;356)

G-protein-coupled receptors (GPCRs) constitute a large family of proteins that converts external stimuli into intracellular activity. The receptors share a common deduced structure, and through their intracellular domains, they interact with heterotrimeric G-proteins. The activation of GPCRs initiates intracellular second messenger cascades, which ultimately evoke carefully controlled cellular responses (326).

The structures and functions of GPCRs are complex, and the secondary pathways vary from the cell to the participating ligand. Activation of BK-2Rs can induce vasodilatory effects on vessels via three different mechanisms. First, they can activate eNOS to produce NO, second they can activate PLA_2 , which promotes synthesis of prostacyclin. A third important vasodilating agent, the endothelial-derived hyperpolarizing factor (EDHF), is also related to BK-2Rs. The identity of EDHF is unknown, though a metabolite of arachidonic acid, epoxyeicosatrienoic acid (EET), is a strong candidate. EETs are synthesized in the vascular endothelium, they open calcium-activated potassium channels, and they hyperpolarize the membrane of vascular smooth muscle, conducting

relaxation (148). It has been proposed that EDHF consists of numerous endothelium-derived factors, including NO and PGI₂, but also reactive oxygen species, carbon monoxide, adenosine, peptides, and various metabolites of arachidonic acid, which can hyperpolarize the underlying smooth muscle thus initiating relaxation (139).

EDHF plays a significant role in resistant vessels, and it has been suggested as an important factor also in the vasodilation of coronary arteries, human forearm arteries, and renal arteries. It is thought to have relevance in preconditioning of the heart (410), and to contribute to vascular reactions in diabetes and other cardiovascular associated diseases (431;484).

The vasodilating substances NO, PGI₂, and EDHF partly cover each other, thus a lack of NO can be substituted by EDHF or PGI₂ (198;427). In eNOS knockout mice, EDHF completely replaces the lack of vasodilation normally produced by NO (427). Again, increased NO production inhibits the formation of EDHF, suggesting a physiological role for EDHF (224;363).

3.2 BK-1 Receptors

3.2.1 History

Regoli and co-workers also characterized BK-1Rs in the 1970s. The two G-protein receptors (BK-1R and BK-2R) were shown to differ in pharmacological profile and expression patterns (28;369;370). The gene was cloned from human embryonic lung fibroblast and is also located in chromosome 14, specifically 14q32(24). The receptor cDNA encodes the 353 amino acid protein with a molecular mass of approximately 41 kDa (303). The two human BK receptor genes are located consecutively in the same chromosome sharing a 36% genomic sequence homology. Mouse, rat, and rabbit BK-1Rs display functional and structural homology with human receptors, with rabbit receptors showing the greatest homology (24).

3.2.2 Distribution and Physiological Effects of BK-1 Receptors

In humans, BK-1Rs are weakly expressed under physiological conditions, but can be induced under pathological stimuli such as tissue injury or inflammation (67;369). However, constitutive expression of BK-1Rs is detected in the vascular compartment of rabbits, dogs, and cats (32;109).

BK-1 receptors can be detected in blood vessels, both veins and arteries, in cardiac tissue, kidneys, lungs, the central nervous system, ECs, SMCs and, fibroblasts (121;303;328). The distribution of BK-1Rs in vascular ECs and SMCs differs according to the size of the vessel: no BK-1Rs can be detected

in large veins or in the aorta (364).

Kinins able to induce BK-1Rs are des-Arg⁹-BK and des-Arg¹⁰-kallidin, bradykinin or other BK-2R inducers do not activate BK-1Rs (368).

The signaling of BK-1Rs is usually an endothelium-dependent phenomenon and can cause either vasoconstriction or vasodilation, depending on the species, the vessel, and its location (120;297). In most vessels the main effect is dilation, although in rabbit aorta and mesenteric vein as well as in human umbilical artery and vein, activation of BK-1Rs causes vasoconstriction (1;163;254;298;369;370).

Vasodilation is normally related to inflammation and is attributed to activation of eNOS and subsequently prolonged release of NO (109;120). The vasodilatory effects can be produced by prostanoids, mainly by PGI₂ and PGE₂, like in BK-2Rs (297;416). However, there are also conflicting studies where BK-1R-mediated vasorelaxation was not sensitive to cyclo-oxygenase inhibition, suggesting that prostaglandin release was not involved (362;385). This difference may be related to species and regional variation or to the absence of a specific prostanoid receptor subtype (297).

In some vessels, vasodilation mediated by the induction of BK-1Rs is endothelium-independent, which has been attributed to prostaglandin activity (376). BK-1Rs are positively coupled to phospholipase A₂ and induces arachidonic acid release (419). Despite prostaglandin synthesis, functional responses depend upon local PG sensitivity. Stimulation of BK-1Rs provokes PG synthesis and contraction in rabbit SMCs and bovine ECs, the contraction can not be prevented with indomethacin (121;254). PGI₂ and PGE₂ induce vasodilation in rabbit mesenteric and celiac arteries and rat coronary circulation, but fail to relax rabbit aorta, although BK-1R stimulation induces PGI₂ secretion (151;297;416) This indicates that not all endothelium-independent BK-1R-mediated vasodilatory effects are prostaglandin-mediated, despite the fact that prostaglandins are consistently released upon BK-1R stimulation. The prolonged response of activated BK-1Rs is probably related to the lack of desensitization and internalization of the receptors (22;289). Human BK-1Rs are not phosphorylated to any significant degree in either the absence or presence of ligands (44).

Many of the peripheral effects of BK-1R activation are pro-inflammatory: vasodilation, increased blood flow, increased plasma protein extravasation, and activation of leukocyte-EC interactions and leukocyte accumulation (8;296). Due to this, BK-1R studies have mainly focused on the regulation and expression of receptors in inflammation and infections (67;352).

The hypotensive responses mediated by BK-1R activation generally occur following inflammation and they have been extensively revealed (21;297;371).

Beyond the cardiovascular effects, BK-1Rs have been connected to diabetes

(512), chronic pain, and nociception (153) as well as cancer (365;438).

Although BK-1Rs and BK-2Rs seem to couple to similar signal transduction pathways, their signaling patterns differ. Also the impact on the differences in expression and regulation needs to be explored further.

3.2.3. Induction of BK-1 Receptors

Knowledge of the functionality and signaling mechanisms of BK-1Rs is incomplete. These receptors are present in tissues in negligible quantities and they can be induced by various inflammatory signals that produce vasoactive responses. BK-1R up-regulation occurs under the control of specific cytokines released in inflammation or after trauma (297;455). Unlike BK-2Rs, the stimulation of BK-1Rs does not desensitize receptors, indeed BK-2R desensitization can up-regulate BK-1Rs. This finding supports the hypothesis of an integrated system where BK-2Rs mediate acute inflammatory effects and BK-1Rs mediate sustained or chronic inflammatory responses. ACE inhibitors can also directly up-regulate BK-1Rs (202).

3.2.3.1 Inflammation

Induction of BK-1Rs is associated with production of inflammatory mediators, stimulation of inflammatory cells, and activation of several intracellular signaling pathways (351). Studies have shown that infection, with or without bacterial lipopolysaccharides (LPS), induces BK-1Rs and the vasodilation seen in septic infections is, at least partly, mediated by BK-1Rs (104;352).

LPS induces effects that are mediated through inflammatory molecules such as $\text{NF-}\kappa\text{-}\beta$, $\text{IL-1}\beta$, $\text{TNF-}\alpha$, and PAF, which in turn promote the BK-1R up-regulation (346). LPS up-regulates BK-1R expression in rat coronary circulation, also possibly via direct action within the vessel wall (382).

Evidence from human lung fibroblasts suggests that receptor up-regulation involves activation of PKC, MAPK, and $\text{NF-}\kappa\text{-}\beta$ pathways (391;508). $\text{IL-1}\beta$ is the optimal inducer of BK-1Rs in various cell types, and up-regulation modulates the receptors at both the transcriptional and post-transcriptional levels (391;508).

Activation of BK-1Rs reduces renal plasma flow and the glomerular filtration rate (GFR) in rats. This is attenuated when LPS is introduced prior to receptor induction, suggesting that activation of BK-1 receptors may participate in the modification of renal hemodynamics in inflammatory states (394). Over-expression of BK-1Rs renders transgenic-mice more susceptible to endotoxic shock and also exacerbates induced paw edema. The hemodynamic response to intravenously administered BK-1R agonist surprisingly increased mean arterial blood pressure in transgenic mice (329).

These results support a role for BK-1Rs in the modulation of inflammatory responses and hypertensive responses.

3.2.3.2 Allergy and Pain

Huan *et al* showed that allergen-stimulated bronchial hyper-reactivity in rats increased BK-1R expression, which was negated by blocking the receptor (199). Completely opposite results were published a few years later (130). The majority of animal studies support a role for BK-1Rs in lung inflammation, although the results have not been confirmed in human studies (136;372).

There is compelling evidence indicating that BK-1Rs play a key role in persistent inflammatory pain (67). Recent studies suggest a role for BK-1Rs also in neuropathic pain, which can be addressed as chronic and is in general very resistant to traditional medications. In transgenic mice, expressing human BK-1Rs, promising results were demonstrated when treating opioid-resistant, neuropathic pain with a highly human-specific BK-1R antagonist, (NVP-SAA164) (153). Another substance, namely ELN441958, has shown promising results in diminishing carrageenan-induced hyperalgesia in rhesus monkeys (178). Streptozotocin (STZ) -treated mice show characteristics of hyperalgesia, which can be completely blocked by a BK-1R antagonist (157).

3.2.3.3 Diabetes

In several animal studies, BK -1R up-regulation has been linked to diabetes (73;470). Streptozotocin-treated (STZ) diabetic rats showed long-lasting up-regulation of functional BK-1Rs in peripheral tissues, which could be negated with insulin treatment (73;94). A BK-1R antagonist (Leu⁸-des-Arg⁹-BK) was able to prevent insulin-dependent diabetes, presumably via an anti-inflammatory effect, in STZ -treated mice, but a BK-2R antagonist (Icatibant) was not able to do so (513). There are also some data concerning the development of diabetes that suggest that during the progression of diabetes, BK-1Rs are over-expressed and BK-2Rs are down-regulated (68).

A role for BK-1Rs has been suggested in the development of diabetes-associated retinopathies and neuropathies. A clear increase in BK-1R-specific binding sites is associated with an enhancement of des-Arg⁹-BK induced vasodilation in the retinas of STZ-treated rats (2).

3.2.3.4 Other Inducing Factors

Many studies suggest that ACE inhibitors can also directly activate BK-1Rs. Opposite results have been reported suggesting that ACE inhibitor-

induced BK-1R up-regulation might be an indirect effect, resulting from low endogenous kinin generation (201).

Increased NO production detected with ACE inhibitor treatment has been suggested to relate to direct BK-1R stimulation (201). The induction of BK-1Rs requires Zn^{2+} binding, inducing elevated Ca^{2+} levels and subsequent activation of eNOS and NO production, which can be blunted by BK-1R antagonist Ca-EDTA (201).

There are some indirect data from human heart failures suggesting that some of the vasodilatory effects induced by ACE inhibitor treatment are mediated through BK-1Rs (487).

It has been suggested that BK-2Rs and BK1Rs can “cross-talk” thus potentiating or preventing each other. Persistent stimulation or prolonged desensitization of BK-2Rs results in BK-1 receptor up-regulation, suggesting that the receptors can co-operate. BK-2R agonists can also directly activate the $NF\kappa\beta$ pathway, which in turn can activate BK-1Rs (353;390).

A heterologous BK-1-BK-2 receptor-complex has been presented. Human BK receptors were spontaneously heterodimerized and the complex was associated with proteolytic degradation of BK-2Rs, suggesting a regulatory mechanism that participates in the transduction of a signal from BK-2Rs to BK-1Rs (216). This formation of a receptor complex may indicate a cellular adaptation of kinin signaling in sustained inflammation.

3.2.4 Inhibition of BK-1 Receptors

There are very limited data, on the inhibitors that influence BK-1R signaling.

Glucocorticoids and protein synthesis inhibitors have been shown to suppress the up-regulation of BK-1Rs in the rabbit aorta (103).

The number of BK-1Rs is constantly up-regulated in rats exhibiting low glucocorticoid levels, and by providing glucocorticoids the receptor levels are dropped back to normal levels (65).

The tumor suppressor factor, p53, can transcriptionally activate or silence a number of target genes, and it has been shown to inhibit the BK-1R gene (499).

3.2.5 Second Messengers of BK-1 Receptors

On activation, BK-1Rs couple to G-proteins, preferentially to the pertussis toxin insensitive G_q , or G_i , through which it mediates stimulation (22).

The two BK receptors have many similar signaling pathways that involves elevation of intracellular Ca^{2+} , arachidonic acid release, eicosanoid production, and eNOS activation together with NO production (255;457).

3.3 Knock-out Models

3.3.1 BK-2 Receptor Models

A BK-2R knockout (KO) mouse was first generated by Borkowski in 1995. The mouse failed to respond to BK stimulation, such as smooth muscle contraction or afferent nerve stimulation. These mice are phenotypically normal and fertile and they have normal or only slightly elevated blood pressure, though malignant hypertension occurs with excess dietary salt (10;52). They have several alterations contributing to cardiovascular function: enhanced effects on Ang II, distorted renal development with abnormal nephrons and decreased glomerular capillary surface, and decreased renin and COX-2 expression in the kidneys (76;393).

Emanuelli studied the cardiac phenotype of BK-2R knockout mice and reported the development of LVH accompanied by only a moderate rise in blood pressure in these animals by 2 months of age. LVH progressed to fulminant heart failure in six months without an increase in blood pressure, suggesting that remodeling was the primary process unrelated to systemic hypertension (132).

Progressive degenerative and fibrotic changes were observed in the myocardium of $-/-$ mice and similar but less prominent changes were detected in heterozygous BK-2R $+/-$ mice. The knockout animals were also tachycardic, similar to rats with a prenatal blockade of BK-2Rs, suggesting that kinins are also involved in early programming of the cardiovascular phenotype (274). Increased myocardial weight and blood pressure at the base-line has been reported in knockout mice, though others have failed to repeat these results (11;275;448).

Duka *et al.* reported that BK-2R knockout-mice developed LVH and heart failure after myocardial infarction and that remodeling and cardiac dysfunction responded poorly to ACE inhibitor or ARB treatment. The same group later reported that ACE inhibition worsened the cardiac function in mice lacking BK-2Rs. This effect was linked to up-regulation of pro-inflammatory BK-1Rs (122;124).

In BK-2R knockout mice, BK-1Rs are up-regulated and their sensitivity to stimulation is significantly increased, which partially substitutes for the lack of BK-2Rs. Further, stimulation of BK-1Rs with an agonist provokes hypotension and antagonist-induced hypertension, suggesting an antihypertensive effect of BK-1Rs, possibly through inhibition of vasoconstrictors produced through the AA cascade (123;124).

Kakoki *et al.* reported that in the absence of BK-2Rs, Akita diabetic mice show an additive increase in oxidative stress, mitochondrial DNA damage, and other senescence-associated phenomena, implying that increased senescence

and apoptosis in actively dividing cells are related to BK-2Rs. The authors suggest that the increased oxidative stress and ROS are due to decreased intracellular NO production in BK-2R-deficient diabetic mice (212).

The transgenic mice models have further verified the role of BK-2Rs in hypertension. Over-expression of human BK-2Rs in mice induces hypotension, which can be negated by icatibant. Also rats that over-express human tissue kallikrein gene experience hypotension. Induced hypertension with subsequent LVH was significantly reduced in hTK over-expressing rats compared with controls, and icatibant completely inhibited the effects (413).

3.3.2 BK-1 Receptor Models

The BK-1R knockout mouse was generated five years later than the BK-2R KO (352). The lack of BK-1Rs has no effect on blood pressure or cardiac performance in physiological situations.

Induced myocardial infarction in BK-1R $-/-$ mice did not significantly influence myocardial remodeling (493). In another study BK-1R knockout mice showed higher coronary perfusion pressure, indicating higher coronary resistance. eNOS expression was also significantly diminished in the aortas of $-/-$ mice, which could explain their increased coronary resistance (249). BK-1R knockout-mice have a pain-phenotype that implicates a role for this receptor in inflammatory and neuropathic pain. Knockout-mice exhibit reduced responses compared to wild types, in several pain models (143;352).

The absence of BK-1Rs in LPS-induced septic shock is associated with significant hypotension and significantly increased mortality (297).

The relevance of BK-1Rs in LPS-mediated septic shock has been intensively studied in knockout-models and the results have been conflicting: some report hypotension and others hypertension in septic BK-1R knockout-mice (74;352). Experimental differences explain some of these discrepancies, and it has been suggested that the BK-1Rs are first up-regulated in the initial phase of septicemia and are associated with vasoconstriction, implying that the BK1-Rs protect against acute, significant hypotension, thus reducing mortality (406).

Reduced neutrophil migration to inflamed tissues as well as apoptosis of BK-1R expressing neutrophils as a requirement for the resolution of inflammation was reported in mice lacking BK-1Rs (352).

3.3.3 Double KO

Until recently, the close proximity of the genes encoding the two receptors has prevented the generation of double-KO mice. Surprisingly double- KO -mice are normotensive and their cardiovascular morphology is normal, supporting

previous results indicating that under basal conditions, the kinin system does not play a significant role in cardiovascular physiology, at least in mice (74).

Table I. The main (cardiovascular) effects of BK-Rs

	BK-1R	BK-2R	BK1R ref	BK-2R ref
Vasodilation	++ (artery)	++	(103;120;297)	(85;87)
Vasoconstriction	++ (vein)		(1;163;369)	
SMC effects	anti-proliferation	anti-proliferation	(112)	(381)
VSMC contraction	+	++	(109)	(254;255)
Ischemic preconditioning	+	++	(53;77)	(61;89;468;475)
Angiogenesis	+	+	(112;345)	(133;310;443)
Fibroblast	pro-fibrotic	anti-fibrotic	(181)	(389;395;454)
Inflammation	chronic	acute	(68;73;298)	(298)
Leukocyte trafficking			(35;296;352)	
Cytokine activation			(104;172)	(173;270)
Endotoxic shock	hypotension	early hypotension	(103;297;371)	(298)
Pain	chronic	acute	(351;352)	(67;117;351)
Thrombosis		anti-thrombotic		(59;361)
Bronchoconstriction	+	+	(199)	(37)

4. Smooth Muscle Cells

In addition to endothelial cells, the BK -receptors can also be detected in other cells, though it is believed that their major effects are contributed via ECs.

It is known that in smooth muscle cells, BK-2R stimulation causes vasoconstriction and contributes to vascular injuries. In denuded canine saphenous veins, direct induction of BK onto SMCs induced concentration-dependent vasoconstriction, which seemed to involve BK-2Rs and arachidonic acids, implying PGI₂ involvement (287).

In experiments with intact guinea pig aortic rings or arterial myocyte cultures, direct BK-2R stimulation leads to constriction.

When the vascular endothelium is damaged, stimulants can gain direct access to the receptors in the smooth muscle. In these circumstances it is possible that BK-Rs play a role in the regulation of vascular injury.

5. Angiogenesis

Angiogenesis is the process of forming new capillaries from existing blood vessels. It involves several steps that result in the formation of new vessels when the highly regulated balance is in favour of angiogenesis (149).

The pro-angiogenic effect of BK has been demonstrated in both *in vitro* and *in vivo* studies under different experimental settings (79;131;231;317;345). Some data on cardiac angiogenesis are also available, which show favourable effects of kinin receptors on angiogenesis. It was recently shown that angiotensin II induced angiogenesis in hypoxic mouse hearts in a BK-2R- and NO-dependent way that also involved AT-2Rs (322). Interestingly, neither Ang II nor VEGF could induce angiogenesis in BK-2R knockout-mice (322). BK-2Rs transactivated the VEGF receptor (KDR/Flk1) in cardiac capillary ECs, resulting in further eNOS activation (443). Later the results were confirmed with coronary endothelial cells, where BK induced tube formation through a BK-2R-KDR/Flk1-dependent mechanism. BK and VEGF were suggested to have a synergistic effect on angiogenesis (310).

Angiogenesis has also been linked to BK-1Rs (345). The study showed that BK was able to activate BK-1Rs in rabbit corneal capillary endothelial cells by stimulating the eNOS/FGF-2-pathway. These results were confirmed later in a study that showed a BK-1R dependent up-regulation of eNOS, and FGF-2 was involved in microvascular coronary endothelial cell and migration was induced by ACE inhibitors (114).

Tissue kallikrein has been associated with angiogenesis in several transgenic studies.

Transduction of the human tissue kallikrein (hTK) gene into an ischemic myocardium stimulated arteriole and capillary growth, reduced apoptosis, and increased the number of cardiac progenitor cells. A transient increase in TK was detected and the Akt-eNOS-NO-pathways were activated through increased kinin production (133;176). Unfortunately, these studies did not explore the BK receptors responsible.

In another study, hTK gene delivery improved cardiac function and protected against cardiac remodeling in HF by attenuating cardiac hypertrophy, fibrosis, and LV enlargement and by promoting capillary formation. These effects were seemingly connected to kinin-BK-2R activation (6;48;505).

These studies suggest that BK modulates cell-signaling pathways through BK1-and/or BK2-receptors and thus plays a role in regulating angiogenesis. The specific mechanisms underlying angiogenesis and the significance of BK receptors remain under debate.

6. Endothelial Dysfunction

6.1 Endothelium General

The endothelium is a continuous monolayer of cells linked to each other by different type of adhesive structures or junctions that cover the cardiovascular system. For a long time the endothelium was thought to be an inert layer covering the vessels but today we know that it is a dynamic structure that regulates and maintains cardiovascular homeostasis (315).

The endothelium is only 0.5 μm thick and rests on a thin basement membrane. The single cells are connected to each other by junctions (241). During embryonic development, endothelial cells differentiate from common precursor cells and acquire organ-specific properties. An important determinant of the cell differentiation is the local environment, and especially interactions with surrounding cells. These interactions may occur through the release of soluble cytokines, cell-to-cell adhesion, and communication or synthesis of matrix proteins on which the endothelium adheres and grows. Even in the same organ, the endothelium exhibits significant heterogeneity (375).

Two types of endothelial cells are recognized in the heart: endocardial endothelial cells and microvascular endothelial (MVE) cells (241). Both cells produce the same autacoids and share a similar role in signal transduction. The huge difference between them is that the MVECs contribute to cardiac function indirectly by regulating regional blood flow, but the endocardial ECs function directly by communicating with myocardial cells.

The endothelium contributes to cardiovascular homeostasis by regulating vascular permeability and adjusting the caliber of blood vessels according to hemodynamic and hormonal demands. Endothelial cells perform these functions by expressing, activating, and releasing powerful vasoactive substances and other bioactive molecules. These substances include factors that affect vasoconstriction and -vasodilation, coagulation, thrombosis formation, growth, angiogenesis, and tissue remodeling as well as immune reactions and tissue inflammation. The endothelium thus creates a complex and finely tuned balance of interactions with the immediate environment, which are common to all endothelial cells (204;380).

Despite its stable constitutive properties, a normal endothelium reversibly changes its function on activation. For instance, exposure of ECs to inflammatory cytokines induces reprogramming of genes - activation of some and depression of others (280). Senescence of ECs influence endothelial responses, thus senescent cells have defective signaling pathways that lead to an inability to properly responds to stimulants such as growth factors (161).

6.2 Endothelium -derived Vasoactive Substances

Endothelial cells secrete vasodilating mediators in response to substances released from autonomic and sensory nerves, circulating hormones, coagulation derivatives, or autacoids such as bradykinin, angiotensin, and endothelin, secreted by endothelial or smooth muscle cells. Some of these vasodilating substances possess angiogenic and growth-modulating properties. Inflammation, vascular lesions, and thrombosis often accompany impaired vasodilation. It is generally accepted that the net effect of noxious substances and vasoactive, retrieving substances dispose the condition of the endothelium. Thus, there are several mending mechanisms in the endothelium that oppose dysfunction -promoting stimuli, of which the most important ones are mentioned here.

Activation of the kinin receptors induces three of the important mechanisms by which the endothelium reacts by vasodilation, as shown in **Fig 8**.




Fig 8. A schematic view of the mechanisms of BK-BK-2R activation in the endothelium

6.2.1 Nitric Oxide

Nitric oxide (NO) secreted by vascular endothelial cells is a ubiquitous signaling molecule that is able to diffuse across membranes, producing various physiological effects, of which the most important effect is vasodilation (200;343). Three different NOS isoforms have been described, named after the cell type first isolated. They differ in tissue expression, regulation and the average amount of NO produced. The endothelial (eNOS) and neuronal (nNOS) isoforms are constitutively expressed; iNOS is induced under pathological conditions and can produce vast amounts of NO. The synthesis

of NO requires L-arginine as a substrate and binding of Ca^{2+} / calmodulin. BK also stimulates NO production through Ca^{2+} /calmodulin-mediated eNOS activation.

The NO synthases are NADPH-dependent oxygenases that require several co-factors.

eNOS is the dominant enzyme in vascular cells and it is localized in the endothelial caveolae and negatively regulated by caveolin. Indeed eNOS seems to be mainly regulated by modulation of its activity, e.g. by protein kinases (MAP, Akt) (162;171). The inducible NO synthase (iNOS) is primarily regulated transcriptionally and is independent of agonist stimulation or intracellular calcium level (162;171).

A lack of NO is a predominant finding in endothelial dysfunction, which can be created by blocking eNOS and ameliorated by introducing nitrates. Activation of BK-2Rs, but also BK-1Rs, stimulates eNOS to produce NO, leading to vasodilation (132;493).

6.2.2 EDHF

The endothelial-derived hyperpolarizing factor (EDHF) contributes to vasodilation and involves BK-2Rs. Many molecules have been suggested as being the EDHF, and lately the notion has focused on arachidonic acid metabolites, mainly epoxyeicosatrienoic acids (EETs). The origin of the EDHF is still unclear (148). The effect of the EDHF has been demonstrated in various blood vessels from different species, including humans, and it likely plays an important role in cardiovascular physiology. In human coronary arteries, the importance of the EDHF seems to increase as the arterial diameter decreases (461). EDHF-mediated responses are clearly altered in various pathological conditions such as heart failure, atherosclerosis, and hypertension (140).

6.2.3 PGI_2

The vasodilating prostaglandin -prostacyclin (PGI_2)- is primarily produced by ECs (316). BK-2R stimulation activates PGI_2 release through the PLA_2 and arachidonic acid pathway (AA). However, unlike NO, the vasodilating effects of PGI_2 are determined by the expression of specific cyclo-oxygenases (COX-1 and COX-2) in vascular smooth muscle cells, and if the arteries do not express these binding receptors, prostacyclin can not participate in vasodilation (91). Receptor activation in SMCs elevates cyclic nucleotide levels, inducing hyperpolarization thus inhibiting vasocontraction. The link between NO and PGI_2 is bidirectional: prostacyclin can ease the release of NO from ECs and NO potentiates the effect of prostacyclin in vascular SMCs (106;409).

6.2.4 PGE₂

Prostaglandin E₂ is another prostaglandin participating in a wide range of cardiovascular functions, such as vasoconstriction and vasodilation, control of blood pressure, regulation of smooth muscle tone and modulation of inflammation. It has been connected to angiogenesis, though the mechanisms are not completely understood, thus PGE₂ does not stimulate EC migration (187;417), but can induce vascular endothelial growth factor (VEGF) expression and thus relate to angiogenesis via VEGF (417).

6.3 Endothelial Dysfunction

Endothelial dysfunction refers to a pathophysiological state where there is an imbalance between the endothelium-derived relaxing and contracting factors and the availability of NO, in particular, is diminished (466).

This imbalance leads to reduced vasodilation and to activation of pro-inflammatory and pro-thrombotic processes, promoting endothelial dysfunction.

Endothelial dysfunction plays a central role in the initiation and progression of cardiovascular diseases in which the pathophysiological end point is alteration of vascular homeostasis, which leads to diminished tissue blood supply. Diseases like heart failure, coronary arterial disease, and hypertension are all associated with endothelial dysfunction, where diminished availability of NO and increased reactive oxygen species (ROS) formation is detected. EC dysfunction is also related to factors such as smoking, hyperglycemia, and hypercholesterolemia, which are all major risk factors of cardiovascular diseases, but are also connected to the kinin system. Despite the many provoking factors, by far the most important perpetrator of endothelial dysfunction is hypoxia.

6.3.1 Hypoxia

The origin of various vascular pathologies involves instant or recurrent oxygen deficiency. Hypoxia activates endothelial cells by means of a complex system that varies between species, but also between cell types, where the same hypoxic effect can induce different signaling. At first the activation can be viewed as adaptive or protective, but if the stimulus is severe or last long enough, the reaction changes into a maladaptive or lethal one. Following acute hypoxia, endothelial cell activation, neutrophil adherence with local inflammation, and enhanced production of cytokines (IL-6, IL-1 β , MCP-1) are observed, accounting for the inflammatory response in ischemic tissues (412).

Hypoxia increases the cytosolic calcium concentration that regulates many enzymes and participates in signal transduction, such as phospholipase A₂

(PLA₂), which increases in hypoxic ECs, leading to prostaglandin synthesis (20;306). It also up-regulates COX-2 through the induction of NF-kappa- β expression, which also induces ECs to secrete prostaglandins (400).

Even though the endothelial cell layer is relatively tolerant of low oxygen tensions, many of the effects of chronic hypoxia are thought to take place through alterations of endothelial cell signaling. Hypoxia directly affects the release of vasoactive nitric oxide (NO), which locally regulates vascular smooth muscle tone (473), though the physiological mechanisms regulating NO production in the vasculature are not completely understood. Conflicting results on eNOS expression and regulation under hypoxia have been presented and the results seem to vary between species and organs (301;434).

Together with eNOS, also iNOS can be induced in endothelial cells, so the net effect of NO production is even more complicated (342).

In chronic hypoxia, the expression of growth factors, cytokines, and molecules participating in the coagulation process is augmented, affecting remodeling of the vascular wall and myocardium. Many mitogenic molecules released by ECs induce smooth muscle proliferation in vascular beds (338). By far the most potent mitogen in ECs is the vascular endothelial growth factor (VEGF), which initiates angiogenesis *in vivo*. Hypoxia up-regulates the VEGF, but also its receptor, which may explain the strong mitogenic response during hypoxia (476).

Hypoxia transcriptionally induces several genes. The major regulator in cells and tissues is the hypoxia-inducible factor (HIF), an oxygen-sensitive transcription activator. The complex of HIF consists of two proteins: hypoxia-induced HIF-1 α and constitutively expressed HIF-1 β . The activated complex is associated with a specific binding site in target genes and it binds transcriptional co-activators to induce gene expression (214). Today there are over a hundred HIF downstream genes identified, and more than 2% of all human genes are directly or indirectly regulated by HIF in arterial endothelial cells (277). HIF participates in multiple physiological responses to hypoxia, such as erythropoiesis, glycolysis, angiogenesis, and apoptosis.

Many genes, such as VEGF, involved in angiogenesis are activated in hypoxia and regulated by the HIF. The levels of HIF-1 α and VEGF are up-regulated in the myocardium of patients with acute coronary occlusion (250). Effective vascular remodeling in ischemic injury depends on an integrated program of HIF-dependent gene expression (223). Although the HIF is activated in situations where also the kinin system is activated, there are no data relating these systems.

6.3.2 Other Factors Contributing to Endothelial Dysfunction

Hypercholesterolemia induces impairment of endothelium-dependent

vasodilation. Oxidized low-density lipoproteins (LDL) down-regulate the expression of eNOS protein and inhibit the release of both NO and the EDHF, leading to impaired vasodilation (129;472). By disturbing the balance in the endothelium, LDL promotes several vasoconstricting factors (259).

Hyperglycemia with or without insulin deficiency is a major contributor to endothelial dysfunction. Hyperglycemia and its biochemical sequelae directly affect the endothelial function, but also the synthesis of vasoactive substances, growth factors, and cytokines. It is not clear whether impaired endothelial function is due to hyperglycemia or other factors. It is also closely associated with the presence of insulin resistance, regardless of the presence of diabetes.

In hypertension there is an imbalance between the vasoconstrictory and vasodilatory-systems, in particular a reduced vascular availability of endothelium -derived NO. In SHR aorta the endothelium induces cyclooxygenase-related contraction of the underlying vascular smooth muscle cells in response to contracting factors. Patients with essential hypertension have impaired dilatory responses to BK and other stimulants, which can be partially restored by indomethacin (466). Impaired BK -dependent vasodilation in hypertension is also found in normotensive people with a family history of hypertension (432). Several other factors, such as renal insufficiency, aging, smoking, hyper-homocysteinemia, and estrogen-deficiency, also contribute to endothelial dysfunction (330).

6.3.3 Endothelial Dysfunction in Heart Failure

Endothelial dysfunction has been presented in cardiac failure in both animals and humans (222;440). It is found in HF patients with normal coronary circulation or no predisposing factors, suggesting that dysfunction could represent an independent factor in the development of HF (335;449). These patients have impaired relaxation of the coronary microvasculature and peripheral vasculature, but enhanced basal production of NO (237). The relaxation is heterogeneously affected in different vascular beds (335). The damped vasodilating reaction in HF patients together with the release of vasoconstricting substances facilitates poor perfusion and ischemia of the myocardium (118;335).

In animal models, the induction of LVH up-regulates the expression of eNOS.

However during the transition to HF, the expression of eNOS also reduces the secretion of NO (170). This regulation of eNOS has been shown in human end-stage HF and is consistent with BK-2R expression in the progression of HF (239;240).

7. Treatment

The treatment of cardiovascular diseases has progressed during the last decades. As knowledge about the diseases and the molecular mechanisms involved has increased, more possibilities have become available. Despite the increased knowledge and treatment of heart failure, the prognosis has not improved much.

The medical treatment of HF consists of diuretics and vasodilating drugs, such as nitroglycerin, which both relieve the symptoms but have no impact on the prognosis of HF. The β -blockers, previously believed to increase the mortality in HF, are now recommended for the treatment of HF together with diuretics and ACE inhibitors, and shown to have both symptom relieving but also mortality reducing effects also in HF (110). ACE inhibitors/AT-1 receptor inhibitors and statins are widely used in the treatment of HF and they will be discussed here more closely.

7.1 Statins

Statins are in many ways intriguing drugs that were originally developed to lower plasma cholesterol.

Cholesterol is an essential component of cell membranes and also a precursor of steroid hormones and bile acids. However, in excessive quantities, cholesterol becomes a major risk factor of cardiovascular diseases. The rate -limiting enzyme of cholesterol biosynthesis in the liver is HMG-CoA reductase, which catalyses the conversion of HMG-CoA into mevalonic acid (mevanolate) (**Fig 9.**). Inhibitors of HMG-CoA reductase, statins were isolated in 1976 from *Penicillium citrinum*, later named as mevastatin (134). A few years later a more potent substance, lovastatin, was isolated and later approved for clinical use. Since then, several new potent statins have been developed and they are today in wide clinical use.

Statins are potent drugs that cause a dramatic reduction in circulating LDL-cholesterol and an increase in LDL clearance. However, increasing data show that the beneficial effects of statins are only partially cholesterol -dependent and possess many cholesterol-independent or pleiotropic effects.

The Framingham heart study demonstrated that elevated cholesterol is an important risk factor of cardiovascular diseases. The mechanisms attributed to lipid lowering with statins include modification of atherosclerosis progression, atheromatous plaque stabilization, and improved endothelial function. Hence, statins reduce cardiovascular events in both hypercholesterolemic and normocholesterolemic patients with coronary heart disease.

In several studies (4S, CARE, HOPE, LIPID, WOSCOPS) statins have demonstrated beneficial effects in both primary and secondary prevention

of cardiovascular diseases. They also improve heart function and survival in patients with non-ischemic heart failure (194;331). It has been suggested that short-term statin therapy improves cardiac function, neurohormonal imbalance, and symptoms associated with idiopathic dilating cardiomyopathy, suggesting that a non-cholesterol -dependent mechanism is involved (150;194;331). On the other hand, there are also hypotheses, which compromise the beneficial effects of statins in HF, of which the lipoprotein-endotoxin -hypothesis is well known (366).




Fig 9. The cholesterol pathway

By inhibiting the conversion of HMG-CoA to mevalonate, statins prevent the synthesis of important isoprenoids such as farnesylpyrophosphate (FPP) and geranylgeranylpyrophosphate (GGPP), which are precursors of cholesterol but also serve as important intermediate lipid attachments for translational modification of a variety of proteins including small guanosine triphosphate (GTP)-binding proteins Ras, Rho, Rac, and Rab (464). (**Fig. 9.**) Protein isoprenylation permits covalent attachment, subcellular localization, and intracellular trafficking of membrane-bound proteins. Members of the Ras and Rho-GTPase families are important substrates for prenylation, and statins inhibit this action, leading to accumulation of inactive Ras and Rho proteins in the cytoplasm (464). The Rho GTPase -family consists of RhoA, Rac, and Cdc 42 protein subfamilies. The overall functions of Rho GTPases are related to cell shape, motility, secretion, and proliferation. Thus, statins induce a change in the three-dimensional actin cytoskeleton, affect intracellular transport, membrane trafficking, and mRNA stability as well as gene transcription (246). The activation of Rho and its downstream target-

Rho-associated -protein kinase (ROCK) - increases the sensitivity of vascular smooth muscle cells to calcium in hypertension and coronary spasm (218;460). The activation is connected to smooth muscle contraction, oxidative stress, aortic stiffness, and changes in blood pressure. The pathway is also involved in myogenesis, angiogenesis, adipogenesis, cardiac hypertrophy, hypertension, and perivascular fibrosis as well as in the activation of inflammatory cytokines and chemokines (333;423;460).

The Rac pathway is connected to the formation of reactive oxygen species (ROS) and cytoskeleton formation. The growth factor and cytokine -generated ROS are mediated by Rac, and the activation of Rac in the vascular wall is associated with atherosclerosis, myocardial hypertrophy, and endothelial dysfunction (169).

By inhibiting these (Rho and Rac) pathways, statins create non-cholesterol -dependent, pleiotropic effects that seem to be an important factor in cardiovascular protection. The positive effects have also been exhibited by inhibiting the GTPases individually, thus promoting cardioprotective effects such as endothelial function improvement or inhibition of atherosclerosis (276;332).

Statins affect eNOS expression and NO production in many ways. They affect eNOS expression by stabilizing its mRNA and reduce the binding of eNOS to caveolin-1, which as a complex reduces NO formation and activates the PI3K/ Akt pathway, where Akt induces eNOS activation and NO production. Statins also restore eNOS activity in hypoxia and hypercholesterolemia (244;332).

Retrospective analysis of large statin trials, such as 4S, suggests that statins also reduce the incidence of heart failure (229). Recently two small prospective studies showed beneficial effects of statins in human heart failure. These patients showed improved cardiac function and reduced symptoms (248;331).

On the other hand, in the latest CORONA study, rosuvastatin failed to decrease mortality in elderly heart failure patients (228). The GISSI-HF study unfortunately displayed similar results, in patients treated with rosuvastatin (439).

Statins exert many effects beyond cholesterol lowering, which include diminished endothelial dysfunction, vascular inflammation, and inhibition of smooth muscle cell proliferation. These same molecular mechanisms can also be connected to kinins and their receptors, but there is no information concerning BK receptors and statins.

7.2 ACE and AT-Receptor Inhibitors

Angiotensin-converting enzyme inhibitors (ACEI) are effective anti-hypertensive agents, but also very potent in treating and preventing heart

failure. They are also powerful agents in treating and preventing MI, stroke, diabetes, and renal failure.

The protective effects of ACE inhibitors are due to their blocking the formation of Ang II and inhibiting the degradation of BK. Increased BK levels enhance NO synthesis and prostaglandin release through activation of BK-2Rs (232). ACEIs potentiate the effects of agonists on BK-2 receptors by inducing protein-protein cross talk between ACE and BK-2 receptors (283). They can also directly activate BK-1Rs (202). The antifibrotic effect of ACE inhibitors in heart failure are partly mediated by BK-2Rs (156). Many of the cardiovascular effects of ACEIs can be blocked by BK-2R antagonists or mimicked by exogenous BK, suggesting that the effects are mediated by the potentiation of bradykinin and/or BK-2Rs, rather than inhibition of Ang II conversion (232)

The effects of kinins with AT-1R inhibitors (ARBs) are thought to be mediated via activation of AT-2Rs. This activation seems to antagonize several effects induced by stimulation of AT-1Rs such as vasoconstriction, cellular growth, and hypertrophy (404). With AT-1R inhibitor treatment the increased concentrations of angiotensin II and its metabolites act on AT-2Rs and activate local kinin formation. They may also act on the kinin degrading enzymes to increase the tissue BK level (477).

Experimental studies show that kinins are also involved in the effects of AT-1R inhibitors. In an animal study where MI was induced to rats, losartan prevented LVH and HF and the effects were partly attenuated by concomitant inhibition of BK-2Rs (266). During the development of HF following myocardial infarction, BK-2R knockout-mice showed diminished improvement in cardiac function in response to AT-1R inhibitors than control mice (500). In mice that over-express AT-2Rs, infusion of Ang II induces hypotension, which can be abolished by icatibant (458).

The coupling between AT-2R and kallikrein during AT-1R inhibitor treatment and the increased plasma Ang (1-7) levels, which are known to potentiate the effects of bradykinin, may also contribute to the mechanisms related to AT-1R inhibitor-kinin cardioprotection (5;304).

Mounting evidence indicates that kinins are involved in the cardioprotective actions of ACE- and AT -receptor inhibitors the effects are mainly transmitted by BK-2Rs.

7.3. Anti-inflammatory Agents

7.3.1 COX -Inhibitors

Non-steroidal anti-inflammatory drugs (NSAIDs) block the activity of both COX isozymes, COX-1 and COX-2, which mediate enzymatic conversion of

membrane-derived arachidonic acid (AA) to prostaglandin H₂ (PGH₂) and other prostaglandin (PG) metabolites: PGE₂, PGI₂, PGF₂, PGD₂, or thromboxane A₂ (TXA₂). These prostanoids have cardiovascular effects – they can act as either vasodilators or vasoconstrictors, they influence thrombosis formation via platelets, and they alter renal functions. The inducible isoform- COX-2 - is expressed at sites of inflammation such as atherosclerosis.




Fig. 10 The prostaglandin synthesis.

Prostaglandins are widely recognized as inflammatory mediators, but they can also regulate vascular contractility. COX-inhibitors are used as anti-inflammatory and pain relieving mediators. Cyclooxygenase-2 selective inhibitors (COXIBs) were developed with the prime objective of minimizing gastrointestinal adverse effects, which are seen with the use of traditional non-steroidal anti-inflammatory drugs (NSAIDs). During recent years several trials have connected COXIBs to unfavorable effects on cardiovascular outcomes and to increased risk of thromboembolic complications (334;421). At the same time there are controversial results suggesting that inhibition of COX-2 might be cardioprotective. To the latter belongs the fact that COX-2 can be detected from atherosclerotic lesions but not from normal coronary arteries (26). COX-2 knockout-mice develop myocardial fibrosis (111) and the expression of COX-2 seems to be up-regulated by atherosclerosis-inducing stimuli, including free radicals and increased arterial wall shear stress (3;492). And finally, because inflammation of the endothelium is known to diminish its NO -producing capacity, COX-2-inhibition was shown to improve NO production in CHD and hypertensive patients (83;482).

Despite the numerous studies, any conclusion about their cardiovascular safety is complicated due to the conflicting results.

Atherosclerosis is a process with inflammatory features, and selective cyclooxygenase-2 (COX-2) inhibitors may potentially have anti-atherogenic effects

by virtue of inhibiting inflammation. However, by decreasing vasodilatory and anti-aggregatory prostacyclin production, COX-2 antagonists may lead to increased pro-thrombotic activity (167)

The connection with the kinin system has been suggested earlier. BK is known to induce COX-2 expression in different cell types, such as endothelial and smooth muscle cells (54;55). Indomethacin, an unselective COX inhibitor, was able to cause potentiation of BK-induced contraction of both isolated rat uterus and guinea pig tracheal SMC preparations (80;398).

Down-regulation of BK-2Rs has been detected in rabbit hearts treated with a non-specific COX inhibitor, suggesting kinin-prostanoid-induced regulation of BK-2Rs (281).

It was recently also shown that BK induced COX-2 expression in aortic vascular smooth muscle cells, and the effect was mediated by BK-2Rs and involved MAPK p42/p44, PKC, and eNOS activation (379).

Clearly, there are interactions between kinin and prostanoid systems, and the involved kinin receptor seems to be BK-2R, but again, the specific mechanisms and clinical relevance need further studies.

8. BK- Receptors in Heart Failure

Considerable progress has been made in understanding the biology and function of bradykinin receptors. A large body of animal studies implicates a role for BK receptors in vascular pathophysiology. Little is directly known about the clinical relevancy of kinin receptor activation in humans, though an increasing amount of data indicates a cardioprotective role for kinin receptors in humans, also. Direct evidence of BK-receptor involvement in cardiovascular diseases is difficult to acquire, as the vascular segments needed for direct receptor observation are fairly difficult to obtain. Thus, the involvement of kinin receptors is mostly suggestive and based on receptor agonist/antagonist studies or detection of secondary pathway induction. Knockout-models have also provided valuable information about the regulation and function of BK receptors.

8.1 Lessons from Human Studies

8.1.1 BK-2 Receptors

In a prospective study with healthy army recruits, a ten-week physical training program, associated with LV growth response and BK-2R +9/-9 polymorphism and increased LV mass related to the +9/+9 genotype (60). The absence (-) of

a 9 bp repeat in exon 1 of the gene encoding the bradykinin 2 receptor is associated with higher receptor mRNA expression, higher gene transcriptional activity and lower cardiac trophic response to exercise training (56;60). Similar results were obtained from patients suffering from hypertension and LV hypertrophy. The +9/+9 genotypes responded worse than other genotypes to LV mass regression, independent of blood pressure reduction or treatment (174). There was also a strong link to ACE I/D polymorphism, suggesting a synergistic effect between kinin receptors and ACE. The results support the role for bradykinin in the regulation of left-ventricular growth (60).

Several studies associate BK-2R promoter region polymorphism -58 T/C and hypertension, where the C-allele is associated with LVH and/or hypertension at least in African-Americans, Japanese and Chinese subjects (141;155). African-Americans have a significantly increased prevalence, earlier onset, increased mortality in HF, and worse endothelial dysfunction, which could at least partly be explained by the 58 T/C polymorphism and the decreased activity of cardioprotective BK-2Rs. Also, a significantly higher incidence of the C allele of BK-2R was seen in hypertensive patients suffering from myocardial infarction in a Japanese study (18). There is no significant difference in the 58 T/C polymorphism, in the European population; the frequency of the C allele is 0.595 and for the T allele is 0.405, which are in Hardy-Weinberg equilibrium (56;174).

So far, the only human study on heart failure patients and BK-2 receptors was done a few years ago. In human end-stage heart failure patients, a decrease in BK-2Rs in both protein and mRNA levels was observed in the left ventricles of end-stage heart failure patients. There seemed to be no differences in cell-specific expression of BK-2Rs between normal and diseased hearts. The subsequent down-regulation of eNOS suggests a maladaptive connection between these two proteins (239).

The role of kinins in heart diseases has not received much attention, despite the fact that in the 1960s it was shown that both local and systemic administration of BK increased coronary blood flow and improved myocardial metabolism. It took almost 30 years to observe the same in humans. Infused BK doubled blood flow in the human forearm; the effect was abrogated by BK-2R antagonist icatibant (87).

The TREND (trial on reversing endothelial dysfunction) trial showed that ACE inhibitor treatment improved NO-mediated endothelial function in CHD patients. The results were confirmed in healthy and HF patients, where improved endothelial function was abolished by icatibant, indicating the significant role of kinins and BK-2Rs in the endothelium and cardiovascular diseases (192;193;278).

In atherosclerotic patients BK-induced vasodilation is impaired in the plaque area but preserved in spastic areas, suggesting that the EDHF might

have a role in vasodilation in CHD patients. These results were previously suggested in animal models (238;313;441).

A clear correlation between brachial dilation and epicardial dilation was established when bradykinin was infused into subjects with no signs of CHD, who were undergoing coronary angiography (290). Flow-mediated impaired dilation of the brachial artery is considered a non-invasive assessment of coronary endothelial dysfunction. Also in patients undergoing coronary angioplasty (PTCA), infusion of BK mimicked the effects of an ischemic precondition rendering the myocardium relatively resistant to subsequent ischemia (252).

Several studies have demonstrated that the serum of patients with heart failure can alter the function of endothelial cells by down-regulating the expression of endothelial nitric oxide synthase (7). Unfortunately, the role of BK-2Rs was not reported in these studies. ACE inhibitors improve endothelial vasomotor dysfunction in patients with coronary disease, and the effect is mediated by BK-2Rs (360).

Icatibant reduced the hypertensive effect of captopril (ACEI) in patients with hypertension by 53 percent (159).

8.1.2 BK-1 Receptors

Knowledge about BK-1Rs in human heart failure is scarce. Infusion of a promiscuous B1-B2-R antagonist (B9340) decreased forearm blood flow in HF patients taking ACE inhibitors. The effect was absent in patients with no ACE inhibitors as well as in a control group. Icatibant failed to dilate forearm blood flow, thus suggesting that BK-1Rs were mediating dilation in congestive heart failure concomitant with ACE inhibition (487).

In 1997 Raidoo *et al.* discovered that atheromatous arteries exhibit more intense BK-1R labeling in the endothelial cells, foamy macrophages, inflammatory cells, and fibroblasts within plaque and in the intima (364). This suggests that BK-1Rs are relevant in CHD. No other reports on this matter have been published since (364).

Two small studies recently showed that BK-1Rs did not have a major role in peripheral circulation in HF patients, nor could they show any cross-talk between BK-1 and BK-2 receptors (97;245).

The role of BK receptors in human heart failure is not convincing. Although there are numerous studies concerning kinins, knowledge about kinin receptors is weak. To complicate the issue, results from different animal studies have been somewhat controversial, which may at least partly be explained by the different species involved and the different expressions of receptors.

8.2 Lessons from Animal Studies

8.2.1 BK-2 Receptors

The number of BK-2Rs was dramatically decreased during the progression of heart failure in a pressure overload model in spontaneously hypertensive rats (SHR). The number of receptors first increased, correlating to the progression of LVH without fibrosis or ventricular dysfunction. Marked fibrosis and LV dysfunction were observed in the failing hearts, with a loss of BK-2Rs mainly located in the vascular endothelial cells, suggesting a role of the BK-2Rs in end-stage HF. Similar results were obtained from experiments in which acute pressure overload (aortic banding/Ang II infusion) induced the heart failure (240).

ACE inhibitors are the basis of medication in heart failure. Treatment with ACE inhibitors affects BK-2R regulation and potentiates its responses in many ways, not only by inhibiting kinin degradation (284;309). Numerous studies show that the positive cardiovascular effects of ACEIs are partly transmitted via BK-2Rs.

To elucidate the role of BK-2Rs, an orally active BK-2R antagonist (FR173657) was administered to dogs with HF under long-term concomitant treatment with ACEI.

Inhibition of BK-2Rs did not alter systemic blood pressure or LV chamber dimension. Increased LV filling pressure and suppressed expression of eNOS and sarcoplasmic reticulum Ca^{2+} -ATPase mRNA were detected as well as up-regulation of collagen I and III expression. Collagen deposits were increased in the myocardium of the failing hearts in the BK-2R antagonized group, indicating a detrimental role for BK-2Rs in the progression of HF. The treatment did not alter systemic blood pressure or vascular resistance in HF dogs, suggesting also that endogenous BK does not participate in blood pressure regulation via BK-2Rs (156).

BK inhibited the progression of cardiac hypertrophy and myocardial fibrosis in hypertensive rats due to activation of BK-2Rs (437).

Chronic infusion of bradykinin was also able to preserve ventricular and endothelial function in conscious dogs, while delaying the progression of heart failure (447).

The role of BK-2Rs in ischemic preconditioning has been observed in several studies.

The positive effects are reduced infarct size and elimination of post-ischemic arrhythmias by icatibant in preconditioned rats, rabbits, and mice. The positive preconditioning effects are not evident in BK-2R knockout mice nor in kininogen -deficient Brown Norway rats, indicating the significance of the kinin system to cardioprotection in I/R injury (468;501).

BK seems to have a preventive role in the progression of HF, and the effects seem to relate to activation of BK-2Rs and involve NO and also the EDHF, both induced by BK-2Rs.

8.2.2 BK-1 Receptors

BK can exert cardioprotective effects via BK-1Rs, which are known to be involved in the BK-induced reduction of catecholamine release in ischemic rat hearts and in the maintenance of endothelial function in preconditioned hearts (53;77).

There are controversial results concerning the regulation and effects of BK-1Rs in myocardial ischemia. Contradictory results concerning the cardioprotective role of BK-1Rs in myocardial ischemia have been obtained from BK-1R KO-models (242;249;493).

Myocardial sympathetic fibers express BK-1Rs and release noradrenalin. BK inhibits noradrenalin release in a BK-2R dependent way in ischemic rat hearts.

This inhibition was only observed in the presence of BK-1R antagonists, suggesting that BK-1R activation, induced by hypoxia, is involved in myocardial NA release (77;152).

9. Aims of the Study

The general hypothesis of the study was that the kinin receptor system, notably BK-2Rs and BK-1Rs, plays an important role in the pathophysiology of left ventricular hypertrophy and participate in the pathogenesis and progression of heart failure.

The aims of the study were to investigate the regulation of kinin receptors in normal and failing heart tissues, and in particular, in coronary artery endothelial cells. More specifically the aims were:

1. To clarify the expression of BK receptors in normal and failing human heart tissue.
2. To compare the accumulation of two known BK-2R gene polymorphisms affecting gene expression in normal and failing human hearts.
3. To investigate the regulation of BK-receptors in human coronary endothelial cells by both physiological and pharmacological factors, including hypoxia and lovastatin.

10. Materials and Methods

The biochemical methods central to this thesis and their use in the original publications are summarized in table 2.

The methods are only briefly described here, and detailed descriptions with references are provided in the Methods sections of the original publications.

Table 2. Methods used in the original publications

Method	Publication
Preparation of human heart samples	I, IV
RNA isolation	I, II, III, IV
Competitive RT-PCR	I, III
Real-time RT-PCR	II, III
Protein extraction	I, II, III
SDS-PAGE gel electrophoresis	I, II, III
Culture of human endothelial cells	I, II, III
Isolation of rat endothelial cells	III
Culture of rat micro vascular endothelial cells	III
Histological staining	I, III
Immunocytochemistry	II, III
cGMP immunoassay	II
NO measurement	II, III
"Wound healing" assay	III

Human Heart Samples (I, IV)

Normal heart samples ($n=6$) were obtained from the left ventricles of organ donors who had no history of cardiac disease and had been excluded from organ donation because of age, body size, or tissue-type mismatch. The cause of death in these subjects was subarachnoid hemorrhage. The failing left ventricles were harvested at the time of cardiac transplantation from 13 patients with end-stage heart failure (New York Heart Association functional class IV) due to either idiopathic dilated cardiomyopathy (IDC; $n = 7$) or coronary heart disease (CHD; $n= 6$) at the University Central Hospital, Helsinki, Finland. All the patients had been treated with a combination of drugs, including beta-blockers, ACE inhibitors, loop-diuretics, digoxin, and spironolactone. None of the patients had received angiotensin II type I receptor blocker treatment or statin treatment. After excision, the heart tissues were immediately frozen in liquid nitrogen and stored at -70°C . The left ventricle myocardium devoid of visible scar tissue was used in the experiments. An

institutional review board approved the use of normal and failing human heart samples, and the investigation conforms to the principles outlined in the Declaration of Helsinki.

RNA isolation and PCR (I-IV)

Total RNA was isolated from both hearts and cell cultures using an ultrapure TRIzol reagent (GIBCO-BRL) and a RNeasy Mini Kit (QIAGEN). Purified total RNA (0.25 µg for RT-PCR and 1 µg for real-time RT-PCR) was transcribed into cDNA using a Superscript preamplification system (GIBCO-BRL).

Competitive RT-PCR was performed in 25 µl of a standard PCR buffer containing 1 µl of the RT reaction mixture, 25 pmol of sense and antisense primers, 100 µM of each deoxynucleotide, 0.5 units of *Taq* DNA polymerase (Roche), and competitor DNA. The primers used were BK-1R, BK-2R, TNF- α , and GAPDH as the housekeeping gene. The competitor DNAs for BK-1R, BK-2R, and TNF- α were obtained by subcloning the obtained PCR fragments and inserting a 129-bp external DNA fragment into the *Sma*I, *Sac*I, and *Ava*I sites of the respective PCR fragment. The PCR fragments were verified to represent the corresponding targets either by specific restriction enzyme treatment or by DNA sequencing. The PCR products were quantitated with a gel documentation system (Gel Doc 2000; Bio-Rad, Hercules, CA), and when a competitive method was used, the logarithm of target-to-competitor ratio was plotted against the logarithm of the competitor DNA molecules.

For real-time RT-PCR the DNA and mRNA sequences of the target genes were retrieved from the GenBank® database (National Institutes of Health, Bethesda, MD) and gene-specific primers and fluorogenic probes were identified with the Primer Express® software package. The primers used were BK-1R, BK-2R, and GAPDH as the endogenous control. The analyses were performed in 96-well plates (Applied Biosystems) using a TaqMan® Universal Master Mix with uracyl-N-glycosylase treatment in 25 µl reaction volumes. They were analyzed with an ABI Prism 7500 Sequence Detection System using 45 cycles of a two-step program consisting of 15 s at 95 °C and 1 minute at 60 °C. The threshold was set to the geometric phase of the amplification curve, and the amount of target was calculated using the $2^{-\Delta\Delta C_T}$ formula.

Protein Extraction and Western Blotting (I-IV)

Triton X-100 extracts were prepared from heart tissue and cell cultures, subjected to SDS-PAGE, and electro blotted onto nitrocellulose filters (261).

The proteins were detected by immunoblotting using antibodies listed in Table 3. The primary antibodies were further detected with horseradish peroxidase-labelled secondary antibodies and enhanced chemiluminescence

(ECL) as described by the manufacturer (ECL, Amersham Pharmacia Biotech). The translation products were quantitated with a gel documentation system (Gel Doc 2000, Bio-Rad).

Table 3. List of antibodies used

Primary Antibody	Host	Isotype	Source
BK-1R	rabbit	Polyclonal	Santa Cruz
BK-1R	rabbit	Polyclonal	Own product ref.(364)
BK-2R	mouse	Monoclonal	Transduction Lab
TNF-α	mouse	Monoclonal	Santa Cruz
TNF-R1	mouse	Monoclonal	Santa Cruz
β-actin	mouse	Monoclonal	Abcam
RhoA	mouse	Monoclonal	Santa Cruz
Control	mouse	Monoclonal	Serotec

Histological and Immunocytochemical Staining

The frozen human and rat hearts were dissected for immunohistochemical stainings using a standard immunoperoxidase method. Briefly, the slides were fixed with ice-cold methanol. Endogenous peroxidase activity was blocked by means of incubation in 2% H_2O_2 in PBS for 10 minutes. After washing the slides carefully, they were blocked for 60 minutes with 3% normal serum in order to avoid unspecific binding of antibodies. The slides were subsequently incubated with primary antibodies and diluted in corresponding blocking serum overnight at +4 °C. The slides were then washed and a secondary antibody was applied. The nuclei were stained with 4', 6-diamino-2-phenylindole dihydrochloride (DAPI) or with hematoxylin and after washing the mounting media- fluorescent or not- added and examined under a normal or fluorescent microscope, depending on the antibodies used.

The samples subjected to immunohistochemical staining, were stained using primary antibodies listed in Table 2. The cultured endothelial cells were cultured on coverslips until subconfluent. The cells were incubated in the absence or presence of different substances (BK 10-100nM, Icatibant 1 μ M, lovastatin 1nM-100 μ , mevanololactone 100 μ M, Y27632 10 μ M) or in hypoxic conditions. After this the cells were fixed with 4% paraformaldehyde and kept at +4 °C until staining. The primary antibodies were incubated overnight, carefully washed, and stained with secondary antibodies. The immunostained cells were analyzed using fluorescent microscopy (Nikon Eclipse E6000 microscope).

Culture of Human Coronary Arterial Endothelial Cells

Human coronary artery endothelial cells (HCAEC) were purchased from Promo Cell GmbH and used for experiments during passages 4–6. The cells were grown in culture media supplemented with growth factors as recommended by the manufacture.

The cells were incubated in a humidified incubator at 37 °C with a 5% CO₂ atmosphere. The medium was changed every second day and experiments were performed with subconfluent cultures. For the hypoxia experiments, the cells were changed into a low-oxygen containing incubator (1% or 5% O₂) and grown there for the needed time.

Isolation and Culture of Rat Coronary Artery Microvascular Endothelial Cells

Male Wistar rats (300–500 g) were obtained from the Laboratory Animal Center of the University of Helsinki. The rats were treated in accordance with institutional guidelines, which had been approved by the institutional Ethics Committee. The rat myocardium was dissected for perfusion with an enzyme solution for 20 minutes and supplemented with BSA for an additional 10 minutes to facilitate the recovery of separated cells. The perfused tissue was cut into small pieces, incubated with the enzyme solution, and then filtered. The myocardial perfusate and suspension were combined and treated with trypsin, CaCl₂, and BSA in the perfusion buffer for 30 minutes. The cells were collected by centrifugation and resuspended in a culture medium supplemented with FBS and appropriate growth factors and antibiotics. The cells were identified as RCMCs and cultured in M199 growth media at 37 °C under 5% CO₂. The experiments were performed with ECs of 2nd–6th passages in serum-free conditions.

Measurement of NO in Human Coronary Arterial Endothelial Cells

NO production in cultured HCAECs was studied by using 4,5-diamino-fluorescein diacetate (DAF-2-DA) as a fluorescent indicator of intracellular NO, as previously described (462). Briefly, the HCAECs were grown on glass coverslips and treated in the presence or absence of lovastatin or in hypoxic conditions. The HCAECs were pre-incubated for 30 min with L-arginine, pulsed for 10 minutes with DAF-2DA, and washed with PBS. The cells were resuspended in PBS containing L-arginine and activated for 10 minutes with BK. Intracellular fluorescence was visualized with a Nikon Eclipse E600 fluorescence microscope using excitation at 495 nm and emission at 515 nm.

cGMP Immunoassay

cGMP was analyzed with the direct cGMP Enzyme Immunoassay Kit (Assay Designs, Ann Arbor, MI) according to the protocol described by the manufacturer. Briefly, lovastatin was added to the subconfluent plates and incubated with the cells to obtain a high level of functional BK-2Rs in the HCAECs. The lovastatin-treated HCAECs were activated with BK for 30 or 60 minutes after which the cells were washed and endogenous phosphodiesterase activity was inhibited. The cells were lysed with 0.1 M hydrochloric acid and the cell lysate was subjected to overnight acetylation and then used in the competitive cGMP EIA kit as recommended by the manufacturer. The results were analyzed with a microplate reader (Victor³, Perkin Elmer) at 405 nm.

Wound-healing Assay

An *in vitro* wound-healing assay was performed as previously described by Buzza *et al.* (64). The isolated rat coronary microvascular endothelial cells (RCMEC) were cultured on top of thermanox slips in a culture media at +37 °C in a humidified incubator with 5% CO₂. After 2 hours of incubation, non-attached cells were removed by washing. The cells were cultured until subconfluent; an endothelial wound was applied to the slip with an appropriate sized scrape. The treated slips were then carefully washed to remove detached RCMECs and fresh culture media was added in the absence or presence of bradykinin and/or icatibant. The thermanox slips were then incubated under mild hypoxic culture conditions (5% O₂) for 24 or 48 hours to induce BK-2R expression, and fresh BK and/or icatibant was added at the beginning and after 24 h of incubation. The experiment was terminated by washing the slips 3 times with ice-cold PBS (including Ca²⁺/Mg²⁺) and fixed with ice-cold 4% paraformaldehyde (PFA). The fixed RCMECs were washed and the cells were stained with hematoxylin and photographed with a digital camera attached to a Nikon Eclipse E600 microscope.

- / + 9 Polymorphism in Exon 1 of the BK-2 Receptor Gene

Genomic DNA was isolated from human heart tissue using an RNA/DNA Mini Kit (Qiagen) and stored in TE buffer (pH 8.5) in aliquots at -20 °C. Isolated genomic DNA was subjected to PCR using primers spanning the insertion/deletion (-/+ 9 bp) site in exon 1 of the BK-2R gene. The PCR was run at T_m = 66 °C for 40 + 1 cycles, and the obtained PCR products were separated on a 3% MetaPhor agarose gel (BMA).

-58 T/C BK-2 Receptor Promoter Polymorphism by PCR and SSCP

The genomic DNA was subjected to RT-PCR. The obtained PCR products were denatured by heating at 94 °C for 3 minutes in 95% formamide, 10 mM NaOH, 0.25% bromphenol blue, and 0.25% xylene cyanol, and then rapidly cooled on ice. The denatured PCR samples were mixed with 9 parts of SSCP Sucrose Dye and run on a 14% acrylamide/bis (29:1) gel (Bio-Rad). The gel was stained with SYBR Gold nucleic acid gel stain (Molecular Probes) for 30-40 minutes and analyzed with a Gel Doc 2000 gel documentation system (Bio-Rad).

11. Statistical analysis

Data are expressed as means \pm SEM. The patient groups showed normal distribution when analyzed with the Shapiro-Wilks W-test. Correlation studies were performed by using Pearson's correlation and/or ANOVA statistics. Comparisons between the groups were made by using Student's *t*-test. Statistical significance was accepted as $P < 0.05$.

12. Results

BK-1 Receptors in Human Heart Failure

A series of end-stage HF human hearts were analyzed in regard with BK-1Rs and compared with control hearts.

A significant increase in BK-1Rs in both mRNA and protein levels in the diseased hearts was discovered. The BK-1R mRNA expression was increased 2.1 -fold in CHD patients and 2.8 -fold in IDC patients compared with control hearts.

A low transcript level of BK-1R mRNA was also detected in normal myocardias. This was further confirmed by immunohistochemical staining, where a weak staining was detected in intramyocardial vessels also in normal hearts.

The protein quantities of BK-1Rs were also more abundant in HF hearts compared with control ones. The protein levels were somewhat higher in CHD hearts than in IDC hearts, indicating that BK-1Rs are regulated at both transcriptional and translational levels, which has previously been suspected in HF with other genes (240;428). Moreover, the BK-1R/BK-2R mRNA ratio was calculated and there seems to be clear differences between the hearts, thus in the normal myocardias 79.000 copies, 32.000 copies in CHD and 25.000 copies/ 1 μ g RNA in IDC heart samples were detected.

Intense staining of TNF- α was detected in the diseased hearts and the

expression of TNF- α correlated with BK-1R expression in Pearson's correlation test. The immunoreactivity of TNF- α was strongly labelled in the intra-myocardial vessels and specifically in the endothelial cells, as were also the BK-1Rs. Some staining was also detected in the interstitium and in the myocardial cells. Furthermore, incubation of human coronary endothelial cells with and without recombinant TNF- α , and showed, that BK-1R expression was accelerated with the introduction of TNF- α at both the mRNA and protein levels.

TNFR-1 staining was intense in the intra-myocardial vessels of the diseased hearts and the receptors co-localized with the BK-1Rs in the vessel endothelial cells.

BK-2 Receptor Correlation with Age and Polymorphism

The BK-2R mRNA expression was analyzed and a linear regression analysis was done, comparing age and receptor amounts in the control hearts. A clear positive correlation was found, between age and BK-2R mRNA expression, suggesting that transcription of BK-2Rs increases with age in healthy hearts. A positive correlation between age and BK-2R mRNA was also found in the IDC hearts, but a slightly negative correlation in CHD hearts.

Previously it was suggested that BK-2Rs were regulated on both the transcriptional and translational levels in human heart failure (239). By plotting the BK-2R levels against the protein levels from both normal and failing hearts, a positive, linear correlation was found between the mRNA and protein levels, suggesting that BK-2Rs are regulated on the transcriptional level and not on the translational level, as previously suggested.

In the promoter region polymorphism -58 T/C, a marked genotypical frequency of CC allele was found in the IDC hearts. In IDC hearts 5/7 possessed genotype CC (71%) compared with 2/6 of this allele (33%) in normal hearts. Equal quantities of the TT allele were found in IDC and normal hearts, 1/7 and 1/6 (14% and 17%). The dominating genotype in normal hearts was TC, seen in 3/6 (50%). The genotypic frequencies in the CHD hearts were similar to those in normal hearts. furthermore, the allelic frequencies for -58T/C polymorphism were 0.58 for the C-allele and 0.42 for the T-allele in normal and CHD hearts and 0.79 and 0.21, respectively, in the IDC hearts.

For the -9 bp deletion/insertion SNP in exon 1, all the CHD hearts (100%) were homozygous for this deletion (-9/-9). In normal hearts only 1/6 (17%) and 2/7 in IDC hearts (29%) were homozygous for this deletion. Seventy-one percent of the IDC hearts (5/7) were heterozygous (-9/+9) for the deletion and of the normal hearts, 3/6 (50%) were homozygous for the insertion in the allele (+9/+9). See Table 4.

Table 4. Characteristics of human hearts

<i>Subject No.</i>	<i>Etiology</i>	<i>Sex</i>	<i>Age</i>	<i>Polymorphism</i>	
				<i>-58 T/C</i>	<i>+9/-9</i>
<i>1</i>	<i>Normal</i>	<i>M</i>	<i>19</i>	<i>T/C</i>	<i>+9/-9</i>
<i>2</i>		<i>M</i>	<i>43</i>	<i>T/C</i>	<i>+9/+9</i>
<i>3</i>		<i>M</i>	<i>49</i>	<i>C/C</i>	<i>+9/-9</i>
<i>4</i>		<i>M</i>	<i>31</i>	<i>T/T</i>	<i>-9/-9</i>
<i>5</i>		<i>M</i>	<i>40</i>	<i>T/C</i>	<i>+9/+9</i>
<i>6</i>		<i>F</i>	<i>50</i>	<i>C/C</i>	<i>+9/-9</i>
<i>7</i>	<i>CHD</i>	<i>M</i>	<i>61</i>	<i>T/C</i>	<i>-9/-9</i>
<i>8</i>		<i>M</i>	<i>60</i>	<i>T/T</i>	<i>-9/-9</i>
<i>9</i>		<i>M</i>	<i>55</i>	<i>C/C</i>	<i>-9/-9</i>
<i>10</i>		<i>M</i>	<i>52</i>	<i>T/C</i>	<i>-9/-9</i>
<i>11</i>		<i>M</i>	<i>60</i>	<i>C/C</i>	<i>-9/-9</i>
<i>12</i>		<i>M</i>	<i>54</i>	<i>T/C</i>	<i>-9/-9</i>
<i>13</i>	<i>IDC</i>	<i>M</i>	<i>29</i>	<i>C/C</i>	<i>-9/-9</i>
<i>14</i>		<i>M</i>	<i>60</i>	<i>C/C</i>	<i>-9/-9</i>
<i>15</i>		<i>M</i>	<i>46</i>	<i>C/C</i>	<i>+9/-9</i>
<i>16</i>		<i>M</i>	<i>56</i>	<i>C/C</i>	<i>+9/-9</i>
<i>17</i>		<i>M</i>	<i>55</i>	<i>T/T</i>	<i>+9/-9</i>
<i>18</i>		<i>M</i>	<i>56</i>	<i>T/C</i>	<i>+9/-9</i>
<i>19</i>		<i>M</i>	<i>63</i>	<i>C/C</i>	<i>+9/-9</i>

Induction of BK -Receptors by Lovastatin in Cultured Endothelial Cells

Incubation with lovastatin up-regulated BK-2R mRNA expression in cultured human endothelial cells (hCAEC). A dual curve was presented regarding the induction of BK-2R expression. At first a rapid and significant ($> 11x$) augmentation of the receptor mRNA was detected after 12-hour incubation with lovastatin. The BK-2R receptor expression mRNA declined back to the baseline after 24-hour incubation, where after incubating the ECs with a concentration of 100nM lovastatin, increasing levels of BK-2R expression was presented in the cultured cells during 72 hours of incubation. The level of BK-2R mRNA was approximately five-fold greater than in the untreated control cells after 72-hour incubation.

Incubating the cells with lovastatin also induced the expression BK-1R mRNA. Already, after 6-hour lovastatin incubation the BK-1R expression was increased over 23-fold. Also the level of BK-1R expression was declined back to control levels during 24-hour incubation. The expression level of BK-1Rs also showed a tendency of induction, during the prolonged (72 h) incubation of cells with lovastatin, reaching a seven-fold expression level compared with the control cells, not incubated with lovastatin.

The increased BK-2 receptor mRNA transcripts were translated into protein, shown by Western-Blotting analysis, as the protein levels were also up-regulated. The staining of BK-2 receptors was also clearly more intense in endothelial cells treated with lovastatin compared to control EC cultures.

The receptor induction was concentration-dependent and the greatest up-regulation was obtained with a fairly low lovastatin concentration (100 nM). The receptor inducing effect was abolished with higher concentrations ($> 1 \mu\text{M}$) of lovastatin, and the highest concentrations of lovastatin were actually toxic to the endothelial cells.

To establish the functionality of the induced receptors, two indirect methods were used to show production of NO in lovastatin treated cultured hCAECs.

First, in cultured endothelial cells, incubated with lovastatin and at the presence of BK, the production of intra cellular cGMP was significantly increased. The induction required the presence of BK and lovastatin, thus in cells not boosted with BK, or incubated with lovastatin, only a slight elevation of cGMP production was detected.

Secondly, the increased production of NO was verified by a fluorescent DAF-2DA stain. Again, incubating the cells with lovastatin and exogenous BK, the measured fluorescence increased in a BK concentration-dependent way. The increased production of NO was estimated as three times greater than in cells not pulsed by BK.

The fluorescence also markedly intensified with increasing the concentration of lovastatin.

Mevalonate completely abolished the BK-2R induction in lovastatin treated cells, indicating that downstream mediators of the cholesterol pathway were involved in the induction of BK-2Rs.

Lovastatin induced the amount of RhoA protein in cultured endothelial cells but prevented the activation of RhoA, which was shown by immunocytochemical stainings. This was confirmed by incubating the cells with a specific inhibitor (Y-27632), of Rho-associated proteins (ROCK). By inhibiting ROCKs, a time-dependent increase in the expression of BK-2R mRNA was detected, which was similar to the results presented with lovastatin treated cells.

Incubating the ECs together with lovastatin and a specific COX-2 inhibitor (NS398), the BK-2R induction was abrogated. The incubation of mere COX-2 inhibitor had no effect on the BK-2R expression.

Induction of BK-2 Receptors in Hypoxia

In mild hypoxic conditions (5% O_2) the level of both BK-2R mRNA and protein expression was significantly induced compared with cells grown in normoxic conditions. The amount of BK-2R mRNA was time-dependently induced in mild hypoxic conditions. In contrast, severe (1% O_2) hypoxia clearly reduced the BK-2R expression.

The hypoxic induction was abolished by incubating the cells with icatibant, a specific BK-2R antagonist. The protein expression was also induced in mild

hypoxic conditions and where icatibant again abrogated the BK-2R induction.

Functionality of the receptors was verified by DAF-2DA immunocytochemical fluorescent stainings. In hypoxic cells the increased NO production was shown to be approximately three times greater than in normoxic cells. The NO production was induced in hypoxic cells but a clear difference was also detected in cells incubated with or without additional BK in mildly hypoxic chamber, where additional BK increased the amount of NO detected.

These *in vitro* results were then compared with *in vivo* rat experiments. Rats subjected to myocardial infarction by LAD ligation, showed an increased number of BK-2Rs in the endothelium of the myocardial vessels at the infarct border zone, suggesting that the BK-2Rs were also induced *in vivo* by hypoxia. To address this hypothesis, a “wound healing” assay was conducted, previously described by Buzza *et al.* (64). This experiment showed that hypoxic conditions induced endothelial cell migration, which was further augmented by additional BK. The migratory effects were almost abrogated by incubating the hypoxic cells with icatibant. Proliferation of the migrating cells was also detected, thus the cells expressed Ki-76, a commonly used proliferation marker (data not shown).

Hypoxia induced functionally active BK-2Rs in a BK-K-2R-NO -dependent way. The induction of migration and proliferation in hypoxia was increased by BK and abrogated by icatibant.

The main results

In human end-stage HF:

- The BK-1R expression is increased in the intra-myocardial vessels.
- The BK-1R induction is accompanied with TNF- α induction and TNFR-1 co-localized with BK-1Rs.
- The expression of BK-2Rs increases with age in healthy human myocardias.
- The CC -allele of -58 T/C BK-2R SNP is more abundant in IDC hearts, compared to normal or CHD hearts.

In cultured endothelial cells:

- Lovastatin treatment increases the expression of functional BK-2Rs in a time and concentration dependent way. The induction can be abrogated by mevanolate.
- The induction of BK-2Rs in statin treated cultures is also inhibited by a COX-2 inhibitor, NS398.
- Hypoxia induces BK-2R expression in cultured cells. The induction of receptors, NO production but also angiogenesis was clearly increased in hypoxic cultures pulsed with BK. Icatibant abrogated all these effects.

13. Discussion

In the present study, I have focused on understanding the function and regulation of BK receptors in the pathophysiology of human heart failure. HF is a clinical syndrome caused by different cardiovascular diseases and the kinin system seems to be implicated in its pathogenesis. Kinins are powerful bioactive autacoids that have been shown to reduce the myocardial preload and oxygen consumption, thus improve the cardiac metabolism and function. Kinins exert their biological functions by activating specific kinin receptors. In particular, BK activated BK-2Rs are widely accepted as cardioprotective molecules, as they mediate both vasodilatory, antifibrotic, and antiproliferative effects via an increased production of NO, EDHF and PGI₂. BK is also known to counteract endothelial dysfunction, a phenomenon involving a decreased expression of eNOS with subsequent reduction in NO bioavailability, present in the hypoxic myocardium during HF.

The role of BK-1Rs in heart diseases and especially in HF has remained vague. The differences in kinin signaling in normal and failing hearts may depend on either kinin concentrations, kinin degrading enzymes, or on the regulation of BK-receptors, that may account for kinin mediated effects in endothelial cells, myocytes, and fibroblasts.

BK-1Rs and Inflammation

Kinins are proinflammatory peptides acting as local hormones and kinins induce the release of endothelium derived relaxing factors. Many of these effects are mediated through activation of BK-2Rs but in pathological conditions, also BK-1Rs mediate the cardiovascular responses of kinins.

Various studies show an induction of BK-1Rs in pathological conditions, such as tissue trauma, inflammation, or anoxia. Indeed, it has been suggested that BK-1Rs partly account for the initiation of inflammation (132;242). Moreover, the level of inflammatory cytokines is increased in HF, which are known to induce the expression of BK-1Rs (256). The role of BK-1Rs in cardiovascular diseases derives mainly from animal studies and has remained controversial (53;242;292).

In the present study, we showed an increased expression of BK-1Rs in human end-stage failing hearts, implying that induction of BK-1Rs reflects the pathological progression of HF and ongoing inflammation. Previous studies in hypertensive rats show that up-regulation of BK-1Rs seems to be associated with the transition from compensated LV hypertrophy to heart failure, with the onset of myocardial fibrosis and with the development of diastolic dysfunction (240;452).

In previous study, Kuoppala *et al* showed low levels of BK-2Rs in the

LV of end-stage HF patients and the study detected a significant increase in fibrosis in IDC hearts compared with normal hearts (239). In the present study we showed increased levels of BK-1Rs, as well as an increased ratio of BK-1R/BK-2R in the myocardium of the HF hearts, suggesting that inflammation accompanies the increased level of fibrosis in these failing hearts. This has previously been addressed in human lung fibroblasts, where induction of BK-1Rs activated fibrogenesis. Moreover, in stenotic aortic valves, a significant induction of BK-1Rs was detected in the diseased valves compared to control valves. Myofibroblasts, cultured from the stenotic valves, were also more susceptible towards TNF- α induced BK-1R up-regulation (181;293).

Our study also showed a correlation of the increased TNF- α levels and TNFR-1 expression in the intra-myocardial coronary vessels with induced BK-1R levels as well as co-localization of BK-1 and TNFR-1s. The results imply that an inflammatory context in HF is a regulator element in BK-1Rs expression.

Whether the upregulation of BK-1Rs in the failing human hearts reflects increased production of cytokines or a compensation of the BK-2R down-regulation can presently not be answered. However, knockout studies have shown that BK-1Rs are up-regulated in the heart of BK-2R knockout-mice and they can takeover some of the hemodynamic properties of BK-2Rs, suggesting that the BK-1Rs can take a compensatory role at least in the absence of BK-2Rs, and thus be cardioprotective (124).

The highest level of BK-1Rs and the lowest of BK-2Rs were detected in IDC hearts, suggesting that the etiology of HF and BK-receptor regulation are interrelated. The observation that the level of fibrosis in the end-stage HF patients was higher in the IDC hearts compared to CHD hearts supports this notion (239). The differences in receptor levels between CHD and IDC may also be explained by differences in the genetics, a notion supported by previously observed differential expression of several genes in coronary arteries from CHD and IDC patients (388).

Deletion of BK-1Rs does not, under basal conditions, affect cardiovascular function or growth and BK-1R knockout-mice do not develop hypertension or myocardial hypertrophy nor HF (242;352), suggesting that BK-1Rs have a diminutive role on cardiovascular regulation under normal situations. On the other hand, animal models and human artery *in vitro* studies have shown that BK-1Rs can regulate vascular tone (230;286;487). Under healthy conditions, the level of BK-1Rs is undetectable or extremely low in human myocardias. Our perception of a low BK-1R level also in control hearts, suggests that BK-1Rs may exert some function in the normal cardiovascular regulation. It may also merely reflect a normal aging process, previously suggested in animal studies (225).

BK-2Rs and Aging

In our study, the level of BK-2Rs correlated positively with age in healthy hearts, suggesting that healthy human heart adapts to ageing or senescence by inducing BK-2R expression.

The association between BK-2Rs and senescence has been recently reported in bovine endothelial cells, where BK dose dependently protected the cells from ROS induced senescence through pathways involving BK-2R and NO (336). In addition, in a report from Kakoki *et al*, BK-2R knockout-mice displayed a slightly accelerated ageing phenotype. The ageing phenotype was strongly promoted when the BK-2R knockout was combined with diabetes (212). These results strongly suggest, that BK-2Rs are associated with senescence and protect against its damaging effects.

We also reported a positive correlation between BK-2R mRNA and age in IDC hearts, where as the correlation was negative in CHD patients. This can reflect the some what older patient material of the CHD group, but may again also imply disease-dependent regulation of BK-2Rs or a genetic variation.

Contradictory results were reported from animal studies, where unchanged levels of BK-2R mRNA were detected, from young and old rats. Whether this reflects normal ageing process in rats or differences in cellular composition in ageing hearts can not be answered. A discrepancy was detected in the study by Kintsurashvili *et al*, between the gene expression and protein translation, thus in old rats the BK-2R protein levels were two-fold lower as compared to young ones (225). The differences between gene expression and protein translation in the study suggests, that BK-2Rs are regulated post-transcriptionally, which was also our finding in the end-stage human hearts.

BK-2 Receptor Polymorphism

Racial differences in survival outcomes, point to genetics in the predisposition of HF and to date two BK-2R polymorphisms have been connected to hypertension and HF (119). In our studies the C-allele in -58 T/C SNP was significantly more abundant in IDC patients which could account for the development of HF and also the lower level of BK-2Rs detected in IDC patients. The age-related up-regulation of BK-2Rs was impaired in IDC hearts, suggesting that high frequency of the C-allele, detected in IDC hearts, might relate to the lower BK-2Rs levels detected.

As for the +9/-9 insertion/deletion SNP, all of our CHD hearts possessed the -9 deletion. None of the diseased hearts and 33% of the control hearts were homozygous for the +9 insertion, suggesting that the known genetics of +/-9 was not associated with down-regulation of BK-2Rs in our patients.

BK-2Rs and Statins

As the endothelium is the major contributor of kinin receptor function, we have also studied endothelium and factors modulating kinin receptors in endothelial cells. In many clinical studies, statins have reduced mortality in CHD patients. Subclass analyses have suggested that statins might also be beneficial in HF patients without CHD, though these results have not been confirmed.

Our novel results show that lovastatin is able to induce both BK-receptors types in cultured human endothelial cells. The induction of BK-2Rs is concentration-dependent and clinically-relevant doses (lovastatin 80mg) seem to correlate with the greatest BK-2R induction. The effects were abrogated by mevanolate, implying that the down-stream mediators of cholesterol biosynthesis pathways are involved. More specifically, it involved inhibition of RhoA-Rho kinase-mediated downstream signaling pathways, previously been shown to have an important role in cellular functions in the pathogenesis of cardiovascular diseases. Rho-kinase up-regulates several molecules that accelerate inflammation and fibrosis, and down-regulates eNOS (247;434).

The present study showed that exogenous BK accelerated the up-regulated BK-2R expression and increased NO production in lovastatin-treated human ECs. A direct regulatory link between BK-2R and eNOS has been shown, where the level of eNOS and also production of NO depend on the level of BK-2R. (23;509).

In a dog model BK induced NO production in renal and coronary arteries, which was attenuated with HF dogs and retrieved by simvastatin.(4;451). Moreover, the BK induced relaxation in rat mesentery arteries was enhanced by simvastatin, which resulted from inhibition of RhoA and attenuation of the vascular contractility through augmenting the endothelial NO production in mesenteric arteries (408). Unfortunately, no data concerning BK receptors was reported.

By reducing the prenylation of isoprenoids, statins inhibit Rho-kinases, accelerated by several inflammatory stimuli such as IL-1 β and angiotensin II (186). In our studies, the statin effect on BK-2Rs was reversed by both mevanolate and icatibant and mimicked by Rho inhibitor, indicating the involvement of Rho in the signal transduction leading to BK-2R expression.

The results suggest that BK increases NO production via activating BK-2Rs and eNOS in statin-treated endothelial cells. This involves the RhoA-Rho-kinase pathway, thus implying that some of the pleiotropic effects of statins may be explained by induction on BK-2Rs, at least in endothelial cells.

The relevance of BK-1R induction by lovastatin remains unclear. Statin treatment attenuates inflammatory signaling pathways, implying another mechanism for the induction of BK-1Rs, in statin treated cells.

It has previously been shown in a murine model that ACE inhibitors can up-regulate functional BK-1Rs, suggesting that also BK-1Rs have beneficial

cardiovascular effects in the BK-BK-R signaling cascade, without inflammation (286). Moreover, deletion of BK-2R has been shown to induce the expression of BK-1Rs, suggesting a role for these receptors in the cardiovascular regulation (213).

Conflicting results were reported recently, where disruption of BK-2Rs resulted in early development of LVH and down-regulation of eNOS. The effects were abrogated by simvastatin and the authors concluded that simvastatin reversed the effects by activating the myocardial eNOS and MAP kinase (339). Moreover, no differences in BK-1R expression were detected between wild type or BK-2R knockout mice nor had simvastatin treatment any effect on the BK-1R expression. The discrepancies between these studies can relate to differences in species studied, to different knockout variants used or differences in experimental set-ups.

The exact mechanisms, by which statins induce BK-receptors, is presently not known, although a direct gene activating effect may be involved. In the light of former studies our *in vitro* results suggest that in heart failure, patients might benefit from statin treatment through induction of BK-2Rs resulting in increased NO production.

BK-2Rs and COX-2

Of the two known COX-isoforms, COX-1 is traditionally viewed as the constitutive enzyme, responsible for the production of the physiological amounts of prostaglandins needed. COX-2 is induced by various inflammatory stimuli and cytokines, and considered as a mediator of inflammation and pain.

COX-2 expression can be found both in the vascular ECs and SMCs and in the ECs, it mainly forms prostacyclin (PGI_2), which induces vasodilation. Prostacyclin is an important secondary mediator, often released by kinins from the endothelium, which stimulates cyclic AMP in vascular smooth muscle cells leading to vasorelaxation. Prostanoid formation occurs in many cell types, which possess kinin receptors.

Statin treatment has previously been shown to inhibit the generation of COX-2-derived prostanoid production and also down-regulate COX-2 expression in both endothelial and vascular smooth muscle cells (98). Recently several opposing reports have been published, where statins induce COX-2 expression and PGI_2 formation in human endothelial cells through inhibition of Rho (105;288).

In our study, the induction of statin treatment on BK-2R expression was reversed by COX-2 inhibitor (NS398). Moreover, no effect on BK-2Rs was detected when incubating the cells merely with a COX-2 inhibitor, nor could we detect any differences in COX-1 expression, suggesting a direct link

between BK-2R and COX-2 pathways in endothelial cells.

A connection between BK-2Rs and COX-2 has previously been suggested by Marceau *et al.* In this study diclofenac induced BK-2R down-regulation in rat hearts, although suspected, they could not show a feed-back mechanism between COX-2 and BK-2Rs (281).

Previous studies reported, that exogenous BK was able to induce COX-2 expression, in different cell types (55;344). Moreover, in rat aortic VSMC, BK induced the expression of COX-2, and the effect was reversed by icatibant and L-NAME, but not by a BK-1R inhibitor. (379). Also, in rabbits infusion of COX-2 inhibitor (diclofenac) induced BK-2R down-regulation in the rabbit heart (281). In a recent study, the positive post-conditioning effect of BK was abolished by a COX-inhibitor, which was thought to relate to decreased production of PGI₂ (349). COX-2 induction by BK can be blocked by icatibant or L-NAME, suggesting, that at least in ECs, the induction of COX-2, involves activation of BK-2 receptors and eNOS. Several studies show, that RhoGTPases are involved in the regulation of COX-2, although the exact mechanisms are still not clear.

Thus, it can be suggested that, in lovastatin treated endothelial cells, BK induces BK-2Rs, increasing the production of NO and PGI₂ via eNOS and COX-2, respectively, leading to increased vasodilation in the vasculature. Whether the COX-2 effect relates to increased prostacyclin production, to decreased production of contracting prostanoids or to some other mechanism, can not be addressed.

It can also be speculated, whether the negative cardiovascular effects of COX-2 inhibitors observed in large clinical studies, partly relates to the down-regulation of BK-2Rs.

Endothelial Hypoxia and Preconditioning

Hypoxia is a common factor in cardiovascular diseases, regardless of etiology. Hypoxia stimulates BK production and the expression of eNOS is known to increase in hypoxic conditions. Several studies focusing on the effects of BK in both animals and humans exists, but again the significance of BK-receptors in hypoxia is not conclusively studied. In our *in vitro* studies mild hypoxia induced functional BK-2Rs in hCAECs, but also in coronary artery smooth muscle cells (CASMC), and the effect was reversed by icatibant. This might give a mechanistical explanation for pre- and postconditioning, as BK is known to be one of the main effector molecules in preconditioning. In experimental models, short intervals of repeated hypoxia are known to be beneficial to the heart, improving the outcome after induced myocardial infarction (323). Postconditioning also augments the deleterious effects induced by reperfusion, and activation of BK-BK-2Rs is involved in this

cardioprotection (350). Hence, our results from hypoxic ECs, might reflect the protection seen in pre-and postconditioning. Indeed, the cardioprotective effects of preconditioning are abolished in BK-2R knockout mice and BK induces and icatibant attenuates the cardioprotective effects in a rabbit model of preconditioning (475). Moreover, in a recent study, postconditioning or intermittent BK infusion was shown to protect the myocardium and involve activation of COX-2 and subsequent PGI₂ formation (349).

We showed that hypoxia also induced migration of endothelial cells, which was abrogated by icatibant, suggesting a role for BK-2Rs in angiogenesis. This is consistent with previous results in which BK-2Rs were shown to transactivate VEGF receptors (KDR/Flk-1) and to induce tube formation in hCAECs (310). Similar results have been shown in mouse hearts, where angiotensin II and hypoxia significantly increased angiogenesis via AT-2Rs and the effects were BK-BK-2R and eNOS -dependent (322). Enalapril was also able to induce sprouting in hypoxic mouse hearts through both BK-1Rs and BK-2Rs, requiring the presence of both BK and BK-2Rs. Thus, BK and other vasoactive molecules can promote angiogenesis in hypoxic mouse hearts by activating BK-2Rs (384).

Our results showed that under hypoxic conditions, induction of BK-2Rs in cultured ECs was clearly attenuated by exogenous BK, which also increased the basal NO production.

Our results from rats, recovering from myocardial infarction, showed an induction of BK-2Rs in the endothelium of vessels forming at the border zone between the scar tissue and healthy myocardium, i.e. where potential ischemia augments BK formation and BK-2R expression and enhances angiogenesis. This further implies that the pathophysiological mechanisms activated during ischemic injuries also involve activation of the kinin receptor system.

Supporting our results are also the observations, that ACE inhibitor treatment or tissue kallikrein gene delivery increased myocardial capillary density and accelerated cardiac remodeling in SHR in a BK-2R-dependent way (48;164).

Our results show that mild hypoxia induces the expression of functional BK-2Rs capable of increasing NO production and that BK-2Rs play a role in angiogenesis and EC migration in hypoxic conditions. There are also indications that PGI₂ via COX-2 induction might enhance the cardioprotective effects of kinin system.

In cardiovascular diseases, such as hypertension and CHD, which often lead to clinical HF, the cardioprotective role of BK-2Rs has been shown.

BK and BK-2Rs oppose endothelial dysfunction and contribute to angiogenesis(132;322). The kinin system decreases preload and cardiac output, thus improving myocardial metabolism and performance. BK also exerts a unique pattern of injury limiting actions in preconditioning (444;475). It exerts

cardioprotective effects when administered prior to the onset of ischemia, participates in early preconditioning, and is also an important factor in late preconditioning (235;402;468). ACE inhibitors and NEP inhibitors increase bradykinin concentration in the myocardium, thus they increase the protective effects of BK in ischemic diseases (325;446;483). Exogenous bradykinin, augmenting endogenous BK levels or activating BK-2Rs, possibly also BK-1Rs, could have an impact on the treatment of ischemic heart diseases.

From previous results hypothesis can be made that during the progression from compensated left ventricular hypertrophy to fulminate heart failure, the heart tries to compensate for ongoing changes by increasing the number of BK-2Rs, but in the end fails to do so and clinical HF develops, accompanied by diminished BK-2R levels and increased BK-1R levels(239;260). Is the diseased myocardium unable to react to BK and induce the BK-2Rs, in end-stage hearts, where the induced myocardial growth is accompanied with diminished NO production and unbalanced Ang II secretion. Or, are there other factors contributing to the down-regulation of BK-2Rs and progression of HF, can not be answered. Also, wheater the induction of BK-1Rs is a compensatory mechanism towards the down-regulation of BK-2Rs or a response to the inflammatory cascades activated, are not known.

The relevance of BK-1Rs in heart failure is still debatable. If the up-regulation of BK-1Rs in end-stage HF is cardioprotective, the induction of BK-1R could represent compensatory effect for BK-2R down-regulation. Several studies show, that BK-1Rs take over the protective effects of BK-2Rs, when the BK-2R are genetically or pharmacologically deleted. There are also several contradictory reports, were the induction of BK-1Rs accompanies activation of cardiotoxic pathways and progression of the cardiovascular disease.

Clinical Implications

Statin treatment has been shown to have several cholesterol-independent mechanisms that counteract endothelial function. Inhibition of the Rho-Rho kinase pathway partly accounts for the beneficial pleiotropic effects of statins and relates to the induction of BK-2Rs, thus improving the vascular endothelial dysfunction.

The beneficial cardiovascular effects of ACE inhibitor treatment are generally considered to arise from the prevention of BK degradation and the induction of BK-2Rs thus promoting vasodilation in endothelial cells, and inhibition of myocyte hypertrophy. Blockade of the BK-2R, partially prevents the cardiovascular benefits of ACE inhibition in different disease models, demonstrating the pharmacological relevance of bradykinin signaling in cardiovascular disease (263).

As BK is degraded by both ACE and NEP, simultaneous inhibition prevents the degradation more efficiently than ACE inhibition alone, leading to increased plasma and tissue concentrations of BK (40). The most extensively studied dual inhibitor, omapatrilat, has been studied in human heart failure, and it has shown superior results in HF when compared to ACE inhibitors. Unfortunately severe angioedema, in higher frequencies in patients receiving omapatrilat, has been reported and because of these serious concerns, it has not been approved for the treatment of HF (19).

Several chemical and physiological factors regulate the BK-R expression. Moreover, kinins are known to affect most pathological conditions leading to LVH and HF. Actions intensifying either BK and/or BK-2Rs, might give additional tools for the treatment of HF. The molecular mechanisms leading to HF are not clear, and among other pathways, the kinin system and factors regulating it, need more studies, in order to find therapeutical drugs for treating HF.

14. Summary and Conclusions

1. In the human end-stage heart failure functional BK-1Rs were significantly increased in the intra myocardial coronary vessels. This induction was accompanied by accumulation of TNF- α and the receptors co-localized with TNFR1 in the myocardial vessels, referring that the induced BK-1Rs relate to inflammation and increased fibrosis in heart failure. This implies that inflammation regulates the expression of BK-1Rs in human HF.

Because the known cardioprotective actions of BK are mediated via BK-2Rs, whereas BK-1Rs are induced by tissue damage, these results suggest that the decreased BK-2R levels in the end-stage HF, may leave the hearts vulnerable to damages, and increases in the BK-1R expression and activity may represent a compensatory reaction in the diseased hearts.

2. The expression of BK-2Rs increased with age in healthy human hearts, which suggests that the human heart adapts to age-related changes in heart, by up-regulating the expression of cardioprotective BK-2Rs.

From previously known polymorphisms, the 9bp insertion/deletion SNP didn't seem to have a role in the progression of HF. The C-allele of -58 T/C SNP was more abundant in the IDC patients, which may account for the lower BK-2R mRNA levels detected and the development of HF.

Our results also suggest that different signaling cascades are activated in CHD and IDC patients.

-
3. Lovastatin treatment clearly increased the amount of functional BK-2Rs in the endothelial cells, involving Rho-Rho kinase pathways, previously shown to have an impact on the progression of cardiovascular diseases. Thus, some of the beneficial cardiovascular effects of statins may relate to increased number of BK-2Rs in the endothelium.

A connection between COX-2 expression and BK-2R was also presented. Our results arose an intriguing question of whether the use of COX-inhibitors and the increased risk of myocardial infarction, relates to down-regulation of BK-2Rs.

Our results suggest that HF patients benefit from statin treatment also via induction of BK-2Rs and increased NO production, thus improving the endothelial function. We also suggest that by decreasing the level of BK-2Rs, COX-inhibitors promote dysfunction of the endothelium and together with increased thrombosis activation, may worsen the outcome of HF patients.

4. Mild hypoxia induced the expression of BK-Rs in cultured endothelial cells. Exogenous BK induced the expression of BK-2Rs and also the migration of endothelial cells in hypoxic conditions. This migration was abrogated by inhibiting the BK-2Rs, suggesting a role for BK-2Rs on EC migration and proliferation in hypoxic conditions. The angiogenesis promoting effects were also shown in an infarct model, where BK-2Rs were localized at the infarct border zone, where hypoxia likely exists. Hypoxia induces BK and BK-2R formation and NO production, which promote angiogenesis and the healing of the myocardium.

15. Acknowledgments

The research for this thesis was carried out at the Wihuri Research Institute during 2001-2009. I wish to express my gratitude to Wihuri Research Institute and to the Jenny and Antti Wihuri Foundation, who made my research possible.

I wish to express my gratitude to Professor Petri Kovanen, head of the Research Institute for providing the tools and facilities for my thesis. The atmosphere of old Salus Hospital has been inspiring.

I especially want to thank my supervisor Docent Ken Lindstedt who, over the years, always had a positive and encouraging attitude. During these years I did several periods of clinical work, when my motivation for this thesis, was completely lost. Ken's inspiring attitude, support, and help has been incredible and allowed me to finish my thesis. Thank you Ken!

I also want to thank Docent Jorma Kokkonen, who in the first place, guided me to Wihuri. Over the years, I have many times thought about your words, concerning the difficulties of laboratory work, which at the time I did not believe. How right you were!

I also want to acknowledge technical assistance of Jaana Tuomikangas, Elina Kaperi, Leena Saikko, Suvi Socolnicki and Mari Jokinen who all helped me with my, some times tangled experiments. Jaana taught me the basic skills of laboratory work and was always there to support me when my experiments didn't succeed. Your positive attitude is admirable and I'm happy for our friendship. Lennu has also been a great help and assisted me with my experiments, even when not working in Wihuri anymore. I have enjoyed our weekly badminton games, where new ideas and resolutions to problems have popped out. Not to mention the regular gym sessions with Jaana and Lennu...

I am grateful to Naotaka Shiota and Antti Kuoppala who taught me the basics of kinins and helped me to get started with the laboratory work. I also want to thank Mikko Mäyränpää, Markus Leskinen, Hanna Heikkilä, Jukka Hakala, Riina Oksjoki, Anna Oksaharju, Mia Sneek, Jani Lappalainen, Laura Nurmi, Kati Öörni, Katariina Lähdesmäki, and Mikko Simolin for their help and support during these years. The help of Laura Fellman is also appreciated.

I also wish to thank all the people working in Wihuri for a pleasant atmosphere and enjoyable coffee breaks as well as all our parties.

Hanna Leskinen and Heikki Ruskoaho, from the University of Oulu, are warmly thanked for collaboration.

I am very grateful to Professor Heikki Vapaatalo and Docent Risto Kerkelä for their thorough review of this thesis. I sincerely thank Keith Kosola, for careful revision of the language of this thesis.

I also want to thank Professor Ville Valtonen, Asko Järvinen and all my colleges at the Division of Infectious Diseases, for a pleasant and inspiring working atmosphere and I look forwards to working with you in the future.

I warmly thank my father, Matti Romo, for his never ending support and interest in my thesis. There have been times when your belief towards my work, has been my only motivation, in continuing the work! Your vice advices expand beyond medicine or my thesis, and they have been an incredible help in many things over the years.

I am also very grateful to my mother-in-law, Pippe and Kurre, for their support as well as help with our children.

I want to give special thanks to my sister Tilli as well as numerous friends, who have believed in my work; especially I wish to thank Minna Thelen and Mats Wolontis, for their interest and support for my work during these years. Finally, I owe my deepest thanks to my husband Kai and our sons Alex and Anton for their support, love, and patience towards my thesis.

I am thankful for financial support from Aarne Koskelo Foundation, Finnish Foundation of Cardiovascular Diseases, Duodecim, and Finnish Society of Angiology.

16. References

- (1) Abbas F, Clayton JK, Marshall KM, Senior J. Characterisation of kinin receptors on the human isolated umbilical artery. *J Endocrinol* 1998;156:389-94.
- (2) Abdouh M, Khanjari A, Abdelaziz N, Ongali B, Couture R, Hassessian HM. Early upregulation of kinin B1 receptors in retinal microvessels of the streptozotocin-diabetic rat. *Br J Pharmacol* 2003;140:33-40.
- (3) Adderley SR, Fitzgerald DJ. Oxidative damage of cardiomyocytes is limited by extracellular regulated kinases 1/2-mediated induction of cyclooxygenase-2. *J Biol Chem* 1999;274:5038-46.
- (4) Adler S, Huang H, Trochu JN, Xu X, Gupta S, Hintze TH. Simvastatin reverses impaired regulation of renal oxygen consumption in congestive heart failure. *Am J Physiol Renal Physiol* 2001;281:F802-F809.
- (5) Agata J, Ura N, Yoshida H, Shinshi Y, Sasaki H, Hyakkoku M, Taniguchi S, Shimamoto K. Olmesartan is an angiotensin II receptor blocker with an inhibitory effect on angiotensin-converting enzyme. *Hypertens Res* 2006;29:865-74.
- (6) Agata J, Chao L, Chao J. Kallikrein gene delivery improves cardiac reserve and attenuates remodeling after myocardial infarction. *Hypertension* 2002;40:653-9.
- (7) Agnoletti L, Curello S, Bachetti T, Malacarne F, Gaia G, Comini L, Volterrani M, Bonetti P, Parrinello G, Cadei M, Grigolato PG, Ferrari R. Serum from patients with severe heart failure downregulates eNOS and is proapoptotic: role of tumor necrosis factor-alpha. *Circulation* 1999;100:1983-91.
- (8) Ahluwalia A, Perretti M. Involvement of bradykinin B1 receptors in the polymorphonuclear leukocyte accumulation induced by IL-1 beta in vivo in the mouse. *J Immunol* 1996;156:269-74.
- (9) Akbary AM, Wirth KJ, Scholkens BA. Efficacy and tolerability of Icatibant (Hoe 140) in patients with moderately severe chronic bronchial asthma. *Immunopharmacology* 1996;33:238-42.
- (10) Alfie ME, Sigmon DH, Pomposiello SI, Carretero OA. Effect of high salt intake in mutant mice lacking bradykinin-B2 receptors. *Hypertension* 1997;29:483-7.
- (11) Alfie ME, Yang XP, Hess F, Carretero OA. Salt-sensitive hypertension in bradykinin B2 receptor knockout mice. *Biochem Biophys Res Commun* 1996;224:625-30.
- (12) Allen TJ, Cao Z, Youssef S, Hulthen UL, Cooper ME. Role of angiotensin II and bradykinin in experimental diabetic nephropathy. Functional and structural studies. *Diabetes* 1997; 46:1612-8.
- (13) Alric C, Pecher C, Schanstra JP, Bascands JL, Girolami JP. Bradykinin-induced inhibition of cell proliferation and tyrosine kinase activity in rat mesangial cells. *Int J Mol Med* 2000;5:85-93.
- (14) Altamura M, Meini S, Quartara L, Maggi CA. Nonpeptide antagonists for kinin receptors. *Regul Pept* 1999;80:13-26.
- (15) Am Heart Association. Heart Disease and Stroke Statistics: 2005 Update. Am Heart Association, Dallas Texas, 2005.
- (16) Anderson RG. Caveolae: where incoming and outgoing messengers meet. *Proc Natl Acad Sci U S A* 1993;90:10909-13.
- (17) Antonio A. The relaxing effect of bradykinin on intestinal smooth muscle. *Br J Pharmacol Chemother* 1968;32:78-86.
- (18) Aoki S, Mukae S, Itoh S, Sato R, Nishio K, Iwata T, Katagiri T. The genetic factor in acute myocardial infarction with hypertension. *Jpn Circ J* 2001;65:621-6.
- (19) Armstrong PW, Lorell BH, Nissen S, Borer J. Omapatrilat. *Circulation* 2002;106:e9011-e9012.
- (20) Arnould T, Michiels C, Alexandre I, Remacle J. Effect of hypoxia upon intracellular calcium concentration of human endothelial cells. *J Cell Physiol* 1992;152:215-21.

-
- (21) Audet R, Rioux F, Drapeau G, Marceau F. Cardiovascular effects of Sar-[D-Phe⁸]des-Arg⁹-bradykinin, a metabolically protected agonist of B1 receptor for kinins, in the anesthetized rabbit pretreated with a sublethal dose of bacterial lipopolysaccharide. *J Pharmacol Exp Ther* 1997;280:6-15.
- (22) Austin CE, Faussner A, Robinson HE, Chakravarty S, Kyle DJ, Bathon JM, Proud D. Stable expression of the human kinin B1 receptor in Chinese hamster ovary cells. Characterization of ligand binding and effector pathways. *J Biol Chem* 1997;272:11420-5.
- (23) Bachetti T, Comini L, Pasini E, Cargnoni A, Curello S, Ferrari R. Ace-inhibition with quinapril modulates the nitric oxide pathway in normotensive rats. *J Mol Cell Cardiol* 2001; 33:395-403.
- (24) Bachvarov DR, Hess JF, Menke JG, Larrivee JF, Marceau F. Structure and genomic organization of the human B1 receptor gene for kinins (BDKRB1). *Genomics* 1996;33:374-81.
- (25) Bachvarov DR, Houle S, Bachvarova M, Bouthillier J, Adam A, Marceau F. Bradykinin B(2) receptor endocytosis, recycling, and down-regulation assessed using green fluorescent protein conjugates. *J Pharmacol Exp Ther* 2001;297:19-26.
- (26) Baker CS, Hall RJ, Evans TJ, Pomerance A, Maclouf J, Creminon C, Yacoub MH, Polak JM. Cyclooxygenase-2 is widely expressed in atherosclerotic lesions affecting native and transplanted human coronary arteries and colocalizes with inducible nitric oxide synthase and nitrotyrosine particularly in macrophages. *Arterioscler Thromb Vasc Biol* 1999;19:646-55.
- (27) Bandeira-Melo C, Calheiros AS, Silva PM, Cordeiro RS, Teixeira MM, Martins MA. Suppressive effect of distinct bradykinin B2 receptor antagonist on allergen-evoked exudation and leukocyte infiltration in sensitized rats. *Br J Pharmacol* 1999;127:315-20.
- (28) Barabe J, Marceau F, Theriault B, Drouin JN, Regoli D. Cardiovascular actions of kinins in the rabbit. *Can J Physiol Pharmacol* 1979;57:78-91.
- (29) Barabe J, Park WK, Regoli D. Application of drug receptor theories to the analysis of the myotropic effects of bradykinin. *Can J Physiol Pharmacol* 1975;53:345-53.
- (30) Barnes PJ. Bradykinin and asthma. *Thorax* 1992;47:979-83.
- (31) Bas M, Bier H, Greve J, Kojda G, Hoffmann TK. Novel pharmacotherapy of acute hereditary angioedema with bradykinin B2-receptor antagonist icatibant. *Allergy* 2006;61:1490-2.
- (32) Belichard P, Loillier B, Paquet JL, Luccarini JM, Pruneau D. Haemodynamic and cardiac effects of kinin B1 and B2 receptor stimulation in conscious instrumented dogs. *Br J Pharmacol* 1996;117:1565-71.
- (33) Benzing T, Fleming I, Blaukat A, Muller-Esterl W, Busse R. Angiotensin-converting enzyme inhibitor ramiprilat interferes with the sequestration of the B2 kinin receptor within the plasma membrane of native endothelial cells. *Circulation* 1999;99:2034-40.
- (34) Bergaya S, Meneton P, Bloch-Faure M, Mathieu E, Alhenc-Gelas F, Levy BI, Boulanger CM. Decreased flow-dependent dilation in carotid arteries of tissue kallikrein-knockout mice. *Circ Res* 2001;88:593-9.
- (35) Bhoola K, Ramsaroop R, Plendl J, Cassim B, Dlamini Z, Naicker S. Kallikrein and kinin receptor expression in inflammation and cancer. *Biol Chem* 2001;382:77-89.
- (36) Bhoola KD, Bewley J, Crothers DM, Cingi MI, Figueroa CD. Kinin receptors on epithelial cells and smooth muscle of the trachea. *Adv Exp Med Biol* 1989;247:421-7.
- (37) Bhoola KD, Figueroa CD, Worthy K. Bioregulation of kinins: kallikreins, kininogens, and kininases. *Pharmacol Rev* 1992;44:1-80.
- (38) Bhoola R, Ramsaroop R, Naidoo S, Muller-Esterl W, Bhoola KD. Kinin receptor status in normal and inflamed gastric mucosa. *Immunopharmacology* 1997;36:161-5.
- (39) Biyashev D, Tan F, Chen Z, Zhang K, Deddish PA, Erdos EG, Hecquet C. Kallikrein activates bradykinin B2 receptors in absence of kininogen. *Am J Physiol Heart Circ Physiol* 2006; 290: H1244-H1250.
- (40) Blais C Jr., Lapointe N, Rouleau JL, Clement R, Gervais N, Geadah D, Adam A. Effects of the vasopeptidase inhibitor omapatrilat on cardiac endogenous kinins in rats with acute myocardial

- infarction Peptides 2001;22:953-62.
- (41) Blais C Jr., Marc-Aurele J, Simmons WH, Loute G, Thibault P, Skidgel RA, Adam A. Des-Arg9-bradykinin metabolism in patients who presented hypersensitivity reactions during hemodialysis: role of serum ACE and aminopeptidase P. Peptides 1999;20:421-30.
- (42) Blais C Jr., Marceau F, Rouleau JL, Adam A. The kallikrein-kininogen-kinin system: lessons from the quantification of endogenous kinins. Peptides 2000;21:1903-40.
- (43) Blaukat A, Alla SA, Lohse MJ, Muller-Esterl W. Ligand-induced phosphorylation/dephosphorylation of the endogenous bradykinin B2 receptor from human fibroblasts. J Biol Chem 1996;271:32366-74.
- (44) Blaukat A, Herzer K, Schroeder C, Bachmann M, Nash N, Muller-Esterl W. Overexpression and functional characterization of kinin receptors reveal subtype-specific phosphorylation. Biochemistry 1999;38:1300-9.
- (45) Blaukat A, Micke P, Kalatskaya I, Faussner A, Muller-Esterl W. Downregulation of bradykinin B2 receptor in human fibroblasts during prolonged agonist exposure. Am J Physiol Heart Circ Physiol 2003;284:H1909-H1916.
- (46) Blaukat A, Pizard A, Breit A, Wernstedt C, Alhenc-Gelas F, Muller-Esterl W, Dikic I. Determination of bradykinin B2 receptor in vivo phosphorylation sites and their role in receptor function. J Biol Chem 2001;276:40431-40.
- (47) Blaukat A, Alla SA, Lohse MJ, Muller-Esterl W. Ligand-induced Phosphorylation/Dephosphorylation of the Endogenous Bradykinin B2 Receptor from Human Fibroblasts. J Biol Chem 1996;271:32366-74.
- (48) Bledsoe G, Chao L, Chao J. Kallikrein gene delivery attenuates cardiac remodeling and promotes neovascularization in spontaneously hypertensive rats. Am J Physiol Heart Circ Physiol 2003;285:H1479-H1488.
- (49) Boels K, Schaller HC. Identification and characterisation of GPR100 as a novel human G-protein-coupled bradykinin receptor. Br J Pharmacol 2003;140:932-8.
- (50) Boissonnas RA, Guttman S, Jaqueoud PA, Konzett H, Stuermer E. Synthesis and biological activity of peptides related to bradykinin. Experientia 1960;16:326.
- (51) Bonner G, Preis S, Schunk U, Toussaint C, Kaufmann W. Hemodynamic effects of bradykinin on systemic and pulmonary circulation in healthy and hypertensive humans. J Cardiovasc Pharmacol 1990;15:S46-S56.
- (52) Borkowski JA, Ransom RW, Seabrook GR, Trumbauer M, Chen H, Hill RG, Strader CD, Hess JF. Targeted disruption of a B2 bradykinin receptor gene in mice eliminates bradykinin action in smooth muscle and neurons. J Biol Chem 1995;270:13706-10.
- (53) Bouchard JF, Chouinard J, Lamontagne D. Role of kinins in the endothelial protective effect of ischaemic preconditioning. Br J Pharmacol 1998;123:413-20.
- (54) Bradbury DA, Corbett L, Knox AJ. PI 3-kinase and MAP kinase regulate bradykinin induced prostaglandin E(2) release in human pulmonary artery by modulating COX-2 activity. FEBS Lett 2004;560:30-4.
- (55) Bradbury DA, Newton R, Zhu YM, El-Haroun H, Corbett L, Knox AJ. Cyclooxygenase-2 induction by bradykinin in human pulmonary artery smooth muscle cells is mediated by the cyclic AMP response element through a novel autocrine loop involving endogenous prostaglandin E2, E-prostanoid 2 (EP2), and EP4 receptors. J Biol Chem 2003;278:49954-64.
- (56) Braun A, Maier E, Kammerer S, Muller B, Roscher AA. A novel sequence polymorphism in the promoter region of the human B2-bradykinin receptor gene. Hum Genet 1996;97:688-9.
- (57) Braz JC, Gregory K, Pathak A, Zhao W, Sahin B, Klevitsky R, Kimball TF, Lorenz JN, Nairn AC, Liggett SB, Bodi I, Wang S, Schwartz A, Lakatta EG, Paoli-Roach AA, Robbins J, Hewett TE, Bibb JA, Westfall MV, Kranias EG, Molken JD. PKC-alpha regulates cardiac contractility and propensity toward heart failure. Nat Med 2004;10:248-54.
- (58) Bristow MR. beta-adrenergic receptor blockade in chronic heart failure. Circulation 2000;101:558-69.

-
- (59) Brown NJ, Gainer JV, Murphey LJ, Vaughan DE. Bradykinin stimulates tissue plasminogen activator release from human forearm vasculature through B(2) receptor-dependent, NO synthase-independent, and cyclooxygenase-independent pathway. *Circulation* 2000; 102:2190-6.
- (60) Brull D, Dhamrait S, Myerson S, Erdmann J, Woods D, World M, Pennell D, Humphries S, Regitz-Zagrosek V, Montgomery H. Bradykinin B2BKR receptor polymorphism and left-ventricular growth response. *Lancet* 2001;358:1155-6.
- (61) Bugge E, Ytrehus K. Bradykinin Protects Against Infarction but Does not Mediate Ischemic Preconditioning in the Isolated Rat Heart. *Journal of Molecular and Cellular Cardiology* 1996; 28:2333-41.
- (62) Burrell LM, Risvanis J, Kubota E, Dean RG, MacDonald PS, Lu S, Tikellis C, Grant SL, Lew RA, Smith AI, Cooper ME, Johnston CI. Myocardial infarction increases ACE2 expression in rat and humans. *Eur Heart J* 2005;26:369-75.
- (63) Busse R, Fleming I. Regulation and functional consequences of endothelial nitric oxide formation. *Ann Med* 1995;27:331-40.
- (64) Buzza MS, Zamurs L, Sun J, Bird CH, Smith AI, Trapani JA, Froelich CJ, Nice EC, Bird PI. Extracellular Matrix Remodeling by Human Granzyme B via Cleavage of Vitronectin, Fibronectin, and Laminin. *J Biol Chem* 2005;280:23549-58.
- (65) Cabrini DA, Campos MM, Tratsk KS, Merino VF, Silva JA, Jr., Souza GE, Avellar MC, Pesquero JB, Calixto JB. Molecular and pharmacological evidence for modulation of kinin B(1) receptor expression by endogenous glucocorticoids hormones in rats. *Br J Pharmacol* 2001;132:567-77.
- (66) Calabro P, Limongelli G, Riegler L, Maddaloni V, Palmieri R, Golia E, Roselli T, Masarone D, Pacileo G, Golino P, Calabro R. Novel insights into the role of cardiotrophin-1 in cardiovascular diseases *J Mol Cell Cardiol* 2009;46:142-8.
- (67) Calixto JB, Cabrini DA, Ferreira J, Campos MM. Kinins in pain and inflammation. *Pain* 2000; 87:1-5.
- (68) Calixto JB, Medeiros R, Fernandes ES, Ferreira J, Cabrini DA, Campos MM. Kinin B1 receptors: key G-protein-coupled receptors and their role in inflammatory and painful processes. *Br J Pharmacol* 2004;143:803-18.
- (69) Calvillo L, Latini R, Kajstura J, Leri A, Anversa P, Ghezzi P, Salio M, Cerami A, Brines M. Recombinant human erythropoietin protects the myocardium from ischemia-reperfusion injury and promotes beneficial remodeling. *Proc Natl Acad Sci U S A* 2003;100:4802-6.
- (70) Campbell DJ, Duncan AM, Kladis A. Angiotensin-converting enzyme inhibition modifies angiotensin but not kinin peptide levels in human atrial tissue. *Hypertension* 1999;34:171-5.
- (71) Campbell DJ, Kladis A, Briscoe TA, Zhuo J. Type 2 bradykinin-receptor antagonism does not modify kinin or angiotensin peptide levels. *Hypertension* 1999;33:1233-6.
- (72) Campbell DJ, Kladis A, Duncan AM. Bradykinin peptides in kidney, blood, and other tissues of the rat. *Hypertension* 1993;21:155-65.
- (73) Campos MM, Cabrini DA, Cardozo AH, Rae GA, Toro JH, Calixto JB. Changes in paw oedema triggered via bradykinin B(1) and B(2) receptors in streptozotocin-diabetic rats. *Eur J Pharmacol* 2001;416:169-77.
- (74) Cayla C, Todiras M, Iliescu R, Saul VV, Gross V, Pilz B, Chai G, Merino VF, Pesquero JB, Baltatu OC, Bader M. Mice deficient for both kinin receptors are normotensive and protected from endotoxin-induced hypotension. *FASEB J* 2007;21:1689-98.
- (75) Ceconi C, Francolini G, Olivares A, Comini L, Bachetti T, Ferrari R. Angiotensin-converting enzyme (ACE) inhibitors have different selectivity for bradykinin binding sites of human somatic ACE. *European Journal of Pharmacology* 2007;577:1-6.
- (76) Cervenka L, Maly J, Karasova L, Simova M, Vitko S, Hellerova S, Heller J, El-Dahr SS. Angiotensin II-induced hypertension in bradykinin B2 receptor knockout mice. *Hypertension* 2001;37:967-73.
- (77) Chahine R, Adam A, Yamaguchi N, Gaspo R, Regoli D, Nadeau R. Protective effects of

- bradykinin on the ischaemic heart: implication of the B1 receptor. *Br J Pharmacol* 1993; 108:318-22.
- (78) Chambers LS, Black JL, Ge Q, Carlin SM, Au WW, Poniris M, Thompson J, Johnson PR, Burgess JK. PAR-2 activation, PGE₂, and COX-2 in human asthmatic and nonasthmatic airway smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 2003;285:L619-L627.
- (79) Chao J, Bledsoe G, Yin H, Chao L. The tissue kallikrein-kinin system protects against cardiovascular and renal diseases and ischemic stroke independently of blood pressure reduction. *Biol Chem* 2006;387:665-75.
- (80) Chaud M, Franchi AM, Rettori V, McCann SM, Gimeno MF. Nitric oxide in the contractile action of bradykinin, oxytocin, and prostaglandin F₂ alpha in the estrogenized rat uterus. *Proc Natl Acad Sci U S A* 1997;94:11049-54.
- (81) Chen X, Orfanos SE, Ryan JW, Chung AY, Hess DC, Catravas JD. Species variation in pulmonary endothelial aminopeptidase P activity. *J Pharmacol Exp Ther* 1991;259:1301-7.
- (82) Chen Z, Tan F, Erdos EG, Deddish PA. Hydrolysis of angiotensin peptides by human angiotensin I-converting enzyme and the resensitization of B2 kinin receptors. *Hypertension* 2005;46:1368-73.
- (83) Chenevard R, Hurlimann D, Bechir M, Enseleit F, Spieker L, Hermann M, Riesen W, Gay S, Gay RE, Neidhart M, Michel B, Luscher TF, Noll G, Ruschitzka F. Selective COX-2 inhibition improves endothelial function in coronary artery disease. *Circulation* 2003; 107:405-9.
- (84) Cheng CP, Onishi K, Ohte N, Suzuki M, Little WC. Functional effects of endogenous bradykinin in congestive heart failure. *J Am Coll Cardiol* 1998;31:1679-86.
- (85) Cherry PD, Furchgott RF, Zawadzki JV, Jothianandan D. Role of endothelial cells in relaxation of isolated arteries by bradykinin. *Proc Natl Acad Sci U S A* 1982;79:2106-10.
- (86) Christopher J, Velarde V, Zhang D, Mayfield D, Mayfield RK, Jaffa AA. Regulation of B(2)-kinin receptors by glucose in vascular smooth muscle cells. *Am J Physiol Heart Circ Physiol* 2001;280:H1537-H1546.
- (87) Cockcroft JR, Chowienzyk PJ, Brett SE, Bender N, Ritter JM. Inhibition of bradykinin-induced vasodilation in human forearm vasculature by icatibant, a potent B2-receptor antagonist. *Br J Clin Pharmacol* 1994;38:317-21.
- (88) Cohen MC, Huberman MS, Nesto RW. Recombinant alpha 2 interferon-related cardiomyopathy. *Am J Med* 1988;85:549-51.
- (89) Cohen MV, Yang XM, Liu GS, Heusch G, Downey JM. Acetylcholine, bradykinin, opioids, and phenylephrine, but not adenosine, trigger preconditioning by generating free radicals and opening mitochondrial K(ATP) channels. *Circ Res* 2001;89:273-8.
- (90) Cohn JN, Levine TB, Olivari MT, Garberg V, Lura D, Francis GS, Simon AB, Rector T. Plasma norepinephrine as a guide to prognosis in patients with chronic congestive heart failure. *N Engl J Med* 1984;311:819-23.
- (91) Coleman RA, Smith WL, Narumiya S. International Union of Pharmacology classification of prostanoid receptors: properties, distribution, and structure of the receptors and their subtypes. *Pharmacol Rev* 1994;46:205-29.
- (92) Colman RW, Schmaier AH. Contact system: a vascular biology modulator with anticoagulant, profibrinolytic, antiadhesive, and proinflammatory attributes. *Blood* 1997;90:3819-43.
- (93) Colucci WS, Braunwald E. Pathophysiology of Heart Failure in Heart Diseases. A Textbook of cardiovascular medicine, 6th edition, 503-533. 2001.
- (94) Couture R, Harrisson M, Vianna RM, Cloutier F. Kinin receptors in pain and inflammation. *Eur J Pharmacol* 2001;429:161-76.
- (95) Cowie MR, Mosterd A, Wood DA, Deckers JW, Poole-Wilson PA, Sutton GC, Grobbee DE. The epidemiology of heart failure. *Eur Heart J* 1997;18:208-25.
- (96) Crackower MA, Sarao R, Oudit GY, Yagil C, Kozieradzki I, Scanga SE, Oliveira-dos-Santos AJ, da Costa J, Zhang L, Pei Y, Scholey J, Ferrario CM, Manoukian AS, Chappell MC, Backx PH, Yagil Y, Penninger JM. Angiotensin-converting enzyme 2 is an essential regulator of heart function. *Nature* 2002;417:822-8.

-
- (97) Cruden NL, Tse GH, Fox KA, Ludlam CA, Megson I, Newby DE. B1 kinin receptor does not contribute to vascular tone or tissue plasminogen activator release in the peripheral circulation of patients with heart failure. *Arterioscler Thromb Vasc Biol* 2005;25:772-7.
- (98) Cuccurullo C, Mezzetti A, Cipollone F. COX-2 and the vasculature: angel or evil? *Curr Hypertens Rep* 2007;9:73-80.
- (99) Cyr M, Lepage Y, Blais C, Jr., Gervais N, Cugno M, Rouleau JL, Adam A. Bradykinin and des-Arg(9)-bradykinin metabolic pathways and kinetics of activation of human plasma. *Am J Physiol Heart Circ Physiol* 2001;281:H275-H283.
- (100) Damas J, Cession-Fossion A. Are bradykinin receptors of smooth muscle and adrenal glands different? *C R Seances Soc Biol Fil* 1973;167:1478-82.
- (101) de Bold AJ, Borenstein HB, Veress AT, Sonnenberg H. A rapid and potent natriuretic response to intravenous injection of atrial myocardial extract in rats. *Life Sci* 1981;28:89-94.
- (102) De Weerd WF, Leeb-Lundberg LM. Bradykinin sequesters B2 bradykinin receptors and the receptor-coupled G alpha subunits G alphaq and G alpha i in caveolae in DDT1 MF-2 smooth muscle cells. *J Biol Chem* 1997;272:17858-66.
- (103) deBlois D, Bouthillier J, Marceau F. Effect of glucocorticoids, monokines and growth factors on the spontaneously developing responses of the rabbit isolated aorta to des-Arg9-bradykinin. *Br J Pharmacol* 1988;93:969-77.
- (104) deBlois D, Horlick RA. Endotoxin sensitization to kinin B(1) receptor agonist in a non-human primate model: haemodynamic and pro-inflammatory effects. *Br J Pharmacol* 2001;132:327-35.
- (105) Degraeve F, Bolla M, Blaie S, Creminon C, Quere I, Boquet P, Levy-Toledano S, Bertoglio J, Habib A. Modulation of COX-2 expression by statins in human aortic smooth muscle cells. Involvement of geranylgeranylated proteins. *J Biol Chem* 2001;276:46849-55.
- (106) Delpy E, Coste H, Gouville AC. Effects of cyclic GMP elevation on isoprenaline-induced increase in cyclic AMP and relaxation in rat aortic smooth muscle: role of phosphodiesterase. *Br J Pharmacol* 1996;119:471-8.
- (107) Dendorfer A, Wolfrum S, Schafer U, Stewart JM, Inamura N, Dominiak P. Potentiation of the vascular response to kinins by inhibition of myocardial kininases. *Hypertension* 2000;35:32-7.
- (108) Dendorfer A, Wolfrum S, Wellhoner P, Korsman K, Dominiak P. Intravascular and interstitial degradation of bradykinin in isolated perfused rat heart. *Br J Pharmacol* 1997;122:1179-87.
- (109) DeWitt BJ, Cheng DY, Kadowitz PJ. des-Arg9-bradykinin produces tone-dependent kinin B1 receptor-mediated responses in the pulmonary vascular bed. *Circ Res* 1994;75:1064-72.
- (110) Dickstein K, Cohen-Solal A, Filippatos G, McMurray JJ, Ponikowski P, Poole-Wilson PA, Stromberg A, Van Veldhuisen DJ, Atar D, Hoes AW, Keren A, Mebazaa A, Nieminen M, Priori SG, Swedberg K. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2008. *Rev Esp Cardiol* 2008;61:1329-70.
- (111) Dinchuk JE, Car BD, Focht RJ, Johnston JJ, Jaffee BD, Covington MB, Contel NR, Eng VM, Collins RJ, Czerniak PM, et al. Renal abnormalities and an altered inflammatory response in mice lacking cyclooxygenase II. *Nature* 1995;378:406-9.
- (112) Dixon BS, Evanoff D, Fang WB, Dennis MJ. Bradykinin B1 receptor blocks PDGF-induced mitogenesis by prolonging ERK activation and increasing p27Kip1. *Am J Physiol Cell Physiol* 2002;283:C193-C203.
- (113) Donaldson CJ, Sutton SW, Perrin MH, Corrigan AZ, Lewis KA, Rivier JE, Vaughan JM, Vale WW. Cloning and characterization of human urocortin. *Endocrinology* 1996;137:2167-70.
- (114) Donnini S, Solito R, Giachetti A, Granger HJ, Ziche M, Morbidelli L. Fibroblast growth factor-2 mediates Angiotensin-converting enzyme inhibitor-induced angiogenesis in coronary endothelium. *J Pharmacol Exp Ther* 2006;319:515-22.
- (115) Donoghue M, Hsieh F, Baronas E, Godbout K, Gosselin M, Stagliano N, Donovan M, Woolf B, Robison K, Jeyaseelan R, Breitbart RE, Acton S. A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circ Res* 2000; 87: E1-E9.

- (116) Dray A. Kinins and their receptors in hyperalgesia. *Can J Physiol Pharmacol* 1997;75:704-12.
- (117) Dray A, Perkins M. Bradykinin and inflammatory pain. *Trends Neurosci* 1993;16:99-104.
- (118) Drexler H, Hayoz D, Munzel T, Hornig B, Just H, Brunner HR, Zelis R. Endothelial function in chronic congestive heart failure. *Am J Cardiol* 1992;69:1596-601.
- (119) Dries DL, Exner DV, Gersh BJ, Cooper HA, Carson PE, Domanski MJ. Racial differences in the outcome of left ventricular dysfunction. *N Engl J Med* 1999;340:609-16.
- (120) Drummond GR, Cocks TM. Endothelium-dependent relaxation to the B1 kinin receptor agonist des-Arg9-bradykinin in human coronary arteries. *Br J Pharmacol* 1995;116:3083-5.
- (121) Drummond GR, Cocks TM. Endothelium-dependent relaxations mediated by inducible B1 and constitutive B2 kinin receptors in the bovine isolated coronary artery. *Br J Pharmacol* 1995;116:2473-81.
- (122) Duka A, Kintsurashvili E, Duka I, Ona D, Hopkins TA, Bader M, Gavras I, Gavras H. Angiotensin-converting enzyme inhibition after experimental myocardial infarct: role of the kinin B1 and B2 receptors. *Hypertension* 2008;51:1352-7.
- (123) Duka I, Duka A, Kintsurashvili E, Johns C, Gavras I, Gavras H. Mechanisms mediating the vasoactive effects of the B1 receptors of bradykinin. *Hypertension* 2003;42:1021-5.
- (124) Duka I, Kintsurashvili E, Gavras I, Johns C, Bresnahan M, Gavras H. Vasoactive potential of the b(1) bradykinin receptor in normotension and hypertension. *Circ Res* 2001;88:275-81.
- (125) Dumoulin MJ, Adam A, Blais C Jr., Lamontagne D. Metabolism of bradykinin by the rat coronary vascular bed. *Cardiovasc Res* 1998;38:229-36.
- (126) Duncan AM, James GM, Anastasopoulos F, Kladis A, Briscoe TA, Campbell DJ. Interaction between neutral endopeptidase and angiotensin converting enzyme inhibition in rats with myocardial infarction: effects on cardiac hypertrophy and angiotensin and bradykinin peptide levels. *J Pharmacol Exp Ther* 1999;289:295-303.
- (127) Duncan AM, Kladis A, Jennings GL, Dart AM, Esler M, Campbell DJ. Kinins in humans. *Am J Physiol Regul Integr Comp Physiol* 2000;278:R897-R904.
- (128) Eggerickx D, Raspe E, Bertrand D, Vassart G, Parmentier M. Molecular cloning, functional expression and pharmacological characterization of a human bradykinin B2 receptor gene. *Biochem Biophys Res Commun* 1992;187:1306-13.
- (129) Eizawa H, Yui Y, Inoue R, Kosuga K, Hattori R, Aoyama T, Sasayama S. Lysophosphatidylcholine inhibits endothelium-dependent hyperpolarization and N{omega}-nitro-L-arginine/indomethacin resistant endothelium-dependent relaxation in the porcine coronary artery. *Circulation* 1995;92:3520-6.
- (130) Ellis KM, Cannet C, Mazzoni L, Fozard JR. Airway hyperresponsiveness to bradykinin induced by allergen challenge in actively sensitised Brown Norway rats. *Naunyn Schmiedebergs Arch Pharmacol* 2004;369:166-78.
- (131) Emanuelli C, Madeddu P. Angiogenesis therapy with human tissue kallikrein for the treatment of ischemic diseases. *Arch Mal Coeur Vaiss* 2004;97:679-87.
- (132) Emanuelli C, Maestri R, Corradi D, Marchione R, Minasi A, Tozzi MG, Salis MB, Straino S, Capogrossi MC, Olivetti G, Madeddu P. Dilated and failing cardiomyopathy in bradykinin B(2) receptor knockout mice. *Circulation* 1999;100:2359-65.
- (133) Emanuelli C, Minasi A, Zacheo A, Chao J, Chao L, Salis MB, Straino S, Tozzi MG, Smith R, Gaspa L, Bianchini G, Stillo F, Capogrossi MC, Madeddu P. Local delivery of human tissue kallikrein gene accelerates spontaneous angiogenesis in mouse model of hindlimb ischemia. *Circulation* 2001;103:125-32.
- (134) Endo A, Kuroda M, Tsujita Y. ML-236A, ML-236B, and ML-236C, new inhibitors of cholesterol synthesis produced by *Penicillium citrinum*. *J Antibiot (Tokyo)* 1976;29:1346-8.
- (135) Engelhardt S, Hein L, Wiesmann F, Lohse MJ. Progressive hypertrophy and heart failure in beta1-adrenergic receptor transgenic mice. *Proc Natl Acad Sci U S A* 1999;96:7059-64.
- (136) Eric J, Gabra BH, Sirois P. Implication of the bradykinin receptors in antigen-induced pulmonary inflammation in mice. *Br J Pharmacol* 2003;13:1589-97.

-
- (137) Ersahin C, Simmons WH. Inhibition of both aminopeptidase P and angiotensin-converting enzyme prevents bradykinin degradation in the rat coronary circulation. *J Cardiovasc Pharmacol* 1997;30:96-101.
- (138) Fang J, Mensah GA, Croft JB, Keenan NL. Heart failure-related hospitalization in the U.S., 1979 to 2004. *J Am Coll Cardiol* 2008;52:428-34.
- (139) Feletou M, Tang EH, Vanhoutte PM. Nitric oxide the gatekeeper of endothelial vasomotor control. *Front Biosci* 2008;13:4198-217.
- (140) Feletou M, Vanhoutte PM. EDHF: new therapeutic targets? *Pharmacol Res* 2004;49:565-80.
- (141) Ferdinand KC. African American heart failure trial: role of endothelial dysfunction and heart failure in African Americans. *Am J Cardiol* 2007;99:3D-6D.
- (142) Fernandes L, Fortes ZB, Nigro D, Tostes RC, Santos RA, Catelli De Carvalho MH. Potentiation of bradykinin by angiotensin-(1-7) on arterioles of spontaneously hypertensive rats studied in vivo. *Hypertension* 2001;37:703-9.
- (143) Ferreira J, Campos MM, Pesquero JB, Araujo RC, Bader M, Calixto JB. Evidence for the participation of kinins in Freund's adjuvant-induced inflammatory and nociceptive responses in kinin B1 and B2 receptor knockout mice. *Neuropharmacology* 2001;41:1006-12.
- (144) Ferreira SH, Vane JR. The disappearance of bradykinin and eledoisin in the circulation and vascular beds of the cat. *Br J Pharmacol Chemother* 1967;30:417-24.
- (145) Field JL, Butt SK, Morton IK, Hall JM. Bradykinin B2 receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci. *Br J Pharmacol* 1994;113:607-13.
- (146) Figueroa CD, Marchant A, Novoa U, Forstermann U, Jarnagin K, Scholkens B, Muller-Esterl W. Differential distribution of bradykinin B(2) receptors in the rat and human cardiovascular system. *Hypertension* 2001;37:110-20.
- (147) Fleming I, Fisslthaler B, Busse R. Calcium signaling in endothelial cells involves activation of tyrosine kinases and leads to activation of mitogen-activated protein kinases. *Circ Res* 1995;76:522-9.
- (148) Fleming I, Rueben A, Popp R, Fisslthaler B, Schrodt S, Sander A, Haendeler J, Falck JR, Morisseau C, Hammock BD, Busse R. Epoxyeicosatrienoic acids regulate Trp channel dependent Ca²⁺ signaling and hyperpolarization in endothelial cells. *Arterioscler Thromb Vasc Biol* 2007;27:2612-8.
- (149) Folkman J, Shing Y. Angiogenesis. *J Biol Chem* 1992;267:10931-4.
- (150) Foody JM, Shah R, Galusha D, Masoudi FA, Havranek EP, Krumholz HM. Statins and mortality among elderly patients hospitalized with heart failure. *Circulation* 2006;113:1086-92.
- (151) Forstermann U, Hertting G, Neufang B. The importance of endogenous prostaglandins other than prostacyclin, for the modulation of contractility of some rabbit blood vessels. *Br J Pharmacol* 1984;81:623-30.
- (152) Foucart S, Grondin L, Couture R, Nadeau R. Modulation of noradrenaline release by B1 and B2 kinin receptors during metabolic anoxia in the rat isolated atria. *Can J Physiol Pharmacol* 1997;75:639-45.
- (153) Fox A, Kaur S, Li B, Panesar M, Saha U, Davis C, Dragoni I, Colley S, Ritchie T, Bevan S, Burgess G, McIntyre P. Antihyperalgesic activity of a novel nonpeptide bradykinin B1 receptor antagonist in transgenic mice expressing the human B1 receptor. *Br J Pharmacol* 2005;144:889-99.
- (154) Fredrik Leeb-Lundberg LM. Bradykinin specificity and signaling at GPR100 and B2 kinin receptors. *Br J Pharmacol* 2004;143:931-2.
- (155) Fu Y, Katsuya T, Matsuo A, Yamamoto K, Akasaka H, Takami Y, Iwashima Y, Sugimoto K, Ishikawa K, Ohishi M, Rakugi H, Ogihara T. Relationship of bradykinin B2 receptor gene polymorphism with essential hypertension and left ventricular hypertrophy. *Hypertens Res* 2004;27:933-8.
- (156) Fujii M, Wada A, Tsutamoto T, Ohnishi M, Isono T, Kinoshita M. Bradykinin improves left ventricular diastolic function under long-term angiotensin-converting enzyme inhibition in heart failure. *Hypertension* 2002;39:952-7.

- (157) Gabra BH, Sirois P. Role of bradykinin B(1) receptors in diabetes-induced hyperalgesia in streptozotocin-treated mice. *Eur J Pharmacol* 2002;457:115-24.
- (158) Gainer JV, Brown NJ, Bachvarova M, Bastien L, Maltais I, Marceau F, Bachvarov DR. Altered frequency of a promoter polymorphism of the kinin B2 receptor gene in hypertensive African-Americans. *Am J Hypertens* 2000;13:1268-73.
- (159) Gainer JV, Morrow JD, Loveland A, King DJ, Brown NJ. Effect of bradykinin-receptor blockade on the response to angiotensin-converting-enzyme inhibitor in normotensive and hypertensive subjects. *N Engl J Med* 1998;339:1285-92.
- (160) Gallagher AM, Yu H, Printz MP. Bradykinin-induced reductions in collagen gene expression involve prostacyclin. *Hypertension* 1998;32:84-8.
- (161) Garfinkel S, Hu X, Prudovsky IA, McMahon GA, Kapnik EM, McDowell SD, Maciag T. FGF-1-dependent proliferative and migratory responses are impaired in senescent human umbilical vein endothelial cells and correlate with the inability to signal tyrosine phosphorylation of fibroblast growth factor receptor-1 substrates. *J Cell Biol* 1996;134:783-91.
- (162) Ghosh DK, Salerno JC. Nitric oxide synthases: domain structure and alignment in enzyme function and control. *Front Biosci* 2003;8:d193-d209.
- (163) Gobeil F, Pheng LH, Badini I, Nguyen-Le XK, Pizard A, Rizzi A, Blouin D, Regoli D. Receptors for kinins in the human isolated umbilical vein. *Br J Pharmacol* 1996;118:289-94.
- (164) Gohlke P, Kuwer I, Schnell A, Amann K, Mall G, Unger T. Blockade of bradykinin B2 receptors prevents the increase in capillary density induced by chronic angiotensin-converting enzyme inhibitor treatment in stroke-prone spontaneously hypertensive rats. *Hypertension* 1997;29:478-82.
- (165) Golser R, Gorren AC, Leber A, Andrew P, Habisch HJ, Werner ER, Schmidt K, Venema RC, Mayer B. Interaction of endothelial and neuronal nitric-oxide synthases with the bradykinin B2 receptor. Binding of an inhibitory peptide to the oxygenase domain blocks uncoupled NADPH oxidation. *J Biol Chem* 2000;275:5291-6.
- (166) Goto M, Liu Y, Yang XM, Ardell JL, Cohen MV, Downey JM. Role of bradykinin in protection of ischemic preconditioning in rabbit hearts. *Circ Res* 1995;77:611-21.
- (167) Graham DJ, Campen D, Hui R, Spence M, Cheetham C, Levy G, Shoor S, Ray WA. Risk of acute myocardial infarction and sudden cardiac death in patients treated with cyclo-oxygenase 2 selective and non-selective non-steroidal anti-inflammatory drugs: nested case-control study. *Lancet* 2005;365:475-81.
- (168) Greengard JS, Griffin JH. Receptors for high molecular weight kininogen on stimulated washed human platelets. *Biochemistry* 1984;23:6863-9.
- (169) Gregg D, Rauscher FM, Goldschmidt-Clermont PJ. Rac regulates cardiovascular superoxide through diverse molecular interactions: more than a binary GTP switch. *Am J Physiol Cell Physiol* 2003;285:C723-C734.
- (170) Grieve DJ, MacCarthy PA, Gall NP, Cave AC, Shah AM. Divergent biological actions of coronary endothelial nitric oxide during progression of cardiac hypertrophy. *Hypertension* 2001;38:267-73.
- (171) Griffith OW, Stuehr DJ. Nitric oxide synthases: properties and catalytic mechanism. *Annu Rev Physiol* 1995;57:707-36.
- (172) Groves P, Kurz S, Just H, Drexler H. Role of endogenous bradykinin in human coronary vasomotor control. *Circulation* 1995;92:3424-30.
- (173) Haddad EB, Fox AJ, Rousell J, Burgess G, McIntyre P, Barnes PJ, Chung KF. Post-transcriptional regulation of bradykinin B1 and B2 receptor gene expression in human lung fibroblasts by tumor necrosis factor-alpha: modulation by dexamethasone. *Mol Pharmacol* 2000;57:1123-31.
- (174) Hallberg P, Lind L, Michaelsson K, Karlsson J, Kurland L, Kahan T, Malmqvist K, Ohman KP, Nystrom F, Melhus H. B2 bradykinin receptor (B2BKR) polymorphism and change in left ventricular mass in response to antihypertensive treatment: results from the Swedish Irbesartan Left Ventricular Hypertrophy Investigation versus Atenolol (SILVHIA) trial. *J Hypertens* 2003;21:621-4.

-
- (175) Han ED, MacFarlane RC, Mulligan AN, Scafidi J, Davis AE 3rd.. Increased vascular permeability in C1 inhibitor-deficient mice mediated by the bradykinin type 2 receptor. *J Clin Invest* 2002;109:1057-63.
- (176) Harris MB, Ju H, Venema VJ, Liang H, Zou R, Michell BJ, Chen ZP, Kemp BE, Venema RC. Reciprocal phosphorylation and regulation of endothelial nitric-oxide synthase in response to bradykinin stimulation. *J Biol Chem* 2001;276:16587-91.
- (177) Hasan AA, Amenta S, Schmaier AH. Bradykinin and its metabolite, Arg-Pro-Pro-Gly-Phe, are selective inhibitors of alpha-thrombin-induced platelet activation. *Circulation* 1996;94:517-28.
- (178) Hawkinson JE, Szoke BG, Garofalo AW, Hom DS, Zhang H, Dreyer M, Fukuda JY, Chen L, Samant B, Simmonds S, Zeitz KP, Wadsworth A, Liao A, Chavez RA, Zmolek W, Ruslim L, Bova MP, Holcomb R, Butelman ER, Ko MC, Malmberg AB. Pharmacological, pharmacokinetic, and primate analgesic efficacy profile of the novel bradykinin B1 Receptor antagonist ELN441958. *J Pharmacol Exp Ther* 2007;322:619-30.
- (179) Hecquet C, Tan F, Marcic BM, Erdos EG. Human bradykinin B(2) receptor is activated by kallikrein and other serine proteases. *Mol Pharmacol* 2000;58:828-36.
- (180) Heller LJ, Opsahl JA, Wernsing SE, Saxena R, Katz SA. Myocardial and plasma renin-angiotensinogen dynamics during pressure-induced cardiac hypertrophy. *Am J Physiol* 1998;274:R849-R856.
- (181) Helske S, Laine M, Kupari M, Lommi J, Turto H, Nurmi L, Tikkanen I, Werkkala K, Lindstedt KA, Kovanen PT. Increased expression of profibrotic neutral endopeptidase and bradykinin type 1 receptors in stenotic aortic valves. *Eur Heart J* 2007;28:1894-903.
- (182) Henderson LM, Figueroa CD, Muller-Esterl W, Stain A, Bhoola KD. Immunovisualisation of plasma prekallikrein and H-kininogen on human neutrophils and in human hepatocytes. *Agents Actions Suppl* 1992;38:590-4.
- (183) Hess JF, Borkowski JA, Young GS, Strader CD, Ransom RW. Cloning and pharmacological characterization of a human bradykinin (BK-2) receptor. *Biochem Biophys Res Commun* 1992;184:260-8.
- (184) Hilgenfeldt U, Puschner T, Riester U, Finsterle J, Hilgenfeldt J, Ritz E. Low-salt diet downregulates plasma but not tissue kallikrein-kinin system. *Am J Physiol* 1998;275:F88-F93.
- (185) Hirawa N, Uehara Y, Suzuki T, Kawabata Y, Numabe A, Gomi T, Ikeda T, Kizuki K, Omata M. Regression of glomerular injury by kallikrein infusion in Dahl salt-sensitive rats is a bradykinin B2-receptor-mediated event. *Nephron* 1999;81:183-93.
- (186) Hiroki J, Shimokawa H, Higashi M, Morikawa K, Kandabashi T, Kawamura N, Kubota T, Ichiki T, Amano M, Kaibuchi K, Takeshita A. Inflammatory stimuli upregulate Rho-kinase in human coronary vascular smooth muscle cells. *J Mol Cell Cardiol* 2004;37:537-46.
- (187) Hla T, Ristimaki A, Appleby S, Barriocanal JG. Cyclooxygenase gene expression in inflammation and angiogenesis. *Ann N Y Acad Sci* 1993;696:197-204.
- (188) Hock FJ, Wirth K, Albus U, Linz W, Gerhards HJ, Wiemer G, Henke S, Breipohl G, Konig W, Knolle J, et al. Hoe 140 a new potent and long acting bradykinin-antagonist: in vitro studies. *Br J Pharmacol* 1991;102:769-73.
- (189) Hofmann U, Heuer S, Meder K, Boehler J, Lange V, Quaschnig T, Ertl G, Bonz A. The proinflammatory cytokines TNF-alpha and IL-1 beta impair economy of contraction in human myocardium. *Cytokine* 2007;39:157-62.
- (190) Hopkins TA, Ouchi N, Shibata R, Walsh K. Adiponectin actions in the cardiovascular system. *Cardiovasc Res* 2007;74:11-8.
- (191) Horio T, Nishikimi T, Yoshihara F, Nagaya N, Matsuo H, Takishita S, Kangawa K. Production and secretion of adrenomedullin in cultured rat cardiac myocytes and nonmyocytes: stimulation by interleukin-1beta and tumor necrosis factor-alpha. *Endocrinology* 1998;139:4576-80.
- (192) Hornig B, Arakawa N, Drexler H. Effect of ACE inhibition on endothelial dysfunction in patients with chronic heart failure. *Eur Heart J* 1998;19:G48-G53.
- (193) Hornig B, Kohler C, Drexler H. Role of bradykinin in mediating vascular effects of angiotensin-

- converting enzyme inhibitors in humans. *Circulation* 1997;95:1115-8.
- (194) Horwich TB, MacLellan WR, Fonarow GC. Statin therapy is associated with improved survival in ischemic and non-ischemic heart failure. *J Am Coll Cardiol* 2004;43:642-8.
- (195) Houel R, Su J, Barbe F, Choussat R, Crozatier B, Hittinger L. Regional myocardial effects of intracoronary bradykinin in conscious dogs. *Am J Physiol* 1997;72:H1266-H1274.
- (196) Houle S, Landry M, Audet R, Bouthillier J, Bachvarov DR, Marceau F. Effect of allelic polymorphism of the B(1) and B(2) receptor genes on the contractile responses of the human umbilical vein to kinins. *J Pharmacol Exp Ther* 2000;294:45-51.
- (197) Houle S, Molinaro G, Adam A, Marceau F. Tissue kallikrein actions at the rabbit natural or recombinant kinin B2 receptors. *Hypertension* 2003;1:611-7.
- (198) Huang A, Sun D, Carroll MA, Jiang H, Smith CJ, Connetta JA, Falck JR, Shesely EG, Koller A, Kaley G. EDHF mediates flow-induced dilation in skeletal muscle arterioles of female eNOS-KO mice. *Am J Physiol Heart Circ Physiol* 2001;280:H2462-H2469.
- (199) Huang TJ, Haddad EB, Fox AJ, Salmon M, Jones C, Burgess G, Chung KF. Contribution of bradykinin B(1) and B(2) receptors in allergen-induced bronchial hyperresponsiveness. *Am J Respir Crit Care Med* 1999;160:1717-23.
- (200) Ignarro LJ, Buga GM, Wood KS, Byrns RE, Chaudhuri G. Endothelium-derived relaxing factor produced and released from artery and vein is nitric oxide. *Proc Natl Acad Sci U S A* 1987;84:9265-9.
- (201) Ignjatovic T, Tan F, Brovkovich V, Skidgel RA, Erdos EG. Activation of bradykinin B1 receptor by ACE inhibitors. *Int Immunopharmacol* 2002;2:1787-93.
- (202) Ignjatovic T, Tan F, Brovkovich V, Skidgel RA, Erdos EG. Novel mode of action of angiotensin I converting enzyme inhibitors: direct activation of bradykinin B1 receptor. *J Biol Chem* 2002;277:16847-52.
- (203) Inagami T. Atrial natriuretic factor as a volume regulator. *J Clin Pharmacol* 1994;34:424-6.
- (204) Inagami T, Naruse M, Hoover R. Endothelium as an endocrine organ. *Annu Rev Physiol* 1995;57:171-89.
- (205) Innis RB, Manning DC, Stewart JM, Snyder SH. [3H]Bradykinin receptor binding in mammalian tissue membranes. *Proc Natl Acad Sci U S A* 1981;78:2630-4.
- (206) Isami S, Kishikawa H, Araki E, Uehara M, Kaneko K, Shirotani T, Todaka M, Ura S, Motoyoshi S, Matsumoto K, Miyamura N, Shichiri M. Bradykinin enhances GLUT4 translocation through the increase of insulin receptor tyrosine kinase in primary adipocytes: evidence that bradykinin stimulates the insulin signalling pathway. *Diabetologia* 1996;39:412-20.
- (207) Ishigai Y, Mori T, Ikeda T, Fukuzawa A, Shibano T. Role of bradykinin-NO pathway in prevention of cardiac hypertrophy by ACE inhibitor in rat cardiomyocytes. *Am J Physiol* 1997;273:H2659-H2663.
- (208) Jaffa AA, Rust PF, Mayfield RK. Kinin, a mediator of diabetes-induced glomerular hyperfiltration. *Diabetes* 1995;4:156-60.
- (209) Jin LS, Seeds E, Page CP, Schachter M. Inhibition of bradykinin-induced bronchoconstriction in the guinea-pig by a synthetic B2 receptor antagonist. *Br J Pharmacol* 1989;97:598-602.
- (210) Jong YJ, Dalem LR, Wilhelm B, Baenziger NL. Human bradykinin B2 receptors isolated by receptor-specific monoclonal antibodies are tyrosine phosphorylated. *Proc Natl Acad Sci USA* 1993;90:10994-8.
- (211) Ju H, Venema VJ, Liang H, Harris MB, Zou R, Venema RC. Bradykinin activates the Janus-activated kinase/signal transducers and activators of transcription (JAK/STAT) pathway in vascular endothelial cells: localization of JAK/STAT signalling proteins in plasmalemmal caveolae. *Biochem J* 2000;351:257-64.
- (212) Kakoki M, Kizer CM, Yi X, Takahashi N, Kim HS, Bagnell CR, Edgell CJ, Maeda N, Jennette JC, Smithies O. Senescence-associated phenotypes in Akita diabetic mice are enhanced by absence of bradykinin B2 receptors. *J Clin Invest* 2006;116:1302-9.
- (213) Kakoki M, Takahashi N, Jennette JC, Smithies O. Diabetic nephropathy is markedly enhanced

- in mice lacking the bradykinin B2 receptor. *Proc Natl Acad Sci U S A* 2004;101:13302-5.
- (214) Kallio PJ, Pongratz I, Gradin K, McGuire J, Poellinger L. Activation of hypoxia-inducible factor 1alpha: posttranscriptional regulation and conformational change by recruitment of the Arnt transcription factor. *Proc Natl Acad Sci U S A* 1997;94:5667-72.
- (215) Kammerer S, Braun A, Arnold N, Roscher AA. The human bradykinin B2 receptor gene: full length cDNA, genomic organization and identification of the regulatory region. *Biochem Biophys Res Commun* 1995;211:226-33.
- (216) Kang DS, Ryberg K, Morgelin M, Leeb-Lundberg LM. Spontaneous formation of a proteolytic B1 and B2 bradykinin receptor complex with enhanced signaling capacity. *J Biol Chem* 2004;279:22102-7.
- (217) Kaplan AP, Silverberg M. The coagulation-kinin pathway of human plasma. *Blood* 198;70:1-15.
- (218) Katsumata N, Shimokawa H, Seto M, Kozai T, Yamawaki T, Kuwata K, Egashira K, Ikegaki I, Asano T, Sasaki Y, Takeshita A. Enhanced myosin light chain phosphorylations as a central mechanism for coronary artery spasm in a swine model with interleukin-1beta. *Circulation* 1997;96:4357-63.
- (219) Katz AM. Heart failure: pathophysiology, molecular biology and clinical management. Lippincott/Williams&Wilkins, Philadelphia; 2000.
- (220) Katz AM. Heart failure: a hemodynamic disorder complicated by maladaptive proliferative responses. *Journal of Cellular and Molecular Medicine* 2003;7:1-10.
- (221) Katz AM. Proliferative signaling and disease progression in heart failure. *Circulation Journal* 2002;66:225-31.
- (222) Katz SD, Biasucci L, Sabba C, Strom JA, Jondeau G, Galvao M, Solomon S, Nikolic SD, Forman R, LeJemtel TH. Impaired endothelium-mediated vasodilation in the peripheral vasculature of patients with congestive heart failure. *J Am Coll Cardiol* 1992;19:918-25.
- (223) Ke Q, Costa M. Hypoxia-inducible factor-1 (HIF-1). *Mol Pharmacol* 2006;70:1469-80.
- (224) Kessler P, Popp R, Busse R, Schini-Kerth VB. Proinflammatory mediators chronically downregulate the formation of the endothelium-derived hyperpolarizing factor in arteries via a nitric oxide/cyclic GMP-dependent mechanism. *Circulation* 1999;99:1878-84.
- (225) Kintsurashvili E, Duka A, Ignjacev I, Pattakos G, Gavras I, Gavras H. Age-related changes of bradykinin B1 and B2 receptors in rat heart. *Am J Physiol Heart Circ Physiol* 2005;289:H202-H205.
- (226) Kitakaze M, Node K, Minamino T, Asanuma H, Ueda Y, Kosaka H, Kuzuya T, Hori M. Inhibition of angiotensin-converting enzyme increases the nitric oxide levels in canine ischemic myocardium. *J Mol Cell Cardiol* 1998;30:2461-6.
- (227) Kitamura K, Kangawa K, Kawamoto M, Ichiki Y, Nakamura S, Matsuo H, Eto T. Adrenomedullin: a novel hypotensive peptide isolated from human pheochromocytoma. *Biochem Biophys Res Commun* 1993;192:553-60.
- (228) Kjekshus J, Dunselman P, Blideskog M, Eskilson C, Hjalmarson A, McMurray JV, Waagstein F, Wedel H, Wessman P, Wikstrand J. A statin in the treatment of heart failure? Controlled rosuvastatin multinational study in heart failure (CORONA): study design and baseline characteristics. *Eur J Heart Fail* 2005;7:1059-69.
- (229) Kjekshus J, Pedersen TR, Olsson AG, Faergeman O, Pyorala K. The effects of simvastatin on the incidence of heart failure in patients with coronary heart disease. *J Card Fail* 1997;3:249-54.
- (230) Kleine L, Liu G, Leblanc N, Hebert RL. Bradykinin stimulates ceramide production by activating specific BK-B(1) receptor in rat small artery. *Am J Physiol Heart Circ Physiol* 2002;282:H175-H183.
- (231) Knox AJ, Corbett L, Stocks J, Holland E, Zhu YM, Pang L. Human airway smooth muscle cells secrete vascular endothelial growth factor: up-regulation by bradykinin via a protein kinase C and prostanoïd-dependent mechanism. *FASEB J* 2001;15:2480-8.

- (232) Kobayashi N, Higashi T, Hara K, Shirataki H, Matsuoka H. Effects of imidapril on NOS expression and myocardial remodelling in failing heart of Dahl salt-sensitive hypertensive rats. *Cardiovasc Res* 1999;44:518-26.
- (233) Kokkonen JO, Kuoppala A, Saarinen J, Lindstedt KA, Kovanen PT. Kallidin- and bradykinin-degrading pathways in human heart: degradation of kallidin by aminopeptidase M-like activity and bradykinin by neutral endopeptidase. *Circulation* 1999;99:1984-90.
- (234) Kokkonen JO, Lindstedt KA, Kuoppala A, Kovanen PT. Kinin-degrading pathways in the human heart. *Trends Cardiovasc Med* 2000;10:42-5.
- (235) Kositprapa C, Ockaili RA, Kukreja RC. Bradykinin B2 receptor is involved in the late phase of preconditioning in rabbit heart. *J Mol Cell Cardiol* 2001;33:1355-62.
- (236) Kozik A, Moore RB, Potempa J, Imamura T, Rapala-Kozik M, Travis J. A novel mechanism for bradykinin production at inflammatory sites. Diverse effects of a mixture of neutrophil elastase and mast cell tryptase versus tissue and plasma kallikreins on native and oxidized kininogens. *J Biol Chem* 1998;273:33224-9.
- (237) Kubo SH, Rector TS, Bank AJ, Williams RE, Heifetz SM. Endothelium-dependent vasodilation is attenuated in patients with heart failure. *Circulation* 1991;84:1589-96.
- (238) Kuga T, Egashira K, Mohri M, Tsutsui H, Harasawa Y, Urabe Y, Ando S, Shimokawa H, Takeshita A. Bradykinin-induced vasodilation is impaired at the atherosclerotic site but is preserved at the spastic site of human coronary arteries in vivo. *Circulation* 1995;92:183-9.
- (239) Kuoppala A, Shiota N, Kokkonen JO, Liesmaa I, Kostner K, Mayranpaa M, Kovanen PT, Lindstedt KA. Down-regulation of cardioprotective bradykinin type-2 receptors in the left ventricle of patients with end-stage heart failure. *J Am Coll Cardiol* 2002;40:119-25.
- (240) Kuoppala A, Shiota N, Lindstedt KA, Rysa J, Leskinen HK, Luodonpaa M, Liesmaa I, Ruskoaho H, Kaaja R, Kovanen PT, Kokkonen JO. Expression of bradykinin receptors in the left ventricles of rats with pressure overload hypertrophy and heart failure. *J Hypertens* 2003;21:1729-36.
- (241) Kuruvilla L, Kartha CC. Molecular mechanisms in endothelial regulation of cardiac function. *Mol Cell Biochem* 2003;253:113-23.
- (242) Lagneux C, Bader M, Pesquero JB, Demege P, Ribaut C. Detrimental implication of B1 receptors in myocardial ischemia: evidence from pharmacological blockade and gene knockout mice. *Int Immunopharmacol* 2002;2:815-22.
- (243) Lamba S, Abraham WT. Alterations in adrenergic receptor signaling in heart failure. *Heart Fail Rev* 2000;5:7-16.
- (244) Landmesser U, Bahlmann F, Mueller M, Spiekermann S, Kirchhoff N, Schulz S, Manes C, Fischer D, de Groot K, Fliser D, Fauler G, Marz W, Drexler H. Simvastatin versus ezetimibe: pleiotropic and lipid-lowering effects on endothelial function in humans. *Circulation* 2005;111:2356-63.
- (245) Lang NN, Cruden NL, Tse GH, Bloomfield P, Ludlam CA, Fox KA, Newby DE. Vascular B1 kinin receptors in patients with congestive heart failure. *J Cardiovasc Pharmacol* 2008;52:438-44.
- (246) Laufs U, Endres M, Custodis F, Gertz K, Nickenig G, Liao JK, Bohm M. Suppression of endothelial nitric oxide production after withdrawal of statin treatment is mediated by negative feedback regulation of rho GTPase gene transcription. *Circulation* 2000;102:3104-10.
- (247) Laufs U, Fata VL, Liao JK. Inhibition of 3-hydroxy-3-methylglutaryl (HMG)-CoA reductase blocks hypoxia-mediated down-regulation of endothelial nitric oxide synthase. *J Biol Chem* 1997;272:31725-9.
- (248) Laufs U, Wassmann S, Schackmann S, Heeschen C, Bohm M, Nickenig G. Beneficial effects of statins in patients with non-ischemic heart failure. *Z Kardiol* 2004;93:103-8.
- (249) Lauton-Santos S, Guatimosim S, Castro CH, Oliveira FA, Almeida AP, as-Peixoto MF, Gomes MA, Pessoa P, Pesquero JL, Pesquero JB, Bader M, Cruz JS. Kinin B1 receptor participates in the control of cardiac function in mice. *Life Sciences* 2007;81:814-22.

-
- (250) Lee SH, Wolf PL, Escudero R, Deutsch R, Jamieson SW, Thistlethwaite PA. Early expression of angiogenesis factors in acute myocardial ischemia and infarction. *N Engl J Med* 2000;342:626-33.
- (251) Leeb-Lundberg LM, Mathis SA. Guanine nucleotide regulation of B2 kinin receptors. Time-dependent formation of a guanine nucleotide-sensitive receptor state from which [3H]bradykinin dissociates slowly. *J Biol Chem* 1990;265:9621-7.
- (252) Leesar MA, Stoddard MF, Manchikalapudi S, Bolli R. Bradykinin-induced preconditioning in patients undergoing coronary angioplasty. *J Am Coll Cardiol* 1999;34:639-50.
- (253) Leskinen H, Rauma-Pinola T, Szokodi I, Kerkela R, Pikkariainen S, Uusimaa P, Hautala T, Vuolteenaho O, Ruskoaho H. Adaptive or maladaptive response to adenoviral adrenomedullin gene transfer is context-dependent in the heart. *J Gene Med* 2008;10:867-77.
- (254) Levesque L, Drapeau G, Grose JH, Rioux F, Marceau F. Vascular mode of action of kinin B1 receptors and development of a cellular model for the investigation of these receptors. *Br J Pharmacol* 1993;109:1254-62.
- (255) Levesque L, Larrivee JF, Bachvarov DR, Rioux F, Drapeau G, Marceau F. Regulation of kinin-induced contraction and DNA synthesis by inflammatory cytokines in the smooth muscle of the rabbit aorta. *Br J Pharmacol* 1995;116:1673-9.
- (256) Levine B, Kalman J, Mayer L, Fillit HM, Packer M. Elevated circulating levels of tumor necrosis factor in severe chronic heart failure. *N Engl J Med* 1990;323:236-41.
- (257) Liao JK, Homcy CJ. The G proteins of the G alpha i and G alpha q family couple the bradykinin receptor to the release of endothelium-derived relaxing factor. *J Clin Invest* 1993;92:2168-72.
- (258) Liao Z, Brar BK, Cai Q, Stephanou A, O'Leary RM, Pennica D, Yellon DM, Latchman DS. Cardiotrophin-1 (CT-1) can protect the adult heart from injury when added both prior to ischaemia and at reperfusion. *Cardiovasc Res* 2002;53:902-10.
- (259) Libby P, Aikawa M, Kinlay S, Selwyn A, Ganz P. Lipid lowering improves endothelial functions. *Int J Cardiol* 2000;74:S3-S10.
- (260) Liesmaa I, Kuoppala A, Shiota N, Kokkonen JO, Kostner K, Mayranpaa M, Kovanen PT, Lindstedt KA. Increased expression of bradykinin type-1 receptors in endothelium of intramyocardial coronary vessels in human failing hearts. *Am J Physiol Heart Circ Physiol* 2005;288:H2317-H2322.
- (261) Lindstedt KA, Mahon MG, Foisner R, Hermann M, Nimpf J, Schneider WJ. Receptor-associated protein in an oviparous species is correlated with the expression of a receptor variant. *J Biol Chem* 1997;272:30221-7.
- (262) Linz W, Scholkens BA, Han YF. Beneficial effects of the converting enzyme inhibitor, ramipril, in ischemic rat hearts. *J Cardiovasc Pharmacol* 1986;8:S91-S99.
- (263) Linz W, Wiemer G, Schaper J, Zimmermann R, Nagasawa K, Gohlke P, Unger T, Scholkens BA. Angiotensin converting enzyme inhibitors, left ventricular hypertrophy and fibrosis. *Mol Cell Biochem* 1995;147:89-97.
- (264) Lisanti MP, Scherer PE, Tang Z, Sargiacomo M. Caveolae, caveolin and caveolin-rich membrane domains: a signalling hypothesis. *Trends in Cell Biology* 1994;4:231-5.
- (265) Liu C, Chen J, Sutton S, Roland B, Kuei C, Farmer N, Sillard R, Lovenberg TW. Identification of relaxin-3/INSL7 as a ligand for GPCR142. *J Biol Chem* 2003;278:50765-70.
- (266) Liu YH, Yang XP, Sharov VG, Nass O, Sabbah HN, Peterson E, Carretero OA. Effects of angiotensin-converting enzyme inhibitors and angiotensin II type 1 receptor antagonists in rats with heart failure. Role of kinins and angiotensin II type 2 receptors. *J Clin Invest* 1997;99:1926-35.
- (267) Lohmeier TE, Mizelle HL, Reinhart GA, Montani JP, Hord CE, Jr., Didlake RH. Atrial natriuretic peptide and sodium homeostasis in compensated heart failure. *Am J Physiol* 1996; 271:R1353-R1363.
- (268) Loot AE, Roks AJ, Henning RH, Tio RA, Suurmeijer AJ, Boomsma F, van Gilst WH. Angiotensin-(1-7) attenuates the development of heart failure after myocardial infarction in rats. *Circulation* 2002;105:1548-50.

- (269) Lovenberg TW, Liaw CW, Grigoriadis DE, Clevenger W, Chalmers DT, De Souza EB, Oltersdorf T. Cloning and characterization of a functionally distinct corticotropin-releasing factor receptor subtype from rat brain. *Proc Natl Acad Sci U S A* 1995;92:836-40.
- (270) Lung CC, Jagels MA, Daffern PJ, Tan EM, Zuraw BL. Induction of human B2 bradykinin receptor mRNA and membrane receptors by IFN γ . *Immunopharmacology* 1998;39: 243-53.
- (271) Ma JX, Wang DZ, Ward DC, Chen L, Dessai T, Chao J, Chao L. Structure and chromosomal localization of the gene (BDKRB2) encoding human bradykinin B2 receptor. *Genomics* 1994; 23:362-9.
- (272) MacCarthy PA, Grocott-Mason R, Prendergast BD, Shah AM. Contrasting inotropic effects of endogenous endothelin in the normal and failing human heart: studies with an intracoronary ET(A) receptor antagonist. *Circulation* 2000;101:142-7.
- (273) MacDonald RJ, Margolius HS, Erdos EG. Molecular biology of tissue kallikrein. *Biochem J* 1988;253:313-21.
- (274) Madeddu P, Pargaglia PP, Demontis MP, Varoni MV, Fattaccio MC, Anania V, Glorioso N. Early blockade of bradykinin B2-receptors alters the adult cardiovascular phenotype in rats. *Hypertension* 1995;25:453-9.
- (275) Maestri R, Milia AF, Salis MB, Graiani G, Lagrasta C, Monica M, Corradi D, Emanuelli C, Madeddu P. Cardiac hypertrophy and microvascular deficit in kinin B2 receptor knockout mice. *Hypertension* 2003;41:1151-5.
- (276) Mallat Z, Gojova A, Sauzeau V, Brun V, Silvestre JS, Esposito B, Merval R, Groux H, Loirand G, Tedgui A. Rho-associated protein kinase contributes to early atherosclerotic lesion formation in mice. *Circ Res* 2003;93:884-8.
- (277) Manalo DJ, Rowan A, Lavoie T, Natarajan L, Kelly BD, Ye SQ, Garcia JGN, Semenza GL. Transcriptional regulation of vascular endothelial cell responses to hypoxia by HIF-1. *Blood* 2005;105:659-69.
- (278) Mancini GB, Henry GC, Macaya C, O'Neill BJ, Pucillo AL, Carere RG, Wargovich TJ, Mudra H, Luscher TF, Klibaner MI, Haber HE, Uprichard AC, Pepine CJ, Pitt B. Angiotensin-converting enzyme inhibition with quinapril improves endothelial vasomotor dysfunction in patients with coronary artery disease. The TREND (Trial on Reversing ENdothelial Dysfunction) Study. *Circulation* 1996;94:258-65.
- (279) Manning DC, Snyder SH. Bradykinin receptors localized by quantitative autoradiography in kidney, ureter, and bladder. *Am J Physiol* 1989;256:F909-F915.
- (280) Mantovani A, Bussolino F, Dejana E. Cytokine regulation of endothelial cell function. *FASEB J* 1992;6:2591-9.
- (281) Marceau F, Larrivee JF, Bouthillier J, Bachvarova M, Houle S, Bachvarov DR. Effect of endogenous kinins, prostanoids, and NO on kinin B1 and B2 receptor expression in the rabbit. *Am J Physiol* 1999;277:R1568-R1578.
- (282) Marchetti J, Helou CM, Chollet C, Rajerison R, Alhenc-Gelas F. ACE and non-ACE mediated effect of angiotensin I on intracellular calcium mobilization in rat glomerular arterioles. *Am J Physiol Heart Circ Physiol* 2003;284:H1933-H1941.
- (283) Marcic B, Deddish PA, Jackman HL, Erdos EG. Enhancement of bradykinin and resensitization of its B2 receptor. *Hypertension* 1999;33:835-43.
- (284) Marcic B, Deddish PA, Skidgel RA, Erdos EG, Minshall RD, Tan F. Replacement of the transmembrane anchor in angiotensin I-converting enzyme (ACE) with a glycosylphosphatidylinositol tail affects activation of the B2 bradykinin receptor by ACE inhibitors. *J Biol Chem* 2000;275:16110-8.
- (285) Marin Castano ME, Schanstra JP, Hirtz C, Pesquero JB, Pecher C, Girolami JP, Bascands JL. B2 kinin receptor upregulation by cAMP is associated with BK-induced PGE2 production in rat mesangial cells. *Am J Physiol* 1998;274:F532-F540.
- (286) Marin-Castano ME, Schanstra JP, Neau E, Pradaude F, Pecher C, Ader JL, Girolami JP, Bascands JL. Induction of functional bradykinin b(1)-receptors in normotensive rats and mice

- under chronic angiotensin-converting enzyme inhibitor treatment. *Circulation* 2002;105:627-32.
- (287) Marsault R, Illiano S, Vanhoutte PM. Bradykinin-induced contractions of canine saphenous veins: mediation by B2 receptors and involvement of eicosanoids. *Br J Pharmacol* 1997; 120:215-20.
 - (288) Massaro M, Zampolli A, Scoditti E, Carluccio MA, Storelli C, Distante A, De Caterina R. Statins inhibit cyclooxygenase-2 and matrix metalloproteinase-9 in human endothelial cells: anti-angiogenic actions possibly contributing to plaque stability. *Cardiovasc Res* 2009, Epub.
 - (289) Mathis SA, Criscimagna NL, Leeb-Lundberg LM. B1 and B2 kinin receptors mediate distinct patterns of intracellular Ca²⁺ signaling in single cultured vascular smooth muscle cells. *Mol Pharmacol* 1996;50:128-39.
 - (290) Matsuo S, Matsumoto T, Takashima H, Ohira N, Yamane T, Yasuda Y, Tarutani Y, Horie M. The relationship between flow-mediated brachial artery vasodilation and coronary vasomotor responses to bradykinin: comparison with those to acetylcholine. *J Cardiovasc Pharmacol* 2004;44:164-70.
 - (291) Mattera R, Hayek S, Summers BA, Grove DL. Agonist-specific alterations in receptor-phospholipase coupling following inactivation of Gi2alpha gene. *Biochem J* 1998;332:263-71.
 - (292) Mazenot C, Loufrani L, Henrion D, Ribaut C, Muller-Esterl W, Godin-Ribuot D. Endothelial kinin B(1)-receptors are induced by myocardial ischaemia-reperfusion in the rabbit. *J Physiol* 2001;530:69-78.
 - (293) McAllister BS, Leeb-Lundberg LM, Javors MA, Olson MS. Bradykinin receptors and signal transduction pathways in human fibroblasts: integral role for extracellular calcium. *Arch Biochem Biophys* 1993;304:294-301.
 - (294) McCarthy DA, Potter DE, Nicolaides ED. An in vivo estimation of the potencies and half-lives of synthetic bradykinin and kallidin. *J Pharmacol Exp Ther* 1965;148:117-22.
 - (295) McEachern AE, Shelton ER, Bhakta S, Obernolte R, Bach C, Zuppan P, Fujisaki J, Aldrich RW, Jarnagin K. Expression cloning of a rat B2 bradykinin receptor. *Proc Natl Acad Sci USA* 1991;88:7724-8.
 - (296) McLean PG, Ahluwalia A, Perretti M. Association between kinin B(1) receptor expression and leukocyte trafficking across mouse mesenteric postcapillary venules. *J Exp Med* 2000; 192:367-80.
 - (297) McLean PG, Perretti M, Ahluwalia A. Inducible expression of the kinin B1 receptor in the endotoxemic heart: mechanisms of des-Arg9bradykinin-induced coronary vasodilation. *Br J Pharmacol* 1999;128:275-82.
 - (298) McLean PG, Perretti M, Ahluwalia A. Kinin B(1) receptors and the cardiovascular system: regulation of expression and function. *Cardiovasc Res* 2000;48:194-210.
 - (299) McMurray JJ, Pfeffer MA. Heart failure. *Lancet* 2005;365:1877-89.
 - (300) McMurray JJ, Ray SG, Abdullah I, Dargie HJ, Morton JJ. Plasma endothelin in chronic heart failure. *Circulation* 1992;85:1374-9.
 - (301) McQuillan LP, Leung GK, Marsden PA, Kostyk SK, Kourembanas S. Hypoxia inhibits expression of eNOS via transcriptional and posttranscriptional mechanisms. *Am J Physiol* 1994;267:H1921-H1927.
 - (302) Meneton P, Bloch-Faure M, Hagege AA, Ruetten H, Huang W, Bergaya S, Ceiler D, Gehring D, Martins I, Salmon G, Boulanger CM, Nussberger J, Crozatier B, Gasc JM, Heudes D, Bruneval P, Doetschman T, Menard J, henc-Gelas F. Cardiovascular abnormalities with normal blood pressure in tissue kallikrein-deficient mice. *Proc Natl Acad Sci U S A* 2001; 98:2634-9.
 - (303) Menke JG, Borkowski JA, Bierilo KK, MacNeil T, Derrick AW, Schneck KA, Ransom RW, Strader CD, Linemeyer DL, Hess JF. Expression cloning of a human B1 bradykinin receptor. *J Biol Chem* 1994;269:21583-6.
 - (304) Messadi-Laribi E, Griol-Charhbili V, Pizard A, Vincent MP, Heudes D, Meneton P, henc-Gelas F, Richer C. Tissue kallikrein is involved in the cardioprotective effect of AT1-receptor blockade

- in acute myocardial ischemia. *J Pharmacol Exp Ther* 2007;323:210-6.
- (305) Messerli FH, Ketelhut R. Left ventricular hypertrophy: an independent risk factor. *J Cardiovasc Pharmacol* 1991;17:S59-S66.
- (306) Michiels C, Arnould T, Knott I, Dieu M, Remacle J. Stimulation of prostaglandin synthesis by human endothelial cells exposed to hypoxia. *Am J Physiol* 1993;264:C866-C874.
- (307) Minshall RD, Nakamura F, Becker RP, Rabito SF. Characterization of bradykinin B2 receptors in adult myocardium and neonatal rat cardiomyocytes. *Circ Res* 1995;76:773-80.
- (308) Minshall RD, Nedumgottil SJ, Igic R, Erdos EG, Rabito SF. Potentiation of the effects of bradykinin on its receptor in the isolated guinea pig ileum. *Peptides* 2000;21:1257-64.
- (309) Minshall RD, Tan F, Nakamura F, Rabito SF, Becker RP, Marcic B, Erdos EG. Potentiation of the actions of bradykinin by angiotensin I-converting enzyme inhibitors. The role of expressed human bradykinin B2 receptors and angiotensin I-converting enzyme in CHO cells. *Circ Res* 1997;81:848-56.
- (310) Miura Si, Matsuo Y, Saku K. Transactivation of KDR/Flk-1 by the B2 Receptor Induces Tube Formation in Human Coronary Endothelial Cells. *Hypertension* 2003;41:1118-23.
- (311) Miyamoto A, Laufs U, Pardo C, Liao JK. Modulation of bradykinin receptor ligand binding affinity and its coupled G-proteins by nitric oxide. *J Biol Chem* 1997;272:19601-8.
- (312) Miyauchi T, Masaki T. Pathophysiology of endothelin in the cardiovascular system. *Annu Rev Physiol* 1999;61:391-415.
- (313) Mombouli JV, Illiano S, Nagao T, Scott-Burden T, Vanhoutte PM. Potentiation of endothelium-dependent relaxations to bradykinin by angiotensin I converting enzyme inhibitors in canine coronary artery involves both endothelium-derived relaxing and hyperpolarizing factors. *Circ Res* 1992;71:137-44.
- (314) Mombouli JV, Vanhoutte PM. Kinins and endothelial control of vascular smooth muscle. *Annu Rev Pharmacol Toxicol* 1995;35:679-705.
- (315) Moncada S, Palmer RM, Higgs EA. Nitric oxide: physiology, pathophysiology, and pharmacology. *Pharmacol Rev* 1991;43:109-42.
- (316) Moncada S, Vane JR. Pharmacology and endogenous roles of prostaglandin endoperoxides, thromboxane A2, and prostacyclin. *Pharmacol Rev* 1978;30:293-331.
- (317) Morbidelli L, Parenti A, Giovannelli L, Granger HJ, Ledda F, Ziche M. B1 receptor involvement in the effect of bradykinin on venular endothelial cell proliferation and potentiation of FGF-2 effects. *Br J Pharmacol* 1998;124:1286-92.
- (318) Motta G, Rojkaer R, Hasan AA, Cines DB, Schmaier AH. High molecular weight kininogen regulates prekallikrein assembly and activation on endothelial cells: a novel mechanism for contact activation. *Blood* 1998;91:516-28.
- (319) Mukae S, Itoh S, Aoki S, Iwata T, Nishio K, Sato R, Katagiri T. Association of polymorphisms of the renin-angiotensin system and bradykinin B2 receptor with ACE-inhibitor-related cough. *J Hum Hypertens* 2002;16:857-63.
- (320) Mukoyama M, Nakao K, Obata K, Jougasaki M, Yoshimura M, Morita E, Hosoda K, Suga S, Ogawa Y, Yasue H, . Augmented secretion of brain natriuretic peptide in acute myocardial infarction. *Biochem Biophys Res Commun* 1991;180:431-6.
- (321) Muller-Esterl W. Kininogens, kinins and kinships. *Thromb Haemost* 1989;61:2-6.
- (322) Munk VC, Sanchez de ML, Petrampil M, Butz N, Banfi A, Eriksson U, Hein L, Humar R, Battegay EJ. Angiotensin II induces angiogenesis in the hypoxic adult mouse heart in vitro through an AT2-B2 receptor pathway. *Hypertension* 2007;49:1178-85.
- (323) Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation* 1986;74:1124-36.
- (324) Nagaya N, Nishikimi T, Horio T, Yoshihara F, Kanazawa A, Matsuo H, Kangawa K. Cardiovascular and renal effects of adrenomedullin in rats with heart failure. *Am J Physiol* 1999;276:R213-R218.

-
- (325) Nakano A, Miura T, Miki T, Nozawa Y, Ichikawa Y, Ura N, Shimamoto K. Effects of neutral endopeptidase 24.11 inhibition on myocardial infarct size and ischemic preconditioning in rabbits. *Naunyn Schmiedebergs Arch Pharmacol* 2002;366:335-42.
- (326) Neer EJ, Smith TF. G protein heterodimers: new structures propel new questions. *Cell* 1996; 84:175-8.
- (327) Newton R, Eddleston J, Haddad E, Hawisa S, Mak J, Lim S, Fox AJ, Donnelly LE, Chung KF. Regulation of kinin receptors in airway epithelial cells by inflammatory cytokines and dexamethasone. *Eur J Pharmacol* 2002;441:193-202.
- (328) Ni A, Chai KX, Chao L, Chao J. Molecular cloning and expression of rat bradykinin B1 receptor. *Biochim Biophys Acta* 1998;1442:177-85.
- (329) Ni A, Yin H, Agata J, Yang Z, Chao L, Chao J. Overexpression of kinin B1 receptors induces hypertensive response to des-Arg9-bradykinin and susceptibility to inflammation. *J Biol Chem* 2003;278:219-25.
- (330) Nittynen L, Nurminen ML, Korpela R, Vapaatalo H. Role of arginine, taurine and homocysteine in cardiovascular diseases. *Ann Med* 1999;31:318-26.
- (331) Node K, Fujita M, Kitakaze M, Hori M, Liao JK. Short-term statin therapy improves cardiac function and symptoms in patients with idiopathic dilated cardiomyopathy. *Circulation* 2003; 108:839-43.
- (332) Nohria A, Grunert ME, Rikitake Y, Noma K, Prsic A, Ganz P, Liao JK, Creager MA. Rho kinase inhibition improves endothelial function in human subjects with coronary artery disease. *Circ Res* 2006;99:1426-32.
- (333) Noma K, Goto C, Nishioka K, Jitsuiki D, Umemura T, Ueda K, Kimura M, Nakagawa K, Oshima T, Chayama K, Yoshizumi M, Liao JK, Higashi Y. Roles of rho-associated kinase and oxidative stress in the pathogenesis of aortic stiffness. *J Am Coll Cardiol* 2007;49:698-705.
- (334) Nussmeier NA, Whelton AA, Brown MT, Langford RM, Hoeft A, Parlow JL, Boyce SW, Verbarg KM. Complications of the COX-2 inhibitors parecoxib and valdecoxib after cardiac surgery. *N Engl J Med* 2005;352:1081-91.
- (335) O'Murchu B, Miller VM, Perrella MA, Burnett JC, Jr. Increased production of nitric oxide in coronary arteries during congestive heart failure. *J Clin Invest* 1994;93:165-71.
- (336) Oeseburg H, Iusuf D, van der HP, van Gilst WH, Henning RH, Roks AJ. Bradykinin protects against oxidative stress-induced endothelial cell senescence. *Hypertension* 2009;53:417-22.
- (337) Orawski AT, Simmons WH. Purification and properties of membrane-bound aminopeptidase P from rat lung. *Biochemistry* 1995;34:11227-36.
- (338) Orton EC, Reeves JT, Stenmark KR. Pulmonary vasodilation with structurally altered pulmonary vessels and pulmonary hypertension. *J Appl Physiol* 1988;65:2459-67.
- (339) Osorio JC, Cheema FH, Martens TP, Mahmut N, Kinnear C, Gonzalez AM, Bonney W, Homma S, Liao JK, Mital S. Simvastatin reverses cardiac hypertrophy caused by disruption of the bradykinin 2 receptor. *Can J Physiol Pharmacol* 2008;86:633-42.
- (340) Ouchi N, Shibata R, Walsh K. Cardioprotection by adiponectin. *Trends Cardiovasc Med* 2006; 16:141-6.
- (341) Pagliaro P, Penna C. Rethinking the renin-angiotensin system and its role in cardiovascular regulation. *Cardiovasc Drugs Ther* 2005;19:77-87.
- (342) Palmer LA, Semenza GL, Stoler MH, Johns RA. Hypoxia induces type II NOS gene expression in pulmonary artery endothelial cells via HIF-1. *Am J Physiol* 1998;274:L212-L219.
- (343) Palmer RM, Ferrige AG, Moncada S. Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* 1987;327:524-6.
- (344) Pang L, Knox AJ. PGE2 release by bradykinin in human airway smooth muscle cells: involvement of cyclooxygenase-2 induction. *Am J Physiol* 1997;273:L1132-L1140.
- (345) Parenti A, Morbidelli L, Ledda F, Granger HJ, Ziche M. The bradykinin/B1 receptor promotes angiogenesis by up-regulation of endogenous FGF-2 in endothelium via the nitric oxide synthase pathway. *FASEB J* 2001;15:1487-9.

- (346) Passos GF, Fernandes ES, Campos MM, Araujo JG, Pesquero JL, Souza GE, Avellar MC, Teixeira MM, Calixto JB. Kinin B1 receptor up-regulation after lipopolysaccharide administration: role of proinflammatory cytokines and neutrophil influx. *J Immunol* 2004; 172:1839-47.
- (347) Pearlman ES, Weber KT, Janicki JS, Pietra GG, Fishman AP. Muscle fiber orientation and connective tissue content in the hypertrophied human heart. *Lab Invest* 1982 ;46:158-64.
- (348) Pelc LR, Gross GJ, Warltier DC. Mechanism of coronary vasodilation produced by bradykinin. *Circulation* 1991;83:2048-56.
- (349) Penna C, Mancardi D, Tullio F, Pagliaro P. Postconditioning and intermittent bradykinin induced cardioprotection require cyclooxygenase activation and prostacyclin release during reperfusion. *Basic Res Cardiol* 2008;103:368-77.
- (350) Penna C, Mancardi D, Raimondo S, Geuna S, Pagliaro P. The paradigm of postconditioning to protect the heart. *Journal of Cellular and Molecular Medicine* 2008;12:435-58.
- (351) Perkins MN, Campbell E, Dray A. Antinociceptive activity of the bradykinin B1 and B2 receptor antagonists, des-Arg9, [Leu8]-BK and HOE 140, in two models of persistent hyperalgesia in the rat. *Pain* 1993;53:191-7.
- (352) Pesquero JB, Araujo RC, Heppenstall PA, Stucky CL, Silva JA Jr., Walther T, Oliveira SM, Pesquero JL, Paiva AC, Calixto JB, Lewin GR, Bader M. Hypoalgesia and altered inflammatory responses in mice lacking kinin B1 receptors. *Proc Natl Acad Sci U S A* 2000; 97:8140-5.
- (353) Phagoo SB, Poole S, Leeb-Lundberg LM. Autoregulation of bradykinin receptors: agonists in the presence of interleukin-1beta shift the repertoire of receptor subtypes from B2 to B1 in human lung fibroblasts. *Mol Pharmacol* 1999;56:325-33.
- (354) Piedimonte G, Nadel JA, Long CS, Hoffman JJ. Neutral endopeptidase in the heart. Neutral endopeptidase inhibition prevents isoproterenol-induced myocardial hypoperfusion in rats by reducing bradykinin degradation. *Circ Res* 1994;75:770-9.
- (355) Pizard A, Blaukat A, Michineau S, Dikic I, Muller-Esterl W, Alhenc-Gelas F, Rajerison RM. Palmitoylation of the human bradykinin B2 receptor influences ligand efficacy. *Biochemistry* 2001;40:15743-51.
- (356) Pizard A, Blaukat A, Muller-Esterl W, Alhenc-Gelas F, Rajerison RM. Bradykinin-induced internalization of the human B2 receptor requires phosphorylation of three serine and two threonine residues at its carboxyl tail. *J Biol Chem* 1999;274:12738-47.
- (357) Pons S, Griol-Charhbil V, Heymes C, Fornes P, Heudes D, Hagege A, Loyer X, Meneton P, Giudicelli JF, Samuel JL, Alhenc-Gelas F, Richer C. Tissue kallikrein deficiency aggravates cardiac remodelling and decreases survival after myocardial infarction in mice. *Eur J Heart Fail* 2008;10:343-51.
- (358) Porsti I, Bara AT, Busse R, Hecker M. Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor. *Br J Pharmacol* 1994 ;111:652-4.
- (359) Powell SJ, Slynn G, Thomas C, Hopkins B, Briggs I, Graham A. Human bradykinin B2 receptor: nucleotide sequence analysis and assignment to chromosome 14. *Genomics* 1993; 15:435-8.
- (360) Prasad A, Husain S, Quyyumi AA. Abnormal flow-mediated epicardial vasomotion in human coronary arteries is improved by angiotensin-converting enzyme inhibition: a potential role of bradykinin. *J Am Coll Cardiol* 1999;33:796-804.
- (361) Pretorius M, Rosenbaum D, Vaughan DE, Brown NJ. Angiotensin-converting enzyme inhibition increases human vascular tissue-type plasminogen activator release through endogenous bradykinin. *Circulation* 2003;107:579-85.
- (362) Pruneau D, Belichard P. Induction of bradykinin B1 receptor-mediated relaxation in the isolated rabbit carotid artery. *Eur J Pharmacol* 1993;239:63-7.
- (363) Puybasset L, Bea ML, Ghaleh B, Giudicelli JF, Berdeaux A. Coronary and systemic hemodynamic effects of sustained inhibition of nitric oxide synthesis in conscious dogs. Evidence for cross talk between nitric oxide and cyclooxygenase in coronary vessels. *Circ Res* 1996;79:343-57.
- (364) Raidoo DM, Ramsaroop R, Naidoo S, Muller-Esterl W, Bhoola KD. Kinin receptors in human vascular tissue: their role in atheromatous disease. *Immunopharmacology* 1997;36:153-60.

-
- (365) Raidoo DM, Sawant S, Mahabeer R, Bhoola KD. Kinin receptors are expressed in human astrocytic tumour cells. *Immunopharmacology* 1999;43:255-63.
- (366) Rauchhaus M, Coats AJ, Anker SD. The endotoxin-lipoprotein hypothesis. *Lancet* 2000; 356:930-3.
- (367) Raut R, Rouleau JL, Blais C, Jr., Gosselin H, Molinaro G, Sirois MG, Lepage Y, Crine P, Adam A. Bradykinin metabolism in the postinfarcted rat heart: role of ACE and neutral endopeptidase 24.11. *Am J Physiol* 1999;276:H1769-H1779.
- (368) Regoli D, Barabe J. Pharmacology of bradykinin and related kinins. *Pharmacol Rev* 1980; 32:1-46.
- (369) Regoli D, Barabe J, Park WK. Receptors for bradykinin in rabbit aortae. *Can J Physiol Pharmacol* 1977;55:855-67.
- (370) Regoli D, Marceau F, Barabe J. De novo formation of vascular receptors for bradykinin. *Can J Physiol Pharmacol* 1978;56:674-7.
- (371) Regoli DC, Marceau F, Lavigne J. Induction of beta 1-receptors for kinins in the rabbit by a bacterial lipopolysaccharide. *Eur J Pharmacol* 1981;71:105-15.
- (372) Reynolds CJ, Togias A, Proud D. Airway neural responses to kinins: tachyphylaxis and role of receptor subtypes. *Am J Respir Crit Care Med* 1999;159:431-8.
- (373) Ribatti D, Presta M, Vacca A, Ria R, Giuliani R, Dell'Era P, Nico B, Roncali L, Dammacco F. Human erythropoietin induces a pro-angiogenic phenotype in cultured endothelial cells and stimulates neovascularization in vivo. *Blood* 1999;93:2627-36.
- (374) Rieder MJ, Carrington DP, Chung M-W. NHLBI Program for Genomic Applications, UW-FHCRC, Seattle SNPs. 2001.
- (375) Risau W. Differentiation of endothelium. *FASEB J* 1995;9:926-33.
- (376) Ritter JM, Doktor HS, Cragoe EJ Jr. Actions of bradykinin and related peptides on rabbit coeliac artery rings. *Br J Pharmacol* 1989;96:23-8.
- (377) Roberts RA, Gullick WJ. Bradykinin receptors undergo ligand-induced desensitization. *Biochemistry* 1990;29:1975-9.
- (378) Rocha E Silva, Rothschild AM. Anaphylatoxin, histamine depletion and skin reactions in the rat. *Nature* 1955;175:987-8.
- (379) Rodriguez JA, De la CP, Collyer E, Decap V, Vio CP, Velarde V. Cyclooxygenase-2 induction by bradykinin in aortic vascular smooth muscle cells. *Am J Physiol Heart Circ Physiol* 2006;290: H30-H36.
- (380) Rosenberg RD, Aird WC. Vascular-bed-specific hemostasis and hypercoagulable states. *N Engl J Med* 1999;340:1555-64.
- (381) Rosenkranz AC, Dusting GJ, Ritchie RH. Hyperglycaemia abolishes the antihypertrophic efficacy of bradykinin in rat ventricular myocytes. *Clin Exp Pharmacol Physiol* 1999;26:519-21.
- (382) Sabourin T, Guay K, Houle S, Bouthillier J, Bachvarov DR, Adam A, Marceau F. Absence of ligand-induced regulation of kinin receptor expression in the rabbit. *Br J Pharmacol* 2001; 133:1154-62.
- (383) Salvino JM, Seoane PR, Douty BD, Awad MM, Dolle RE, Houck WT, Faunce DM, Sawutz DG. Design of potent non-peptide competitive antagonists of the human bradykinin B2 receptor. *J Med Chem* 1993;36:2583-4.
- (384) Sanchez de ML, Neysari S, Jakob S, Petrampil M, Butz N, Banfi A, Zaugg CE, Humar R, Battagay EJ. B2-kinin receptor plays a key role in B1-, angiotensin converting enzyme inhibitor, and vascular endothelial growth factor-stimulated in vitro angiogenesis in the hypoxic mouse heart. *Cardiovasc Res* 2008;80:106-13.
- (385) Santiago JA, Garrison EA, Champion HC, Smith RE, Del Rio O, Kadowitz PJ. Analysis of responses to kallidin, DABK, and DAK in feline hindlimb vascular bed. *Am J Physiol* 1995; 269:H2057-H2064.
- (386) Santos RA, Simoes e Silva AC, Maric C, Silva DM, Machado RP, de Buhr I, Heringer-Walther S,

- Pinheiro SV, Lopes MT, Bader M, Mendes EP, Lemos VS, Campagnole-Santos MJ, Schultheiss HP, Speth R, Walther T. Angiotensin-(1-7) is an endogenous ligand for the G protein-coupled receptor Mas. *Proc Natl Acad Sci U S A* 2003;100:8258-63.
- (387) Sato F, Nagasawa S. Mechanism of kinin release from human low-molecular-mass-kininogen by the synergistic action of human plasma kallikrein and leukocyte elastase. *Biol Chem Hoppe Seyler* 1988;369:1009-17.
- (388) Satterthwaite G, Francis SE, Suvarna K, Blakemore S, Ward C, Wallace D, Braddock M, Crossman D. Differential gene expression in coronary arteries from patients presenting with ischemic heart disease: Further evidence for the inflammatory basis of atherosclerosis. *American Heart Journal* 2005;150:488-99.
- (389) Schafer S, Schmidts HL, Bleich M, Busch AE, Linz W. Nephroprotection in Zucker diabetic fatty rats by vasopeptidase inhibition is partly bradykinin B2 receptor dependent. *Br J Pharmacol* 2004;143:27-32.
- (390) Schanstra JP, Alric C, Marin-Castano ME, Girolami JP, Bascands JL. Renal bradykinin receptors: localisation, transduction pathways and molecular basis for a possible pathological role (review). *Int J Mol Med* 1999;3:185-91.
- (391) Schanstra JP, Bataille E, Marin Castano ME, Barascud Y, Hirtz C, Pesquero JB, Pecher C, Gauthier F, Girolami JP, Bascands JL. The B1-agonist [des-Arg10]-kallidin activates transcription factor NF-kappaB and induces homologous upregulation of the bradykinin B1-receptor in cultured human lung fibroblasts. *J Clin Invest* 1998;101:2080-91.
- (392) Schanstra JP, Duchene J, Desmond L, Neau E, Calise D, Estaque S, Girolami JP, Bascands JL. The protective effect of angiotensin converting enzyme inhibition in experimental renal fibrosis in mice is not mediated by bradykinin B2 receptor activation. *Thromb Haemost* 2003; 89:735-40.
- (393) Schanstra JP, Duchene J, Praddaude F, Bruneval P, Tack I, Chevalier J, Girolami JP, Bascands JL. Decreased renal NO excretion and reduced glomerular tuft area in mice lacking the bradykinin B2 receptor. *Am J Physiol Heart Circ Physiol* 2003;284:H1904-H1908.
- (394) Schanstra JP, Marin-Castano ME, Praddaude F, Tack I, Ader JL, Girolami JP, Bascands JL, Jeunier B. Bradykinin B(1) receptor-mediated changes in renal hemodynamics during endotoxin-induced inflammation. *J Am Soc Nephrol* 2000;11:1208-15.
- (395) Schanstra JP, Neau E, Drogoz P, Arevalo Gomez MA, Lopez Novoa JM, Calise D, Pecher C, Bader M, Girolami JP, Bascands JL. In vivo bradykinin B2 receptor activation reduces renal fibrosis. *J Clin Invest* 2002;110:371-9.
- (396) Scharlt M, Bocksch WG, Dreyse S, Beckmann S, Franke O, Hunten U. Remodeling of myocardium and arteries by chronic angiotensin converting enzyme inhibition in hypertensive patients. *J Hypertens Suppl* 1994;12:S37-S42.
- (397) Schini VB, Boulanger C, Regoli D, Vanhoutte PM. Bradykinin stimulates the production of cyclic GMP via activation of B2 kinin receptors in cultured porcine aortic endothelial cells. *J Pharmacol Exp Ther* 1990;252:581-5.
- (398) Schlemper V, Medeiros R, Ferreira J, Campos MM, Calixto JB. Mechanisms underlying the relaxation response induced by bradykinin in the epithelium-intact guinea-pig trachea in vitro. *Br J Pharmacol* 2005;145:740-50.
- (399) Schmaier AH, Kuo A, Lundberg D, Murray S, Cines DB. The expression of high molecular weight kininogen on human umbilical vein endothelial cells. *J Biol Chem* 1988 ;263:16327-33.
- (400) Schmedtje JF Jr., Ji YS, Liu WL, DuBois RN, Runge MS. Hypoxia induces cyclooxygenase-2 via the NF-kappaB p65 transcription factor in human vascular endothelial cells. *J Biol Chem* 1997;272:601-8.
- (401) Schmidlin F, Scherrer D, Landry Y, Gies JP. Glucocorticoids inhibit the bradykinin B2 receptor increase induced by interleukin-1beta in human bronchial smooth muscle cells. *Eur J Pharmacol* 1998;354:R7-R8.
- (402) Scholkens BA, Linz W, Konig W. Effects of the angiotensin converting enzyme inhibitor, ramipril, in isolated ischaemic rat heart are abolished by a bradykinin antagonist. *J Hypertens Suppl* 1988;6:S25-S28.

-
- (403) Schremmer-Danninger E, Hermann A, Fink E, Fritz H, Roscher AA. Identification and occurrence of mRNAs for components of the kallikrein-kinin system in human skin and in skin diseases. *Immunopharmacology* 1999;43:287-91.
- (404) Searles CD, Harrison DG. The interaction of nitric oxide, bradykinin, and the angiotensin II type 2 receptor: lessons learned from transgenic mice. *J Clin Invest* 1999;104:1013-4.
- (405) Seccia TM, Belloni AS, Guidolin D, Sticchi D, Nussdorfer GG, Pessina AC, Rossi GP. The renal antifibrotic effects of angiotensin-converting enzyme inhibition involve bradykinin B2 receptor activation in angiotensin II-dependent hypertension. *J Hypertens* 2006;24:1419-27.
- (406) Seguin T, Buleon M, Destrube M, Ranera MT, Couture R, Girolami JP, Tack I. Hemodynamic and renal involvement of B(1) and B(2) kinin receptors during the acute phase of endotoxin shock in mice. *Int Immunopharmacol* 2008;8:217-21.
- (407) Shariat-Madar Z, Mahdi F, Warnock M, Homeister JW, Srikanth S, Krijanovski Y, Murphey LJ, Jaffa AA, Schmaier AH. Bradykinin B2 receptor knockout mice are protected from thrombosis by increased nitric oxide and prostacyclin. *Blood* 2006;108:192-9.
- (408) Shiga N, Hirano K, Hirano M, Nishimura J, Nawata H, Kanaide H. Long-term inhibition of RhoA attenuates vascular contractility by enhancing endothelial NO production in an intact rabbit mesenteric artery. *Circ Res* 2005;96:1014-21.
- (409) Shimokawa H, Flavahan NA, Lorenz RR, Vanhoutte PM. Prostacyclin releases endothelium-derived relaxing factor and potentiates its action in coronary arteries of the pig. *Br J Pharmacol* 1988;95:1197-203.
- (410) Shintani Y, Node K, Asanuma H, Sanada S, Takashima S, Asano Y, Liao Y, Fujita M, Hirata A, Shinozaki Y, Fukushima T, Nagamachi Y, Okuda H, Kim J, Tomoike H, Hori M, Kitakaze M. Opening of Ca²⁺-activated K⁺ channels is involved in ischemic preconditioning in canine hearts. *Journal of Molecular and Cellular Cardiology* 2004 ;37:1213-8.
- (411) Shiuchi T, Cui TX, Wu L, Nakagami H, Takeda-Matsubara Y, Iwai M, Horiuchi M. ACE inhibitor improves insulin resistance in diabetic mouse via bradykinin and NO. *Hypertension* 2002;40:329-34.
- (412) Shreeniwas R, Koga S, Karakurum M, Pinsky D, Kaiser E, Brett J, Wolitzky BA, Norton C, Plocinski J, Benjamin W. . Hypoxia-mediated induction of endothelial cell interleukin-1 alpha. An autocrine mechanism promoting expression of leukocyte adhesion molecules on the vessel surface. *J Clin Invest* 1992;90:2333-9.
- (413) Silva J, Araujo RC, Baltatu O, Oliveira SM, Tschöpe C, Fink E, Hoffmann S, Plehm R, Chai KX, Chao L, Chao J, Ganten D, Pesquero JB, Bader M. Reduced cardiac hypertrophy and altered blood pressure control in transgenic rats with the human tissue kallikrein gene. *FASEB J* 2000 8;99-1010.
- (414) Silverberg DS, Wexler D, Sheps D, Blum M, Keren G, Baruch R, Schwartz D, Yachnin T, Steinbruch S, Shapira I, Laniado S, Iaina A. The effect of correction of mild anemia in severe, resistant congestive heart failure using subcutaneous erythropoietin and intravenous iron: a randomized controlled study. *J Am Coll Cardiol* 2001;37:1775-80.
- (415) Silvestre JS, Bergaya S, Tamarat R, Duriez M, Boulanger CM, Levy BI. Proangiogenic effect of angiotensin-converting enzyme inhibition is mediated by the bradykinin B(2) receptor pathway. *Circ Res* 2001;89:678-83.
- (416) Simmet T, Hertting G. On the relation between contraction and prostaglandin release in rabbit mesenteric blood vessels. *Eur J Pharmacol* 1980;65:325-31.
- (417) Simmons DL, Botting RM, Hla T. Cyclooxygenase isozymes: the biology of prostaglandin synthesis and inhibition. *Pharmacol Rev* 2004;56:387-437.
- (418) Siren AL, Fratelli M, Brines M, Goemans C, Casagrande S, Lewczuk P, Keenan S, Gleiter C, Pasquali C, Capobianco A, Mennini T, Heumann R, Cerami A, Ehrenreich H, Ghezzi P. Erythropoietin prevents neuronal apoptosis after cerebral ischemia and metabolic stress. *Proc Natl Acad Sci U S A* 2001;98:4044-9.
- (419) Smith JA, Webb C, Holford J, Burgess GM. Signal transduction pathways for B1 and B2 bradykinin receptors in bovine pulmonary artery endothelial cells. *Mol Pharmacol* 1995; 47:525-34.

- (420) Solomon SD, Dobson J, Pocock S, Skali H, McMurray JJ, Granger CB, Yusuf S, Swedberg K, Young JB, Michelson EL, Pfeffer MA. Influence of nonfatal hospitalization for heart failure on subsequent mortality in patients with chronic heart failure. *Circulation* 2007;116:1482-7.
- (421) Solomon SD, McMurray JJ, Pfeffer MA, Wittes J, Fowler R, Finn P, Anderson WF, Zauber A, Hawk E, Bertagnolli M. Cardiovascular risk associated with celecoxib in a clinical trial for colorectal adenoma prevention. *N Engl J Med* 2005;352:1071-80.
- (422) Sonnenblick M, Rosenmann D, Rosin A. Reversible cardiomyopathy induced by interferon. *BMJ* 1990;300:1174-5.
- (423) Sordella R, Jiang W, Chen GC, Curto M, Settleman J. Modulation of Rho GTPase signaling regulates a switch between adipogenesis and myogenesis. *Cell* 2003;113:147-58.
- (424) Soskic V, Nyakatura E, Roos M, Muller-Esterl W, Godovac-Zimmermann J. Correlations in palmitoylation and multiple phosphorylation of rat bradykinin B2 receptor in Chinese hamster ovary cells. *J Biol Chem* 1999;274:8539-45.
- (425) Staszewska-Barczak J, Dusting GJ. Sympathetic cardiovascular reflex initiated by bradykinin-induced stimulation of cardiac pain receptors in the dog. *Clin Exp Pharmacol Physiol* 1977;4:443-52.
- (426) Steranka LR, Manning DC, Dehaas CJ, Ferkany JW, Borosky SA, Connor JR, Vavrek RJ, Stewart JM, Snyder SH. Bradykinin as a pain mediator: receptors are localized to sensory neurons, and antagonists have analgesic actions. *Proc Natl Acad Sci U S A* 1988;85:3245-9.
- (427) Sun D, Huang A, Smith CJ, Stackpole CJ, Connetta JA, Shesely EG, Koller A, Kaley G. Enhanced release of prostaglandins contributes to flow-induced arteriolar dilation in eNOS knockout mice. *Circ Res* 1999;85:288-93.
- (428) Suo M, Hautala N, Foldes G, Szokodi I, Toth M, Leskinen H, Uusimaa P, Vuolteenaho O, Nemer M, Ruskoaho H. Posttranscriptional control of BNP gene expression in angiotensin II-induced hypertension. *Hypertension* 2002;39:803-8.
- (429) Swedberg K. Importance of neuroendocrine activation in chronic heart failure. Impact on treatment strategies. *European Journal of Heart Failure* 2000;2:229-33.
- (430) Szokodi I, Kinnunen P, Tavi P, Weckstrom M, Toth M, Ruskoaho H. Evidence for cAMP-independent mechanisms mediating the effects of adrenomedullin, a new inotropic peptide. *Circulation* 1998;97:1062-70.
- (431) Taddei S, Ghiadoni L, Virdis A, Buralli S, Salvetti A. Vasodilation to bradykinin is mediated by an ouabain-sensitive pathway as a compensatory mechanism for impaired nitric oxide availability in essential hypertensive patients. *Circulation* 1999;100:1400-5.
- (432) Taddei S, Virdis A, Mattei P, Ghiadoni L, Gennari A, Fasolo CB, Sudano I, Salvetti A. Aging and endothelial function in normotensive subjects and patients with essential hypertension. *Circulation* 1995;91:1981-7.
- (433) Takagaki Y, Kitamura N, Nakanishi S. Cloning and sequence analysis of cDNAs for human high molecular weight and low molecular weight prekininogens. Primary structures of two human prekininogens. *J Biol Chem* 1985;260:8601-9.
- (434) Takemoto M, Sun J, Hiroki J, Shimokawa H, Liao JK. Rho-kinase mediates hypoxia-induced downregulation of endothelial nitric oxide synthase. *Circulation* 2002;106:57-62.
- (435) Tan Y, Wang B, Keum JS, Jaffa AA. Mechanisms through which bradykinin promotes glomerular injury in diabetes. *Am J Physiol Renal Physiol* 2005;288:F483-F492.
- (436) Tan Y, Keum JS, Wang B, McHenry MB, Lipsitz SR, Jaffa AA. Targeted deletion of B2-kinin receptors protects against the development of diabetic nephropathy. *Am J Physiol Renal Physiol* 2007;293:F1026-F1035.
- (437) Tanaka Y, Nagai M, Date T, Okada T, Abe Y, Seki S, Taniguchi M, Taniguchi I, Mochizuki S. Effects of bradykinin on cardiovascular remodeling in renovascular hypertensive rats. *Hypertens Res* 2004;27:865-75.
- (438) Taub JS, Guo R, Leeb-Lundberg LM, Madden JF, Daaka Y. Bradykinin receptor subtype 1 expression and function in prostate cancer. *Cancer Res* 2003;63:2037-41.

-
- (439) Tavazzi L, Tognoni G, Franzosi MG, Latini R, Maggioni AP, Marchioli R, Nicolosi GL, Porcu M. Rationale and design of the GISSI heart failure trial: a large trial to assess the effects of n-3 polyunsaturated fatty acids and rosuvastatin in symptomatic congestive heart failure. *Eur J Heart Fail* 2004;6:635-41.
- (440) Teerlink JR, Clozel M, Fischli W, Clozel JP. Temporal evolution of endothelial dysfunction in a rat model of chronic heart failure. *J Am Coll Cardiol* 1993;22:615-20.
- (441) Thollon C, Bidouard JP, Cambarrat C, Delescluse I, Villeneuve N, Vanhoutte PM, Vilaine JP. Alteration of endothelium-dependent hyperpolarizations in porcine coronary arteries with regenerated endothelium. *Circ Res* 1999;84:371-7.
- (442) Thomas PB, Liu EC, Webb ML, Mukherjee R, Hebbar L, Spinale FG. Exogenous effects and endogenous production of endothelin in cardiac myocytes: potential significance in heart failure. *Am J Physiol* 1996;271:H2629-H2637.
- (443) Thuringer D, Maulon L, Frelin C. Rapid transactivation of the vascular endothelial growth factor receptor KDR/Flk-1 by the bradykinin B2 receptor contributes to endothelial nitric-oxide synthase activation in cardiac capillary endothelial cells. *J Biol Chem* 2002;277:2028-32.
- (444) Tio RA, van Gilst WH, Rett K, Wolters K, Dietze GJ, Wesseling H. The effects of bradykinin and the ischemic isolated rat heart. *Horm Metab Res Suppl* 1990;22:85-9.
- (445) Toda N. Actions of bradykinin on isolated cerebral and peripheral arteries. *Am J Physiol* 1977;232:H267-H274.
- (446) Tom B, de Vries R, Saxena PR, Danser AH. Bradykinin potentiation by angiotensin-(1-7) and ACE inhibitors correlates with ACE C- and N-domain blockade. *Hypertension* 2001;38:95-9.
- (447) Tondangu D, Hittinger L, Ghaleh B, Le Corvoisier P, Sambin L, Champagne S, Badoual T, Vincent F, Berdeaux A, Crozatier B, Su JB. Chronic infusion of bradykinin delays the progression of heart failure and preserves vascular endothelium-mediated vasodilation in conscious dogs. *Circulation* 2004;109:114-9.
- (448) Trabold F, Pons S, Hagege AA, Bloch-Faure M, Alhenc-Gelas F, Giudicelli JF, Richer-Giudicelli C, Meneton P. Cardiovascular phenotypes of kinin B2 receptor- and tissue kallikrein-deficient mice. *Hypertension* 2002;40:90-5.
- (449) Treasure CB, Vita JA, Cox DA, Fish RD, Gordon JB, Mudge GH, Colucci WS, Sutton MG, Selwyn AP, Alexander RW, et al. Endothelium-dependent dilation of the coronary microvasculature is impaired in dilated cardiomyopathy. *Circulation* 1990;81:772-9.
- (450) Trevisani M, Amadesi S, Schmidlin F, Poblete MT, Bardella E, Maggiore B, Harrison S, Figueroa CD, Tognetto M, Navarra G, Turini A, Bunnett NW, Geppetti P, De Giorgio R. Bradykinin B2 receptors mediate contraction in the normal and inflamed human gallbladder in vitro. *Gastroenterology* 2003;125:126-35.
- (451) Trochu JN, Mital S, Zhang X, Xu X, Ochoa M, Liao JK, Recchia FA, Hintze TH. Preservation of NO production by statins in the treatment of heart failure. *Cardiovasc Res* 2003;60:250-8.
- (452) Tschope C, Heringer-Walther S, Koch M, Spillmann F, Wendorf M, Leitner E, Schultheiss HP, Walther T. Upregulation of bradykinin B1-receptor expression after myocardial infarction. *Br J Pharmacol* 2000;129:1537-8.
- (453) Tschope C, Seidl U, Reinecke A, Riester U, Graf K, Schultheiss HP, Hilgenfeldt U, Unger T. Kinins are involved in the antiproteinuric effect of angiotensin-converting enzyme inhibition in experimental diabetic nephropathy. *Int Immunopharmacol* 2003;3:335-44.
- (454) Tschope C, Walther T, Koniger J, Spillmann F, Westermann D, Escher F, Pauschinger M, Pesquero JB, Bader M, Schultheiss HP, Noutsias M. Prevention of cardiac fibrosis and left ventricular dysfunction in diabetic cardiomyopathy in rats by transgenic expression of the human tissue kallikrein gene. *FASEB J* 2004;18:828-35.
- (455) Tsukagoshi H, Shimizu Y, Horie T, Fukabori Y, Shimizu Y, Iwamae S, Hisada T, Ishizuka T, Izuka K, Dobashi K, Mori M. Regulation by interleukin-1beta of gene expression of bradykinin B1 receptor in MH-S murine alveolar macrophage cell line. *Biochem Biophys Res Commun* 1999;259:476-82.
- (456) Tsutamoto T, Hisanaga T, Wada A, Maeda K, Ohnishi M, Fukai D, Mabuchi N, Sawaki M,

- Kinoshita M. Interleukin-6 spillover in the peripheral circulation increases with the severity of heart failure, and the high plasma level of interleukin-6 is an important prognostic predictor in patients with congestive heart failure. *J Am Coll Cardiol* 1998;31:391-8.
- (457) Tsutsui M, Onoue H, Iida Y, Smith L, O'Brien T, Katusic ZS. B(1) and B(2) bradykinin receptors on adventitial fibroblasts of cerebral arteries are coupled to recombinant eNOS. *Am J Physiol Heart Circ Physiol* 2000;278:H367-H372.
- (458) Tsutsumi Y, Matsubara H, Masaki H, Kurihara H, Murasawa S, Takai S, Miyazaki M, Nozawa Y, Ozono R, Nakagawa K, Miwa T, Kawada N, Mori Y, Shibasaki Y, Tanaka Y, Fujiyama S, Koyama Y, Fujiyama A, Takahashi H, Iwasaka T. Angiotensin II type 2 receptor overexpression activates the vascular kinin system and causes vasodilation. *J Clin Invest* 1999;104:925-35.
- (459) Ueda S, Masumori-Maemoto S, Ashino K, Nagahara T, Gotoh E, Umemura S, Ishii M. Angiotensin-(1-7) Attenuates Vasoconstriction Evoked by Angiotensin II but Not by Noradrenaline in Man. *Hypertension* 2000;35:998-1001.
- (460) Uehata M, Ishizaki T, Satoh H, Ono T, Kawahara T, Morishita T, Tamakawa H, Yamagami K, Inui J, Maekawa M, Narumiya S. Calcium sensitization of smooth muscle mediated by a Rho-associated protein kinase in hypertension. *Nature* 1997;389:990-4.
- (461) Urakami-Harasawa L, Shimokawa H, Nakashima M, Egashira K, Takeshita A. Importance of endothelium-derived hyperpolarizing factor in human arteries. *J Clin Invest* 1997;100:2793-9.
- (462) Uruno A, Sugawara A, Kanatsuka H, Kagechika H, Saito A, Sato K, Kudo M, Takeuchi K, Ito S. Upregulation of nitric oxide production in vascular endothelial cells by all-trans retinoic acid through the phosphoinositide 3-kinase/Akt pathway. *Circulation* 2005;112:727-36.
- (463) van der Meer P, Voors AA, Lipsic E, van Gilst WH, Van Veldhuisen DJ. Erythropoietin in cardiovascular diseases. *Eur Heart J* 2004;25:285-91.
- (464) van Aelst L, Souza-Schorey C. Rho GTPases and signaling networks. *Genes Dev* 1997;11:2295-322.
- (465) van Iwaarden F, de Groot PG, Bouma BN. The binding of high molecular weight kininogen to cultured human endothelial cells. *J Biol Chem* 1988;263:4698-703.
- (466) Vanhoutte PM, Boulanger CM. Endothelium-dependent responses in hypertension. *Hypertens Res* 1995;18:87-98.
- (467) Vavrek RJ, Stewart JM. Competitive antagonists of bradykinin. *Peptides* 1985;6:161-4.
- (468) Vegh A, Papp JG, Parratt J. Attenuation of the antiarrhythmic effects of ischaemic preconditioning by blockade of bradykinin B2 receptors. *Br J Pharmacol* 1994;113:1167-72.
- (469) Venema RC. Post-translational mechanisms of endothelial nitric oxide synthase regulation by bradykinin. *Int Immunopharmacol* 2002;2:1755-62.
- (470) Vianna RM, Ongali B, Regoli D, Calixto JB, Couture R. Up-regulation of kinin B1 receptor in the lung of streptozotocin-diabetic rat: autoradiographic and functional evidence. *Br J Pharmacol* 2003;138:13-22.
- (471) Vickers C, Hales P, Kaushik V, Dick L, Gavin J, Tang J, Godbout K, Parsons T, Baronas E, Hsieh F, Acton S, Patane M, Nichols A, Tummino P. Hydrolysis of biological peptides by human angiotensin-converting enzyme-related carboxypeptidase. *J Biol Chem* 2002; 277:14838-43.
- (472) Vidal F, Colome C, Martinez-Gonzalez J, Badimon L. Atherogenic concentrations of native low-density lipoproteins down-regulate nitric-oxide-synthase mRNA and protein levels in endothelial cells. *Eur J Biochem* 1998;252:378-84.
- (473) Villa LM, Salas E, Darley-USmar VM, Radomski MW, Moncada S. Peroxynitrite induces both vasodilatation and impaired vascular relaxation in the isolated perfused rat heart. *Proc Natl Acad Sci U S A* 1994;91:12383-7.
- (474) Vogel R, Werle E, Zickgraf-Rudel G. Current aspects of kinin research. I. Potentiation and blocking of biological kinin activity. *Z Klin Chem Klin Biochem* 1970;8:177-85.
- (475) Wall TM, Sheehy R, Hartman JC. Role of bradykinin in myocardial preconditioning. *J Pharmacol Exp Ther* 1994;270:681-9.
- (476) Waltenberger J, Mayr U, Pentz S, Hombach V. Functional upregulation of the vascular

- endothelial growth factor receptor KDR by hypoxia. *Circulation* 1996;94:1647-54.
- (477) Walther T, Siems WE, Hauke D, Spillmann F, Dendorfer A, Krause W, Schultheiss HP, Tschöpe C. AT1 receptor blockade increases cardiac bradykinin via neutral endopeptidase after induction of myocardial infarction in rats. *FASEB J* 2002;16:1237-41.
- (478) Wang B, Dang A, Liu G. Genetic variation in the promoter region of the β_2 bradykinin receptor gene is associated with essential hypertension in a chinese han population. *Hypertension Research* 2001;24:299-302.
- (479) Watson PJ. Drug receptor sites in the rabbit saphenous nerve. *Br J Pharmacol* 1970;40:102-12.
- (480) Werle E, Götze W. Über die Wirkung des Kallikreins auf den isolierten Darm und über eine neue darmkontrahierende Substanz. *Biochem Z.* 1937;289:217-33.
- (481) Werle E, Grunz M. Zur Kenntnis der darm kontrahierenden, uterus erregenden und blutdrucksenkenden substanz DK. *Biochem Z.* 1939;301:429-36.
- (482) Widlansky ME, Price DT, Gokce N, Eberhardt RT, Duffy SJ, Holbrook M, Maxwell C, Palmisano J, Keaney JF Jr., Morrow JD, Vita JA. Short- and long-term COX-2 inhibition reverses endothelial dysfunction in patients with hypertension. *Hypertension* 2003;42:310-5.
- (483) Wiemer G, Scholkens BA, Becker RH, Busse R. Ramiprilat enhances endothelial autacoid formation by inhibiting breakdown of endothelium-derived bradykinin. *Hypertension* 1991;18:558-63.
- (484) Wigg SJ, Tare M, Tonta MA, O'Brien RC, Meredith IT, Parkington HC. Comparison of effects of diabetes mellitus on an EDHF-dependent and an EDHF-independent artery. *Am J Physiol Heart Circ Physiol* 2001;281:H232-H240.
- (485) Wilhelmssen L, Rosengren A, Eriksson H, Lappas G. Heart failure in the general population of men - morbidity, risk factors and prognosis. *Journal of Internal Medicine* 2001;249:253-61.
- (486) Windischhofer W, Leis HJ. [3H]bradykinin receptor-binding, receptor-recycling, and receptor-internalization of the B2 bradykinin receptor in the murine osteoblast-like cell line MC3T3-E1. *J Bone Miner Res* 1997;12:1615-25.
- (487) Witherow FN, Helmy A, Webb DJ, Fox KA, Newby DE. Bradykinin contributes to the vasodilator effects of chronic angiotensin-converting enzyme inhibition in patients with heart failure. *Circulation* 2001;104:2177-81.
- (488) Wolfrum S, Dendorfer A, Dominiak P. Identification of kallidin degrading enzymes in the isolated perfused rat heart. *Jpn J Pharmacol* 1999;79:117-20.
- (489) Wolfrum S, Richardt G, Dominiak P, Katus HA, Dendorfer A. Apstatin, a selective inhibitor of aminopeptidase P, reduces myocardial infarct size by a kinin-dependent pathway. *Br J Pharmacol* 2001;134:370-4.
- (490) Wollert KC, Drexler H. The renin-angiotensin system and experimental heart failure. *Cardiovasc Res* 1999;43:838-49.
- (491) Wu J, Akaike T, Hayashida K, Miyamoto Y, Nakagawa T, Miyakawa K, Muller-Esterl W, Maeda H. Identification of bradykinin receptors in clinical cancer specimens and murine tumor tissues. *Int J Cancer* 2002;98:29-35.
- (492) Xie W, Herschman HR. Transcriptional regulation of prostaglandin synthase 2 gene expression by platelet-derived growth factor and serum. *J Biol Chem* 1996;271:31742-8.
- (493) Xu J, Carretero OA, Sun Y, Shesely EG, Rhaleb NE, Liu YH, Liao TD, Yang JJ, Bader M, Yang XP. Role of the B1 kinin receptor in the regulation of cardiac function and remodeling after myocardial infarction. *Hypertension* 2005;45:747-53.
- (494) Yamada H, Fabris B, Allen AM, Jackson B, Johnston CI, Mendelsohn AO. Localization of angiotensin converting enzyme in rat heart. *Circ Res* 1991;68:141-9.
- (495) Yamamoto K, Burnett JC Jr., Jougasaki M, Nishimura RA, Bailey KR, Saito Y, Nakao K, Redfield MM. Superiority of brain natriuretic peptide as a hormonal marker of ventricular systolic and diastolic dysfunction and ventricular hypertrophy. *Hypertension* 1996;28:988-94.
- (496) Yamasaki S, Sawada S, Komatsu S, Kawahara T, Tsuda Y, Sato T, Toratani A, Kono Y, Higaki T, Imamura H, Tada Y, Akamatsu N, Tamagaki T, Tsuji H, Nakagawa M. Effects of bradykinin on

- prostaglandin I2 synthesis in human vascular endothelial cells. *Hypertension* 2000;36:201-7.
- (497) Yang CM, Kandaswamy V, Young D, Sen S. Changes in collagen phenotypes during progression and regression of cardiac hypertrophy. *Cardiovasc Res* 1997;36:236-45.
- (498) Yang HY, Erdos EG, Levin Y. A dipeptidyl carboxypeptidase that converts angiotensin I and inactivates bradykinin. *Biochim Biophys Acta* 1970;214:374-6.
- (499) Yang X, Taylor L, Polgar P. p53 down-regulates human bradykinin B1 receptor gene expression. *J Cell Biochem* 2001;82:38-45.
- (500) Yang XP, Liu YH, Mehta D, Cavaşin MA, Shesely E, Xu J, Liu F, Carretero OA. Diminished cardioprotective response to inhibition of angiotensin-converting enzyme and angiotensin II type 1 receptor in B(2) kinin receptor gene knockout mice. *Circ Res* 2001;88:1072-9.
- (501) Yang XP, Liu YH, Scicli GM, Webb CR, Carretero OA. Role of Kinins in the cardioprotective effect of preconditioning : study of myocardial ischemia/reperfusion injury in B2 kinin receptor knockout mice and kininogen-deficient rats. *Hypertension* 1997;30:735-40.
- (502) Yaqoob M, Snell CR, Burgess GM. Carbohydrate analysis of the B2 bradykinin receptor from rat uterus. *J Neurochem* 1995;65:1290-6.
- (503) Yokota K, Kishida M, Ogura T, Suzuki J, Otsuka F, Mimura Y, Takeda M, Nakamura Y, Makino H. Role of bradykinin in renoprotective effects by angiotensin II type 1 receptor antagonist in salt-sensitive hypertension. *Hypertens Res* 2003;26:265-72.
- (504) Yoshida H, Kusama Y, Kodani E, Yasutake M, Takano H, Atarashi H, Kishida H, Takano T. Pharmacological preconditioning with bradykinin affords myocardial protection through NO-dependent mechanisms. *Int Heart J* 2005;46:877-87.
- (505) Yoshida H, Zhang JJ, Chao L, Chao J. Kallikrein gene delivery attenuates myocardial infarction and apoptosis after myocardial ischemia and reperfusion. *Hypertension* 2000;35:25-31.
- (506) Yoshimura M, Yasue H, Okumura K, Ogawa H, Jougasaki M, Mukoyama M, Nakao K, Imura H. Different secretion patterns of atrial natriuretic peptide and brain natriuretic peptide in patients with congestive heart failure. *Circulation* 1993;87:464-9.
- (507) Zhang Y, Adner M, Cardell LO. IL-1beta-induced transcriptional up-regulation of bradykinin B1 and B2 receptors in murine airways. *Am J Respir Cell Mol Biol* 2007;36:697-705.
- (508) Zhou X, Polgar P, Taylor L. Roles for interleukin-1beta, phorbol ester and a post-transcriptional regulator in the control of bradykinin B1 receptor gene expression. *Biochem J* 1998;330:361-6.
- (509) Zhuo JL, Mendelsohn FA, Ohishi M. Perindopril alters vascular angiotensin-converting enzyme, AT(1) receptor, and nitric oxide synthase expression in patients with coronary heart disease. *Hypertension* 2002;39:634-8.
- (510) Zini JM, Schmaier AH, Cines DB. Bradykinin regulates the expression of kininogen binding sites on endothelial cells. *Blood* 1993;81:2936-46.
- (511) Zisman LS, Keller RS, Weaver B, Lin Q, Speth R, Bristow MR, Canver CC. Increased angiotensin-(1-7)-forming activity in failing human heart ventricles: evidence for upregulation of the angiotensin-converting enzyme Homologue ACE2. *Circulation* 2003;108:1707-12.
- (512) Zuccollo A, Navarro M, Catanzaro O. Effects of B1 and B2 kinin receptor antagonists in diabetic mice. *Can J Physiol Pharmacol* 1996;74:586-9.
- (513) Zuccollo A, Navarro M, Frontera M, Cueva F, Carattino M, Catanzaro OL. The involvement of kallikrein-kinin system in diabetes type I (insulinitis). *Immunopharmacology* 1999;45:69-74.